# SOME ASPECTS ON REPRODUCTIVE BIOLOGY OF *GERRES FILAMENTOSUS* CUVIER

DISSERTATION SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF

## **MASTER OF FISHERIES SCIENCE**

# IN MARICULTURE OF THE CENTRAL INSTITUTE OF FISHERIES EDUCATION (DEEMED UNIVERSITY) VERSOVA, MUMBAI - 409 061

BY

### A. RAIIIANATIIU BEEVI



L'May r	As Cantral &	erina Pasteries	
hen	wich Institute	. Cechin	
Bate of sec	:ip1	7. 2000	
Accession M	h D-2	36	
Cherry Dr.	a H	74 K	° /1 /

INDIAN COUNCIL OF AGRICULTURAL RESEARCH CENTRAL MARINE FISHERIES RESEARCH INSTITUTE P.B. NO. 1603, KOCHI - 682 014 INDIA

JULY 1998

# IN THE NAME OF ALLAH, THE MOST MERCIFUL AND GOMPASSIONATE

# **DECLARATION**

I hereby declare that this thesis entitled "Some Aspects on reproductive biology of *Gerres Filamentosus* Cuvier" is based on my research and has not previously formed the basis for the award of any degree, diploma, associateship, fellowship or other similar titles or recognition.

fartanathalut

A. RAIHANATHU BEEVI

Kochi - 682 014 July, 1998

# CERTIFICATE

Certified that the dissertation entitled "Some Aspects on reproductive biology of *Gerres filamentosus* Cuvier" is a bonafide record of work done by Kum. **RAIHANATHU BEEVI A.** under our guidance at the Central Marine Fisheries Research Institute during the tenure of her M.Fsc (Mariculture) programme of 1996-98 and that it has not previously formed the basis for the award of any other degree, deploma or other similar titles or for any publication.

Dr.K.K. JOSH I Scientist C.M.F.R.I.,Kochi Chairman and Major Advisor, Advisory Committee

Grace

Smt. Grace Mathew Senior Scientist, C.M.F.R.I, Kochi. Co-Chairman, Advisory Committee

Dr. N. Gopinatha Menon Senior Scientist, C.M.F.R.I, Kochi. Member, Advisory Committee

# CONTENTS

PAGE NO.

DECLARATION	
CERTIFICATE1	
ABSTRACT IN HINDI	
ABSTRACT IN ENGLISH	1.
ACKNOWLEDGEMENT <b>S</b>	2
INTRODUCTION	4
REVIEW OF LITERATURE	7
MATERIALS AND METHODS	16
RESULTS	19
DISCUSSION	30
SUMMARY	39
REFERENCES	40

# LIST OF TABLES

Table No. 1	fecundity of G. filamentosus, maturity stage III
Table No. 2	fecundity of G. filamentosus, maturity stage IV
Table No. 3	fecundity of G. filamentosus, maturity stage V
Table No.4	Maturity percentage of Female G. filamentosus
Table No.5	Maturity percentage of Male G. Filamentosus
Table No. 6	Sex ratio of G. Filamentosus

# LIST OF FIGURES

1.	a) G. filmentosus cuvier female
	b) G. filamentosus female showing ovary
2.	a) G. filmentosus maturity stages I to V
	b) G. filmentosus maturity stage II
	c) G. filmentosus maturity stage V
3.	a) G. filmentosus stage III oocyte at yolk precursor stage
	b) Stage IV early non staining yolk stage.
4	a) G. filamentosus stage IV
	b) Stage V Early yolk staining stage
5	a) G. filmentosus stage V
	b) G. <i>filmentosus</i> stage V
6	a) Fecundity of G. filmentosus in relation to length, stage III
7	Fecundity of G. filmentosus in relation to length, stage IV
8	Fecundity of G. filmentosus in relation to length, stage V
9	Size at first maturity of G. filmentosus

10	Length frequency distribution of G. filmentosus 4-3-98
11	Length frequency distribution of G. filmentosus 6-3-98
12	Length frequency distribution of G. filmentosus 11-3-98
13	Length frequency distribution of G. $F$ ilmentosus 19-3-98
14	Length frequency distribution of G. filmentosus 25-3-98
15	Length frequency distribution of G. filmentosus 1-4-98
16	Length frequency distribution of G. filmentosus 22-4-98
17	Oocyte diameter frequency graph of different stages of ovary 6-3-98
18	Oocyte diameter frequency graph of different stages of ovary 19-3-98
19	Oocyte diameter frequency graph of different stages of ovary 25-3-98
20	Oocyte diameter frequency graph of different stages of ovary 1-4-98

## ABSTRACT

#### STUDIES ON SOME ASPECTS OF REPRODUCTIVE BIOLOGY OF

#### **GERRES FILAMENTOSUS CUVIER IN COCHIN**

The reproductive biology of *Gerres filamentosus* cuvier was investigated in cochin from the samples collected from the trawl landings. Analysis of reproductive organs, maturity condition, fecundity and sex ratio was done during March - June 1998. Oocyte diameter was measured and histological characteristics of ovary were studied. This species remains in the estuarine environment and undergo migration to nearby coastal areas for the purpose of breeding. Final stage of maturity is attained in the sea. The spawning occurs on the sea and the young ones brigrate to estuarine area. The process of maturation is a continuous process and spawning take place within a prolonged period. The observed maturity stages of females were (i) immature (ii) inactive / resting (iii) Developing (active) iv mature (v) ripe and running (.v) spent and of males were (I) immature (ii) maturing and (iii) spent.

#### सार्यम

कोचीन में ट्राल अवतरण से संग्रहित नमूनों में से जेरस फिलमेन्टस के पुनरूत्पादी जीवविझन का अध्ययन किया गया । मार्च - जून, 1998 के दौरान इस के जननांग, प्रौढावस्था, जननक्षमता और लिंग अनुपात का विश्लेषण किया गया । अंडक व्यास का मापन औश्र अंडाश्य की ऊतकीय विशेषताओं का अध्ययन किया गया । ज्वारनदमूखी परिस्थिति में यह जाति दिखाई पडती है और प्रजनन के लिए आस पास के तटीय क्षेत्रों में प्रवास करती है । प्रौढ़ता की अंतिम अवस्था ये समुद्र में प्राप्त करती है । समुद्र में अंडजनन संपन्न होता है और फिर से छोटे वाले ज्वारनदमुखी क्षेत्र में प्रवास करते हैं । अंडजनन और परिपक्ष्वन की ये निरंतर प्रक्रियाएं निश्चित अवधियों में चलते रहते हैं । मादा जाति की प्रौढ़ावस्थाएं (।) अप्रौढ़ (।।) निणिक्रय / सूप्त (।।।) विकासशील (1v ) सक्रिय ( (v) परिपक्ष और गतिशील (vi ) अंडारिक्त हैं । नर जाति की प्रौढ़ावस्थाएँ ( 1) अपरिपक्ष (11) परिपक्ष होने वाली और (111) अंडारिक्त हैं ।

\*\*\*\*\*\*

## **ACKNOWLEDGEMEN'TS**

I have great pleasure in expressing my deep sense of gratitude to my major advisor **Dr. K.K. JOSHI**. Scientist, C.M.F.R.I. for his excellent guidance and encouragement throughtout the period of my study. His sincere efforts and selfless support is inexpressible.

I sincerely thank **Dr. M. Devaraj**, Director, C.M.F.R.I. for providing me with all the facilities for the study. I would like to express my gratitude to **Dr. C**. **Suseelan**, officer in charge, Post Graduate programme in Mariculture for his help during the course of my work. My thanks are also due to **Dr. N.G. Menon** Sr. Scientist C.M.F.R.I. Co-chairman, advisory committee and **Smt. Grace Mathew**, Scientist, C.M.F.R.I. Member, advisory committee for their valuable suggestions.

My sincere thanks are also due to Shri.. Sanil, Scientist, C.M.F.R.I., Shri.. K.J. Abraham, S.R.F., PGPM, and Shri.. Satish, S.R.F for their valuable help during my work.

I am especially grateful to **Dr. Laxminarayana**, Scientist, C.M.F.R.I. and **Shri. P.M. Aboobackar**, Technical Officer, C.M.F.R.I. for their moral support and encouragement. I extend my sincere thanks to all the staff members of PFD division C.M.F.R.I. especially to **Smt. K.C. Seetha**, Technical Assistant.

I am especially thankful to Ms. Rosali Shaffer, Technical information specialist, National marine fisheries service, Panama City Laboratory library 3500 Dalwood beach road, Panama city, Florida 32408, U.S.A. for her unfailing response to my requests for references and reprints. I am thankful to Shri. B. Sudhir, M/s. Ram's Computech, Mamangalam for typing this dissertation.

My thanks are also due to all my classmates for the support and cooperation. I am grateful to my parents, sisters, brothers and all my family members for their encouragement and support. Finally I wish to express my sincere thanks to all my teachers and friends.

I would like to acknowledge my earnest thank to ICAR for the award of J.R.F. Above all this I give my praises and thanks to GOD Almighty.

### INTRODUCTION

The gerreidae, which comprise of eight genera and approximatly 40 species commonly known as majorras, silver biddies or purse mouths, is found is most warm seas, most of them are seen is tropical region, a few are temperate species. They are small to medium sized fishes living is sandy shallows of tidal creeks, lagoons, coral reefs and some enter into freshwaters. *Gerres filamentosus* cuvier is commonly called whipfin silver biddy. It is widely distributed in all warm seas of Indopacific, from the east coast of Africa through Indomalayan Archipelago, South China sea, Northern Australia and west Pacific islands.

In the estuaries and back waters of India it constitute an important fishery. They live in shallow coastal waters to a depths of atleast 50m, on sandy bottoms including coralline areas. Caught mainly by beach seines and bottom trawls, castnets, gillnets and seines. This species is a bottom feeder, with the protrusible mouth adapted for browsing on epiphytic animals like amphipods, apart from this it picks up other benthic organisms like polychaets bivalves, gastropods etc. This species grows to a maximum size of 25cm, 15cm is most common.

Gerres filamentosus cuvier remain is the estuarine enviornment until attainment of sexual maturity, fully ripe fishes undergo migration to near by coastal areas for the purpose of spawning, final stage of maturity was attained only in the sea. They probabily spawn at sea, the young ones migrate to estuarine area. Because of absence of any external character to distinguish the two sexes, the sex were determined after examining the gonads. Sex ratio was found to be 1:1 (female to male). But during the peak spawning season females were more compared to males.

Five maturity stages were determined on the basis of colour, shape, size and microscopic structure of gonad.

I. Immature:

Ovary occupying less than half of body cavity, pinkish jelly like translucent, very small irregular eggs.

II. Maturing Virgin (Recovering spent)

Ovary occupying half of body cavity, light yellowish, eggs not rounded, granular and yellowish white with yolk.

#### III. Ripening:

Ovary occupy 3/4 or more of body cavity, yellowish, egg opaque and not well rounded with yolk.

#### IV. Ripe :

Ovary occupy the entire body cavity, deep yellow with conspicuous blood vessels most of eggs are transparent with few opaque eggs. Some ova are visible to exterior vacuoles present in yolk.

5

#### V. Spent :

Ovary occupy slightly more than half of body cavity, dark pinkish, flaccid, translusent and shrunken with prominent blood vessels.

A knowledge of reproductive biology is essential in understanding the several features of reproduction in fishes. The present investigation was undertaken to elucidate the spawning activities, maturity stages and fecundity of *G. filamentosus* in the Cochin estuary.

#### Significance of the study

Investigations on maturation and spawning in fishes are of great importance. The fertility of a fish is of great significance in the reproduction of the stock of a species. Information regarding the reproduction, reproductive behaviour forms the basis of the management of broodstock in the aquaculture operations. Efficient management of brood stek depends upon accurate prediction of ripening tissue to facilitate the production of high quality seeds for culture. Reliable collection of high quality gametes requires a precise knowledge of maturation, ovulations and spawning of the candidate species. Accurate estimates of fecundity will enhance the efficiency of brood stock management and larval rearing programmes.

The information gathered from maturation and spawning studies can be used in ascertaining the age and size at which fish attains sexual maturity as well as spawning time and location of spawning sites. Together with fecundity estimate this information can be used to calculate the size of the stock and reproductive potential.

### **REVIEW OF LITERATURE**

Gerreidae is a family of marine Fishes living in coastal waters of tropical areas. Some species migrate to mangrove swamps, lagoons, estuaries and sometimes lower rivers (Bauchot, 1992). They are closely related to leognathidae, some of the earlier taxonomists (Bergh, 1940: Weber and Beaufort, 1931) have treated both the groups under one family leognathidae. But later after considering the characters of gerreidac. Such as, moderate sized scales on head and body, gill membrane free from isthmus; nuchal spines absent, and number of branchiostegial rays six, as against five in leognathidae. It was recognised as a separate family from leognathidae (Smith 1953; Munro, 1955). A systematic key was established to help identification of species of gerreidae through its otoliths (Lemos et al., 1993). Actually 40 gerreid species in seven genera are recognised (Nelson, 1994)

Several workers studied the distribution of different species of gerreidae. Day, 1878 reported that *Gerres setifer* is common in river Hooghly, attaining a length of 100mm, From Chilka lake it was first reported by Chaudhuri, 1923. It has a wide distribution in the sea and coast of India including estuaries. Some information on their landings and distribution was given by Devasundaram (1954), Jones and Sujansingani (1954) and Roy and Sahoo (1962), Jhingran and Natarajan (1966; 1969) have briefly described the fishery and Biology of this species. *Gerres lucidus* is a fish of east coast of India (Day, 1878; Nair, 1952; Hamsa and Kutty, 1972) and Minicoy island (Jones and Kumaran, 1980). Distribution of which in pulicat lake was studied by Kaliyamurthy, 1986.

Although various species of gerreidae are found usually in subtropical or tropical waters (Austin, 1971; Cyrus and Blaber, 1984b; Smith and Heemstra, 1986). One of the Australian species, namely southern silver belly *Parequilla melbournensis* is confined to the marine waters of Temperate southern part of this continent (Hutchins and Swainston, 1986; Gomon *et al.*, 1994). Indeed this was the most abundant fish species collected during extensive trawl netting of lower west coast of Australia in seven consecutive seasons (Laurenson *et al.*, 1993a). Different species of gerreidae utilize different habitat and have different degree of specificity in habitat preference. (Mathewson *et al.*, 1995).

Studies on maturation and spawning of teleostean fishes have been made by various workers from all over the world. An attempt to study maturity by measurements of ova diameter was first made by Clark (1934). Later Hickling and Ruthenberg, 1936 studied ovary as an indicator of spawning periods in hake, haddock, pilchard and herring. Dejong, 1940 investigated the spawning habits of thirteen species of teleostean fishes from Java sea. Hickling, 1940 estimated fecundity of Atlantic herring of Southern North sea by counting the eggs of mature ovaries. Clark (1950) observed the spawning habit of northern pike in North Western China. The investigation carried out on the teleosts like *Melanogrammus aeglefimus* (Robb, 1982) *Scomberomorus cavalla* (Funucane *et al.*, 1986); Euthymmus lineatus (Schaefer, 1987); Lutjanus griseus (Domeier et al., 1996); Sillago burrus and S. Vittata (Hyndus et al., 1996) revealed the reproduction biology of the species.

The relationship between attainment of first maturity and body size together with estimates of age composition and growth rates have been determined for populations of *Diapterus rhomheus* using a combinations of gonadal and length frequency data (Austin, 1971, Etchevers, 1978). *Gerres setifer* grows to a size of 110mm when one year old and 175mm when two year old (Patnaik S 1971). The maximum length attained by *G. filamentosus* cuvier is 175mm similar to that of gerreids *Gerres nigri*, *Diapterus rhombeus*, *Paraquuila melbourmsis*, but is less than that of *Gerres oyena* (300mm) (Austin, 1971, Albert and Desfossez, 1988, Kurup and Samuel, 1991, Sarre *et al* 1997). The use of scales to age *Gerres oyena* (Forskal ) showed that this species which grows to a total length of 300mm lives for up to 7 years (EI - Agamy, 1988). Age and growth rate of *Gerres setifer* was determined based on length - Frequency analysis (Anantha and Santha Joseph, 1995). Sarre *et al* 1997 used length frequency data scales and otolith for age determination of *Paraequila melbournesis*.

Reproductive system is a fundamental element in understanding fish stock variability. Early biological observations on the spanish Mackeral were made by Earil (1883) in Chesapeake bay and by Hildebrand and Cable (1938) along the south Atlantic coast. Reports on reproduction and other aspects of their biology were made for waters of Florida (Klima, 1959,) and of veracruz, Mexico (Medoza 1968).

Mojarras (gerreidae) are omnivorus and euriphagous fishes eating a wide spectrum of items mainly small invertebrates. Job (1940) gave an account of food of *Gerres punctatus* (*G. filamentosus*) along with food of other perches from Madras coast. Chacko (1949) furnished an account of food and feeding habit of *Gerres filamentosus*, *G. oyena and G. abbreviatus* from gulf of Mannar. Observations done on food and feeding habit of *Gerres Oyena* (Forskal) and *Gerres filamentosus* cuvier from pulicat lake by Rao, 1968. Concluded that both Species have identical food habit with similar construction of mouth. Both are bottom feeder with its protrusible mouth adapted for browsing on epiphytic animals like amphipods and also picks up other bethic organisms like polychaetes, bivalves, gastropods etc. (Arenas - G et al 1992)

Alves and Tom (1967) described the ouarian histology and size at first maturity of king mackeral *Scomberomorus cavalla* from Florida. Funucane *et al.*, (1986) studied the reproductive biology of *S. cavalla* from south eastern united states. Reproductive biology of Spanish mackeral *Scomberomorus maculatus* was also studied from this area by Funucane and Collins, (1986). Robb (1982) studied histological changes occur in the ovary of haddock *Melanogrammus aeglefinus* (L)

Histomorphology of ovarian changes during the reproductive cycle of Trachurus mediterraneas (Teleosti, Carangidae) was observed by Elgharabavi - et al., (1988). Grimes et al., (1988) investigated the reproductive biology of Tilefish Lopholatilus chamaeleonticeps Reproductive biological study of the pony fish Gazza minuta (Bloch) from PortoNovo east coast of India was done by Jayabalan (1988). An important component of many studies of fish reproductive biology is . the assessment of stage of gonad development of individual fish. The methods in use vary from highly detailed to cursery, but there are few reviews of their reliability or usefulness. West, 1990 reviews the methods of assessing ovarian development in fishes. This review examines histology, measurement of oocyte size, staging based on appearance of the ovary, and gonad indices. Sexual maturation of blue spotted mud hopper, Boleophthalmus pectinirostris was investigated histologically regarding gonad development and studied by G. S. I. egg diameter, composition etc. by Chung et al., 1991. Lopes et al., 1991 described the oocyte development in Piranha Pygocentrus natterevi knerr 1860 (Pisces, Characidae)

Ovarian germinative tissue of a South American teleost fish, the "Pacu" *Piaractus mesopotamicus* (Holmberg) has been studied through histological techniques, for germinative cellular type identification and maturity cycles charecterization of the fish species. (Lima *et al.*, 1991).

The reproduction biology of serranid fish *Plectropomus maculatus* sampled from inshore waters of the central great barrier reef was studied based on histological analysis of gonad material (Ferreira, 1993) and red drum *Sciaenops ocellatus* from nertic waters of the northern gulf of Mexico (Wilson *et al.*, 1994).

A histological study was carried out on the gonads of amberjack, Seriola dermerili to provide a basic knowledge of its reproductive biology with a view to a possible control of reproduction in captivity (Micale, et al., 1993), spawning in anchovy Engraulis lapensis (Meloy et al., 1994), spawning period of Pacefic Mackeral Scomber japonicus is estimated based on the changes in gonad index and ovarian histology (Murayama 1995); the ovarian morphology, annual reproductive cycle and spawning characteristics of kichipi rockfish Sebastolobus macrochir collected from the Pacific coast of south eastern Hokkaido, was examined on the basis of histological observation (Koya et al., 1995). Histological sections of European horse mackeral Trachurus trachurus (L) ovaries were used to follow oogenesis and to describe the atretic states (Karlous and economidis, 1996)

Prabhu (1956) studied maturation and spawning periodicities of seven species from Madras coast. Later studies on the reproductive biological aspects of *Chanos chanos* (Tampi, 1957); *Sardinella longiceps* (Nair, 1959); *Trichiurus lepturus* (James, 1967); *Trachysurus dussumeriri* (Vasudevappa and James, 1978); *Liza parsia* (Muthukaruppan, 1993); *Sillago sihama* (Jayasankar, 1989); *Mugil*  *cephalus* (Goplakrishnan, 1991) have described maturation, spawning, fecundity of teleosts of Indian coast.

Little information is available on the reproductive cycles of gerreidae, Although Austin (1971) and Etchevers (1978) studied the breeding of *Diapterus rhombeus* in the offshore waters of Puerto rico and Venezuela, Charles (1978) investigated the reproductive cycles of *Eucinostomus gula* in Biscayne Bay, Florida, U.S.A Reproductive biological study of Gerres in Natal estuaries by Cyrus and Blaber, 1984, similar studies on *Diapterus rhombeus* (cuvier) and *Gerres nigri* (Gunther) have been done in tropical esturies (Etchevers, 1978; Alberet and Desfossez, 1988). Despite their very high numbers in many areas and commercial importance is some regions (EI-Agamy, 1988; Kurup and Samuel, 1991). There have been few detailed studies on the biology of species comprising this family. (Castro and Cowen, 1991, Conover, 1992). Several species of gerreidae spent part of their life in estuaries (Cyrus and Blaber, 1982; Kershner *et al* 1985; Blaber *et al*, 1989; Nelson, 1994).

The only study on reproductive biology of gerreids which has incorporated type of histological examination of gonads, that is usually required to determine the precise duration of spawning period and whether spawning occures more than once in a breeding season is that carried out by Cyrus and Blaber (1984), on three species of gerres in estuaries of tropical Southern Africa. Of all methods determining reproductive state of female fish, histology is the most accurate (Hunter and Macewicz, 1985b) and depends upon the nature of study may be the only satisfactory method. Experimental induced spawning of patao *Eugerres brasilianus* (Pisces, Gerreidae) was done by Alvarez *et al.*, 1991. Nivon *et al* 1990 assessed the aquaculture potential of eight marine species from Maxican Pacific which include Silver biddies *Gerres cinerus* and *Eugerres auxillaris*. Sarre *et al* (1997) carried out investigations on the reproductive biology of a temperate gerreid in Australian waters.

A persual of Literature shows that the reproductive biology of gerreids of Indian waters have been reported by several workers (Jones and Sujansingari, 1954; Jhingran and Natarajan, 1969; Patnaik, 1971) Kurup and Samuel, 1991).

#### **DESCRIPTION OF SPECIES**

Gerres filamentosus cuvier

Vernacular name: whipfin Silver biddy

#### **Distinctive Characters**

Body compressed, elevated, its depth contained 2 to 2.5 times in standard length in larger specimens upto 3 times in smaller specimens. Anterodorsal profile ascending steeply at an angle of about 4<sup>o</sup>c to horizondal axis. Second dorsal spine laterally compressed, produced into a filament, whose tip extends past level of first anal spine. Third dorsal spine laterally compressed as long as distance from tip of snout to pre opercular margin., pectoral fin long, tip of depressed fin reaching to level of origin of first anal spine, third dorsal spine laterally compressed as long as distance from tip of snout to pre opercular margin; second anal fin spine much shorter than anal fin base; caudal tin deeply forked, its longest rays three times the length of median rays, (Fig 1)

#### Colour :

Silvery with 7 to 10 columns of ovoid spots on upper portion of sides, coalescing as bars in small specimens. Pectoral, Pelvic, Caudual and anal fins dusty hyaline, tips of pelvics, anal and lower lobe of caudal, white, dorsal fin hyaline except for the end of filamentous spine, which is black, It is silvery with a dark ovoid patch in the anterio dorsal quarter

#### Geographical distribution

Widespread in all warm seas of Indopacefic, from the east coast of Africa through Indo Malayan archipelago, South china sea, Northern Australia and the west Pacific islands





Fig. 1 (a) G. Filamentosus cuvier female (b) G. Filamentosus cuvier female showing ovary

## **MATERIALS AND METHODS**

#### **Collection of Specimen**

Samples of *Gerres filamentosous* were collected from Cochin Fisheries harbour at weekly intervals for a period of three Months (From March 1988 to June 1988). They are caught by trawl nets. The specimens collected had a length range of 120mm to 190 mm

The fish after collection were carried to the laboratory and kept in freezer. The Morphometric measurements like total length, standard length, fork length, head length, depth were measured. The body weight were taken accurately in a sensitive balance. Then the sexes of specimen are identified by cutting open the belly.

Maturity stages of females were identified based on microscopic examination of ovary shape and size in relation to body cavity, colour, extend of yolk formation and microscopic structure such as ova diameter measurement (Clark, 1934).

The ovaries were taken out from body cavity and kept in 10% formalin for preservation. It is used for Fecundity study and ova diameter measurement. A small part of formalin preserved ovary tissue is taken and spread on a microslide. The oocyte were carefully separated with fine needle as evenly as possible to spread out on the slide, Diameter measurements of atleast 100 oocyte from each ovary was carried out under the microscope with an ocular micrometer at a magnification (10x) Ovaries from 30 number of specimens were taken for fecundity estimation.

#### 1. Sex Ratio

A total of 246 fishes collected from commercial catches were examined for determining sex ratio. It was not possible to detect any external characters such as body proportions, meristic charecters or colour which could be useful for sex determination. Sex is determined only by gonadal examination. Total number of individuals belong to each sex were counted. A knowledge of sex composition of catch helps in understanding whether any differential fishing exists between males and females and if so, its possible bearing on the fishable stocks.

#### 2. Size at first Maturity

The fish specimen with gonads undergoing active gametogenesis were considered as mature. The length at which 50% of total individuals are nature was considered as size at first maturity (Beverton and Holt, 1957)

#### 3. Fecundity

Fecundity was estimated by counting the number of mature ova in a weighed portion of ovary and computed for the whole. The regression equation  $\log y = a+b \log x$  was fitted to the data to determine the relationship between fecundity and body weight (grams) as well as total length (mm) here,

#### 4. Oocyte diameter

Size of ova is a good indicater of maturation and gonadal cycle (Davis, 1985). Oocyte diameter was measured along its horizontal axis using ocular micrometer pre caliberated with stage micrometer. Oocytes were measured at random till the count reaches 100 oocyte. The data obtained was used to work out largest size frequency profile.

#### 5. Length frequency analysis

This method is based on the assumption that length of individuals of same age group in a population of fish are approximately normally distributed.

#### 6. Histology

Histological characteristics of ovary at different stages of maturity were studied. Sample pieces of ovary 15mm thick were taken from freshly collected specimens and fixed in Bouins fixative for 12-24 hrs. The tissue was then thoroughly washed with running water for 4 -5 hrs and then stored in 70% ethyl alcohol until further processing. The tissue were dehydrated in graded alcohol series by following standard procedure. This tissues were then cleared with chloroform for 3-4 hrs impregnated with and embedded in paraffin wax . The material was then sectioned at 6-8 µm in a rotary microtome. The sections were spread over slides on which a thin layer of Mayors glycerol albumen adhesive was applied earlier, This adhesive is a combination of egg white and glycerol in 1:1 ratio. The sections were deparaffinised, hydrated and stained with haematoxylin and 1% aques eosin as counterstain. DPX was used as mounting medium for all slides.

# RESULTS

#### 1. Maturity Stages

#### Male

The male gonads were classified into four stages, immature, maturing, mature and spent. The testes were small and occupied very little portion amounting to about 1/10<sup>th</sup> of the body cavity in the immature condition. Maturing testes were thick and little flattened whitish in colour and occupied half of the body cavity. In Mature condition there were white in appearance and occupies 3/4<sup>th</sup> of the body cavity. Spent testes some what whitish red in colour and shrunken.

#### Females :

The following stages of sexual maturity were observed in the case of females (Fig. 2a).

#### Stage I (Immature)

Ovaries appears thick, short, transparent tubes occupies small portion of the body cavity. There were present throughout the study period but is less in number as compared to other stages. Oogonia (Mean size:  $10 \ \mu m$ ) were found in the ovaries. Diametr of the Oogonia ranged forms 2 to  $15 \ \mu m$  in this state.





Fig 2 (b) Fig. 2 (a) G. Filamentosus cuvier female maturity stage 1-V (b) Stage II (c) Stage V

#### Stage II Inactive/Resting (Recovering spent)

Ovaries were yellowish, occupies one third of the body cavity : Ova visible to naked eye. Large number of pre-vitellogenetic oocytes were present in immatures and inactive ovaries which showed no evidence of development towards llIrd type Oocytes. Diameter of the oocytes showed range of 10 µm to20µm (Fig. 2b).

#### . Stage III Developing (Active)

Ovaries occupies about half of the body cavity and yellowish in colour. (Fig. 2a). As the development begins the oocytes increase in size, developing into yolk precursor stage. The nuclei showed irregular cuticle. (Fig 3a). Diameter of the oocytes showed range 15 µm to 30µm.

#### Stage IV - Mature :

Ovaries occupies about three forth of the body cavity and deep yellow colour. At this stage non cellular membrane begins to form between follicular layer and developing oocyte. Fig. 3 b illustrates an early non staining yolk stage oocyte. In stage IV development oocyte contains numerous yolk granules and a well developed chorion (Fig. 4 a). The ova diameter showed a range of 20  $\mu$ m to 35 $\mu$ m. The ovary contains batches of vocculated oocytes, some of which contains developing yolk granules.

#### Stage V (Ripe and Running) :

The mature Oocytes (Fig 2c) in stage V are all in various stages of yolk accumulation. (Fig 4b). Fig.5 (a-b) shows early and peripheral development of the secondary yolk componant appears as stained granuels. The cytoplasom is completely fiilled with secondary yolk granules. During this stage ripe Oocytes undergo hydration, resulting in transparency and considerable increase in volume. This leads to the rupture of the follicle and the eggs passed out though the oviduct.

#### Stage VI - Spent:

Ovaries small occupies less than 1/3rd of body cavity.

### 2. Fecundity

The fecundity of specimens ranging 142 to 161 mm, 141 to 172 mm and 141 to 180 mm corresponding to the maturity stages III, IV and V are given in the table 1, 2 and 3.

In the stage III fecundity showed a range of 5460, to 88695 eggs for the fish ranging in length from 142 to 157 mm The number of eggs produced by the fish of different length groups shows that the fecundity of this fish more or less

increases with length (Fig 6). The fecundity in the length range 140-145mm is below 10,0000 and increases upto 80,000 in the length range and shows slight decline in the higher length groups.

In the stage IV fecundity showed a range of 17400 to 89,560 for fish ranging from 141 to 195 mm. The fecundity in the length range 140 to 145 is below 25,000 and thus shows an increase upto 50,000 eggs in the length group 150-155 and finally to 70,000 eggs in the range 170 to 175 mm group (Fig 7).

In the stage V fecundity showed a range of 21990 to 41750 eggs for the fish ranging in length from 141 to 180mm in this stages. (Fig 8).

#### 3. Maturity condition

#### Female

Maturity percentage of fishes is given in the **Table 4** It is observed that in the length range 100-115 all the fishes were immature condition. From 115 mm onwards number of immature fishes decreases and is 165-170 mm range the immature percentage is zero. Mature females were observed in the 115-120 mm length range (25%) and progressively increases and in the 125-130mm range 50% females were mature. In the length range 145-150mm 80% females were mature. The sample shows that the fishes above 165mm sizes were all in mature condition. A total of 150 samples were analysed and it is found that 73 belongs to immature and 77 belongs to mature condition of fish. Size at first maturity was calculated as 125 mm (Fig 9)

#### Male

Maturity percentage of male fish were given in the **Table 5.** It is observed that in the length range 100-105 all the fishes were in immature condition. As the length increases the percentage of maturity also shows increasing trend and in the 110-115 length range 50% fishes were mature. In the size range 140-145 almost 80% of fishes were in the mature conditions and is 150-155, the maturity is 100%.

#### 4. Sex Ratio

A total of 246 fishes were studied and it is observed that 150 are female fishes and 96 are male ones (**Table No:6**). Sex ratio was found to be female dominated throughout the period of study (IM :1.5F). During the month of march the sex ratio was 1:1.3 (male : female) and showed increase in the female number during the April (IM:2.6F). During June the sex ratio was IM : 1.1 F.

#### 5. Length - Frequency

Length frequency distribution of G. Filamentosus during the period of study is shwon in the Fig 10 to Fig 16.

#### 6. Ova-diameter

The Oocyte - diameter frequency of mature ovary are shown in the Fig. 17 to Fig. 20.

# Table No:1 Fecundiity of G. filamentosus

# Maturity Stage III

SI. No.	Total length	No. of Ova
1	142	5460
2.	147	35640
3.	150	43700
4.	151	26200
5.	155	26150
6.	157	88695
7.	160	50820
8.	161	28250

SI. No.	Total length	No. of Ova
1.	141	17400
2.	144	23700
3.	149	24920
4.	150	43740
5.	155	20160
6.	156	20880
7.	157	53100
8.	160	21680
9.	164	25640
10.	171	65696
11.	172	78560
12.	195	89560

# Table No :2 Fecundity of G. filamentosus. Maturity Stage IV
SI. No.	Total length	No. of Ova	
1.	141	21990	
2.	144	26295	
3.	149	43380	
4.	150	36480	
5.	155	32120	
6.	159	40960	
7.	164	39180	
8.	165	35250	
9.	169	43850	
10.	180	41750	

# Table No : 3 Fecundity of G. filamentosus,Maturity Stage V

Length (mm)	Immature	Mature	Total
100-105	4 (100) -		4
105-110	6 (100)	-	6
110-115	8 (100)	n na	8
115-120	12 (75)	4 (25)	16
120-125	16 (77)	5 (23)	21
125-130	2 (50)	2 (50)	4
130-135	2 (33)	4 (66)	6
135-140	2 (25)	6 (75)	8
140-145	3 (28)	8 (72)	11
145-150	3 (17)	15 (83)	18
150-155	8 (44)	10 (54)	18
155-160	5 (36.8)	9 (64)	14
160-165	2 (28.6)	5 (71.4)	7
165-170	-	2 (100)	2
170-175	-	2 (100)	· 2
175-180	-	2 (100)	2
180-185	-	1 (100)	1
185-190	-	1 (100)	l
190-195		1 (100)	1
	73	77	150

 Table No: 4 Maturty Percentage of Female G. filamentosus

Length (mm)	Immature	Mature	Total
100-105	3 (100)	3 (100) ~	
105-110	2 (66)	1 (33)	3
110-115	1 (50)	1 (50)	2
115-120	2 (40)	3 (60)	5
120-125	1 (33)	2 (66)	3
125-130	1 (33)	2 (66)	3
130-135	2 (17)	10 (83)	12
135-140	5 (25)	15 (75)	20
140-145	2 (16)	11 (84)	13
145-150	1 (12)	8 (88)	9
150-155		9 (100)	9
155-160	-	5 (100)	5
160-165	-	2 (100)	2
165-170	-	2 (100)	2
170-175	-	1 (100)	l
175-180	<b></b>	1 (100)	1
180-185		1 (100)	1
185-190		2 (100)	2
190-195		-	<b>7</b> .
	20	76	96

 Table No: 5 Maturty Percentage of Male G. filamentosus

	Female	Male	Total	Male : Female
March	52 (57%)	38 (43%)	90	1:1.3
April	58 (65%)	22 (35%)	80	1:2.6
May	40 (55%)	36 (45%)	76	1:1.1
Total	150 (60%)	96 (40%)	246	1:1.5

Table No: 6 Sex ratio of G. filamentosus



Fig. 3 (a) Stage III Oocytes at yolk precursor stage



Fig. 3 (b) Stage IV Early non staining yolk stage



Fig. 4 (a) Stage IV Chorion development



Fig. 4 (b) Stage V Early yolk staining stage



Fig. 5 (a) StageV Early yolk stage - Single oocyte



Fig. 5 (b) Stage V

















## Fig.10.Length-frequency distribution of Gerres filementosus

## Fig.12.Length-frequency distribution of Gerres filementosus





# Fig.13.Length-frequency distribution of Gerres filamentosus

## Fig.14.Length-frequency distribution of Gerres filementosus





# Fig.15.Length-frequency distribution of Gerres filementosus









Dogra diamar (mu)

Fig. 18. Oocyte diameter frequency graph of different stages ovary (19.3.98)



Fig. 20. oocyte diameter frequency graph of different stages ovary (1.4.98)



## DISCUSSION

#### Areas and seasons of spawning

Fish reproduction is almost always a seasonal or cyclical phenomena in the tropical and subtropical reagion, were environmental conditions seems relatively stable, there may be still be a cycle of gonadal maturation impossed by the energy demands of maturing a batch of eggs (Hoar, 1969).

The breeding biology of fishes of the family Gerreidae, of Pulicat and Chilka lakes was investigated by Jhingran and Natarajan (1969), Rao (1970) and Patnaik (1971). In Chilka lake, Jones and Sujansingani (1954) observed the male specimens of *G. setifer* in ripe condition with flowing milt and female with ovary almost in ripe condition, but they were not able to say whether the species bred in the lake or not. Jhingran and Natarajan (1969) observed the occurrence of males and females of *G. setifer* with oozing gonads in the southern sector of Chilka lake. Rao 1970 stated that *G. oyena* with oozing gonads were not encountered from Pulikat lake and so he persumed that the final stage of maturity of the species was attained only in the sea. Fishes with fully matured gonads occured through out the year and hence concluded that *G. oyena* bred in the east coast of India in the year round.

Patnaik (1971) noticed that G. setifer bred in Chilka lake and outwardly migration of this species from Chilka lake had not been taking place. Kurup and

Samuel (1991) observed that fishes collected from Cochin barmouth areas of Vembanad lake showed a higher degree of maturity than those specimens collected from adjacent areas and hence assumed that fully ripe fishes undergo migration to near by costal area for the purpose of spawning. Specimens of *G. filementosus* with oozing gonads were not observed in their study, inferred that the final stage of maturity was attained in the sea.

In the present study the specimens collected from trawlers includes G. *filamentosus* with oozing gonads inferred that the final stage of maturity was attained in the sea. Not only the oozing the gonads were present but the matured specimens collected from the inshore area of depth range 10-20 m supports the above conclusion. The present results agreees with the results of Kurup and Samuel (1991), where samples are mainly collected from gillnets, sines and castnets. Cyrus and Blaber (1989) carried out investigation about the reproductive cycle in the Natal estuaries concluded that hatching as well as spawning of G. *filamentosus*, G. *acinacus* and G. *rappi* occurs throughout at the sea.

Sarre *et al.* (1997) observed a temperate Gerried *Parequula melbournensis* spends throughout its life cycle in offshore area. The samples were abundant in the inner continental shelf of lower cost of Australia at depth range of 5-35 m. The Juveniles of some species of closely related family Leognathidae were also found in areas other than along the shore line and in estuaries (Blaber *et al.*, 1994, 1995). Domeir *et al.*, 1996 observed spawning migration of Gray snapper in Florida, they

observed movements of adult Gray snapper from inshore to offshore sites. Similarly Hyndes et al., 1996 noted spawning migration of Sillago burrus and Sillago vittata.

The endocrine system forms the major link between environment and the organs concerned with reproduction. Changing environmental condition, operating through the sensory systems and specific centers in the brain, trigger neurosecretion which inturn regulate the activities of pituiary gland. The pituiary hormons have direct effects on gametogenesis, matabolism and behaviuor. These hormones also regulate the development of gonadal endocrine tissue. Many marine fish species in the tropical or subtropical waters, including those of Gerriedae have long spawning season where as those species that inhabit temperate waters, where conditions in the winter are more severe, typically have shorter more clearly defined breeding periods(Castro and Cowen, 1991; Conovar, 1992)

The spawning seasons of *G. setifer* in Chilka lake was from May to September (Patnaik, 1971). Spawning period of *G. filametosus* in cochin estuary was from October to February (Kurup and Samuel, 1991). In the present study, samples collected during March to June showed all the maturity stages in the sample. The presence of advanced oocyte in some ovary during April, suggest that spawning was imminent in that month. The lack of clearly defined modal length classes in the sample in this season and no conspicuous progression of modal length classes over successive seasons indicate that spawning occurs in this season. Compiling earlier works, it can be assumed that the spawning of G. filamentosus occuring throughout the year. Quasim(1955) and Harder Jones(1968) have studied that protracted spawning seasons are charecteristic in tropical region, because of duration of period of suitable temperatures and food availability, for survival of juveniles

The spawning of *P.melbournensis* occurs throughout the year (Sarre *et al*, 1997)and far less protracted spawning periods for similar species Mugilidae, Clupiedae, Gerreidae(Chubb *et al*, 1981, Chubb and Potter 1984; Sarre *et al.*, 1997). Multiple spawning over a prtracted period or even full year is associated with species found in less seasonal environments (Burt *et al.*, 1988). Multiple spawning increases the number of eggs that can be produced in a year (Burt *et al.*, 1988), spreads the risk of predation on eggs and larvae over an extented period (Lambert and Wase 1984) and acts as a buffer against adverse fluctuation in the amount of food available to the larvae (Mc Evoy and Mc Evoy, 1992).

#### **Gonad development**

Arora (1951) carried out investigations of *L. Splendens* in Rameswaram island and arrived at some conclusion regarding to the breeding habit of this fish. He classified females into three groups based on the gross examination of ova. The three groups are immature, intermediate and maturing groups. Immature ovaries contained number of eggs visible to the naked eye; Intermediate ovaries small, creamy white, opaque eggs visible to the naked eye and maturing ovaries

contained larger eggs which are translucent rather than opaque, According to Arora, *L. splendens* spawns more than once in a season. Obviously the fish spawns from March till August and September but the peak spawning occurs during the months of April and August. He stated that the species first matures at an average length of 60mm. Standard length.

Kurup and Samuel (1991 classified ovaries of *G filamentosus* into five stages, based on the ovadiameter studies. They were **1**. *Immature stage* : Most of the oocytes were less than 0.2 mm with mode at 0.05 mm. This immature oocytes were found in the ovaries of all stages of development. **2**. *Maturity virigin and recovering spent* : Immature oocytes and another batch of ova of 0.2 mm was seen. **3**. *Ripening* : A group of opaque ova with a mode of 0.3 mm were found **4**. *Ripe* : One batch of transparent ova with a mode of 0.45 mm **5**. *Spent* : Mode was 0.05 mm.

Cyrus and Blaber (1984) were recorded seven stages in ovaries of G. *filamentosus, G. acinales* and G. *rappi*. Sarre *et al* (1997) found seven stages in the ovaries of P. *Melbournesis*. On the basis of the histological observation seven stages of Oogenesis was postulated.

State I : Oogonia were present throughout the year in addition and mean size was 10µm.

Stage II : Large number of pre vitellogenic Oocytes were present in immature and inactive ovaries. These Oocytes had large nuclei (23µm).

Stage III : At onset of gonad development the Oocyte increase in size developing into yolk preserved stage (34  $\mu$ m)

Stage IV : In this non staining yolk stage a non cellular membrane begins to form between follicular layer and the developing oocyte.

Stage V: In the red-staining yolk stage development in Gerres is characterised by Cytoplasm which is completely filled with secondary yolk and few primary granules

Stage VI : Classified as ripe

Stage VII : spent

In the present histological observation, it was found that six maturity stages in the case of females. They were 1. Immature 2. Inactive/resting 3. Developing 4. Mature 5. Ripe and running 6. Spent (Modified from Kesteven, 1960; Nikolsky, 1963; Davis, 1977, Cyrus and Blaber 1984; Kurup and Samuel 1991; Sarre *et al.*, 1997). The male gonads were classified into four stages in the present observation, they were immature, maturing, mature and spent.

The main stage of Oogenesis were investigated and documented according to developmental sequence. Similar to studies done by Dipper and Pullin (1979)

which was divided as follows, stage i - Oogonia, Stage ii - Previtallogenic oocytes; stage iii to v - Vitellogenesis; Stage iii - yolk precursor; Stage iv - nonstaining yolk; Stage v- red staining yolk Stage vi - completion of development.

The developing oocytes lie along the ovigerous folds, embedded in the loose connective tissue (Cyrus and Blaber 1984) Main stages of oogenesis of *Gerres* is Natal estuaries was as follows, Oogonia were present throughout the year in adult. Oogonia develops into Pre-vitellogenic oocytes are were present is immature and inactive ovaries. Small numbers present in developing ovaries but fewer were observed in mature ovaries. This oocyte develops into yolk precursor stage. The nuclei were irregular and nucleoli were visible. A non cellular membrane begins to form between the follicular layer of yolk precursor stage and develops into Non staining yolk stage. Next stage was development of small red stained granule (Red-stained yolk) and complets the development.

#### Fecundity

A total count of ova in both ovaries yeilds an estimate of female reproductive potential or fecundity. Patnaik (1971) observed that the fecundity of *G. setifer* varied from 17,2937 to 1,61,505 eggs in length range 88 - 193mm fish. Rao (1970) observed that the fecundity of *G. oyena* varied from 1,04,211 to 14,43,785 eggs in the size rages from 148-282 mm. Kurup and Samuel (1991) observed that the fecundity of *G. filamentosus* varied from 64,278 to 3,87,576 in the size range of 100-148 mm SL. In the present study the observed fecundity varied from 5460 to 89,560 in the size range of 141-195 mm.

According to Simpson (1954) and Beverton and Holt (1957) the number of ova should depend upon the volume of the ovary. Therefore, fecundity has been related to length cubed (Bagenal, 1968). Fecundity is a function of size and size is obviously a reflection of the growth process which is directly affected by temperature interacting with nutrition. Temperature functions primarily as a controlling factor, where as diet and ration function as limiting factor (Honston, 1982).

#### Sex ratios

Table 6 shows the sex ratio obtained in the present study. The ratio was found to be deviated from the normal expected value 1:1 ratio. The sex ratio was significanly different in March, April, and May. The present result was in agreement with the results of Kurup and Samuel (1991). The skewed sex ratio was due to the ponderance of females over males. Quasim (1966) suggested that the ponderance of one sex in a population was because of sexual difference in growth rate between sexes.

The difference in sex ratio have been suggested to be due to difference between males and females with respect to growth, mortality or availability (shaefer, 1987). The diversity in the sex ratio might be due to partial segregation of mature forms, either through habitat preference or because of school formation, then rendering one sex to be more easily caught than the other. During the period March - May season females were more as compared to males in the present species may be due to behavioural difference between two sexes.

#### Size Composition

The maximum length of *G. filamentosus* reported in the present study was 195 mm is similar to that of gerreids *G. filamentosus G. nigri, Diapterus rhombeus,* and *P. melbournensis* but is less than that of *G.oyena* (300 mm) (Austin, 1971; Albaret and Des fossez 1988; El-agamy 1988; Kurup and Samuel, 1991; Sarre *et al.*, 1997). Absence of clearly defined modes in the sequential defined modes, in the sequential length frequency data means that model progression analysis could not be used to elucidate either age composition or growth rate.

## SUMMARY

- 1. A brief study of the reproductive biology of *G.filamentous* occuring at Cochin estuary was carried out from March 1998 to June 1998. A total of 246 specimens of *G. filamentosus* were examined for determining gonadal maturity stages, ova diameter, fecundity and spawning based on the standard methods used.
- 2. In the present study, the specimens collected from trawlers included G. filamentosus with oozing gonads, inferred that final stage of maturity was attained in the sea. Not only the oozing gonads were present but the mature specimen collected from the inshore area supports the above conclusion.
- In the case of female G. filamentous six stages of maturity were observed. They were (i) Immature (ii) Inactive/resting (iii) Developing (iv) mature (v) Ripe and running (vi) Spent. The male maturity stages were (i) immature (ii) maturing (iii) mature (iv) spent
- 4. The fecundity observed in the present study varied from 5,460 to 89,560 in the size range of 141-195 mm. The sex ratio were found to be deviated from the normal expected value 1:1 ratio, during the month of March, April and May.

## REFERENCES

ALBARET. J.J. and DESFOSSEZ. P. 1988. Biology and ecology of Gerreidae (Pisces, Teleostei) in the Ebrie Lagoon (Jvory Coast). Revue d' Hydrobiologie Tropicale 21, 71-88.

\*ALVAREZ- LAJONCHERE, L; PEREZ, L; HERNANDEZ, O.G. 1991. Spawning experimental induction of patao *Eugerres brasilianus* (cuvier) (Piisces,Gerreidae). *Rev.Cub. invest. Pesq.*, Vol.16, No.1-2, pp. 33-39.

\*ALVES, M.J. M., AND TOME, G.S. 1967. Alguns aspectos do desenvolvimento maturativo das gonadas da cavala, *Scomberomorus cavalla* (cuvier, 1829). *Arq.Estac. Biol. Mar. Univ. Fed.ceara* 7 (1):1-9

ANANTHA,C.S.,SANTHA JOSEPH,P. 1995. Age and growth of *Gerres setifer* (Hamilton-Buchanan) inhabiting the coastal waters of Mangalore. *Environ.Ecol.*, Vol. 13, No.2,pp406-409.

\*ARENAS-GRANADOS, P; ACERO, P.A. 1992. Trophic structure of the Mojarra (Gerreidae) from Samta Marta big Marsh (colombias Caribbean).

ARORA, H.L., 1951. A contribution to the biology of the Silver belly, Leiognathus splendens (Cuv). Proc. Indo-Pacific. Fsh. Comm., Sec. III: 1-6 AUSTIN, H.M. 1971. Some Aspects of the biology of rhomboild mojarra *Diapterus rhombeus* in puertovico. Bulletin of marine science **21**, 886-903.

BAGENAL, T.B. 1957 The breeding and Fecundity of Long Rough Dab, *Hippoglossoides platessoides* (Fabr) and the associted Cycle in condition. *J. Mar. biol. Ass.U.K.***36**: 339-75

BAGGER MAN, N. 1980 Photoperiodic and endogenous control of the annual reproductive cycle in teleost fishes pages 533-567. In M.A. Ali, (eds), Environmental physiology of fishes plenum, New York.

BAGLIN, R.E., Jr. 1982 Reproductive biology of western Atlantic bluefin tuna.and *Fish. Bull.*, U.S. 80: 121-134

\*BAUCHOT, M.L.1992. Gerreidae fresh and Brackishwater Fishes of west Africa, Vol. 2, 1992.pp 691-693

BEAUMARIAGE, D.S. 1973. Age, growth and reproduction of king Mackeral, Scomberomorus Cavalla in florids, Fla. Mar.Res.pub: No.: 1 45p

BERGH, L.S. 1940. Classification of fishes both recent and fossil Trav. Inst. Zool. Acad. Sci. U.S.S.R 5(2) : 517 PP (Russian and English texts). Reprint 1947, Ann Arbor, Michigan, J.W. Edwards Brothers.

\*BERTIN, L. 1958A. Sexulite at Fecondation. In "Traite' de Zologie", p.p. Grasser, (ed), Vol XIII, Past 2 pp 1584-1652

BEVERTON, R.J.H.and HOLT, S.J.1957. On the dynamics of exploited. Fish population *Fish.Invest. London*, Ser.2. 19: 533pp

BLABER, S.J.M. BREWER, D.T. and SALINI, J.P.1995. Fishcommunties and the nursery role of the shallow inshore waters of tropical bay in the Gulf of Carpentavia. Australia. *Estuarine, Coastal and Shelf Science* 40, 177-193

BLABER, S.J.M., BREWER, D.T. and SALINI, J.P. 1989. Species Composition and biomasses of fishes in different habitats of a tropical northern Australian estuary;their occurance in the adjoining sea and estuarine dependance *Estuarine*, *coastal and shelf science* **29**, 509-531

BLABER, S.J.M., BREWER, D.T. and HARRIS, A.N. 1994. Distribution, biomass and community structure of Demersal Fishes of Gulf of Carpentaria. Australia.

BLACKBURN, M., R.M. LANURS, R.W. OWEN, AND B.ZEITSCHEL 1970.

Seasonal and areal changes standing stock of phytoplankton, Zooplankton and micronekton in the eastern tropical pacific. Mar. Bio., 7 (1) 14-31

BURT, A., KRAMER, D.L., NAKATSURA, K.and SPRY, C. 1988.

The tempo of reproduction in Hydphessobrycon pulchripinnis (Characidae) with a discussion on the biology of multiple spawning in fishes. *Envi. Bio. of fishes* 28, 15-27.

BYE, V.J. 1984. The role environmental factors in the timing of reproductive Cycles,

CASTRO, L.R. and COWAN, R.K. 1991. Environmental factors affecting the early Life history of bay anchpvy. *Anchoa mitchilli* in Great South Bay, New York.

CHACKO, P.J. 1949. Food and Feeding habits of fishes of Gulf of Mannar. Proc. Indian Acad. Sci., 29 (3), SecB., 83-97.

\*CHARLES, R. 1975. Aspects of the biology of the Mojarra Eucinostomus gula (Quoy and Gaimard) in the Bis cayne Bay, Florida, Msc thesis Unvversity of Miami

CHAUDHURI, B.L., 1923. Fauna of Chilka lake. Fish, Part IV. Mem. Indian. Mus., 5 (11): 711-36

CHUBB, C.F. and POTTER, I.C. 1984. The reproductive biology and estuarine movements of Gizzard shad, Nematalosa vlaminghi (Munro) Journal of Fish Biology 25, 527-543

CHUBB, C.F., POTTER,I.C., GRANT, C.J., LENANTON, R.C.J., and WALLACE, J. 1981. The age structure growth rates and movements of sea mullet *mugil cephalus* L., and yellow eye mullet, *Aldrichetta forsteri* (Valenciennes) in

the Swan-Avon River System. Western Australia. Australian Journal of Marine and Fresh water Research 32, 605-628

CHUNG, - E - y : AN, - C.M.; LEE, T. Y 1991. Sexual maturation of the blue spotted mud hopper *Boleophthalmus pectnirostris* (Linnaeus) *Bull - Kovean - Fish, -Soc. 1991*, Vol.24, No.3, pp : 167-176

CLARK, C.F. 1950. Observations on the spawning habits of the northern pike, Esox lucius Linnaeus, in North Western China Copeia : 4 : 285-288

CLARK, F.N. 1934. Maturity of the California Sardine (Sardina Caerulea) determined by ova diameter measurements Calif. Div. Fish Game : Fish. Bull (42): 1-49.

CONOVER, D.O. 1992. Seasonality and the Scheduling of life history at different latitudes *Journal of Fish Biology* **41**, 161-178

CRIM, L.W., AND B.D. GLEB 1990. Reproduction Methods for Fish Biology. Edt. C.B. Schreck and P.B. Moyle American Fisheries Society. USA.

CYRUS, D.P. and BLABER, S.J.M. 1982. Species identification, distribution and abundance of Gerreidae (Teleostei) Bleeker, 1859 is the estuaries of Natal. South African Journal of Zoology 17, 105-116

CYRUS, D.P and BLABER, S.J.M. 1984 b. Predation and sources of mortality of Gerreidae Bleeker, 1859 Teleostei in Natal estuaries with special references to the Kosi System. *The Lammergeyer* **32**, 14-20

CYRUS, D.P. and BLABER, S.J. M. 1984 a. The reproductive biology of Gerres in Natal estuaries. *Journal of Fish Biology* 24, 491-504

DAVIS, T.L.O., 1977. Reproductive biology of freshwater Catfish *Tandanus tandaners* Mitchel) in the Gwydlr river, Australia. 1- structure of gonads. *Aust. J. Mar. Fresh water Res.*, 1977, **28**; 139-158.

DAY, F., 1878. Fishes of India. Willian Dawson and Sons Ltd., London, 778 pp

DEJONG, J.K. 1940. A preliminary investigations of spawning habits of some fishes of Java Sea, Treubia 17, 307-30.

DEVASUNDARAM, M.P. 1954. A report on the Fisheries of Chilka lake from 1948-1952. Government press, Cuttack: 1-34

DOMEIER, M.L., C. KOENING AND F. COLEMAN 1996. Reproductive biology of the Gray Sanpper, (*Lutjanus gsiseno*), with Notes on spawning for other Western Atalantic Snappen (Lutjanidae). In Biology, fisheries and culture of tropical grupers and snappers 1996. *ICLAM Conf. Proc.* 48, 189-201.
\*EARLL, R.E. 1883. The spanish Mackerel, *Cybium maculatum* (Mitch) Ag. Its natural history and artifical propagation, with an account of the origin and development of the fishery. *Rept. U.S. Comm. Fish.* (1880) pt. 8: 395-4264

Edt. C.B. Schreck and P.B. Moyle American Fisheries Society. USA.

EL GHARABAVY, -M.M., ABDEL-AZIZ, -S.H. 1988. Histomorphology of ovarian changes during the reproductive cycle of *Trachurus mediterraneus* (Teleostei, Carangidae) *Bull, -Natk, -Inst, - Oceanogr. -Fish. -Egypt.* 1988. Vol.14, no.2, pp. 55-69

EL-AGAMY, A.E. 1988. Age determination and growth studies of *Gerres oyena* (Forskal) in the Arabian Gulf waters. *Mahasagar* **21**, 23-34

ETCHEVERS, S.L. 1978. Contribution the biology of *Diapterus rhombeus* (cuvier) (pisces - Gerreidae). South of Margarita island, venezuela. *Bulletn of marine science* 28, 385-389

FERREIRA, -B.P. 1993. Reproduction of the inshore coral trout *plectropomus maculatus* (perciformes : Serranidae) from the central Great Barrier Reef, Australia *J. -Fish - Biol. 1993* Vol.**42**, No.6 pp 831 - 844

FORSBERGH, E.D., 1969. On the climatology, Oceanogsaphy and fisheries of the Panama Bight. Inter-Am-Trop. Tuna Comm., Bull., 14 (2): 45-385.

FORSEBERGH, E.D. 1963. Some ralationships of meterological, hydrographic, and biological variables in the Gulf of Panama. *Inter-Am. Trop. Tuna Comm., Bull.*, 7 (1): 1-109

FUNUCANE J.H., COLLINS L.A., BRUSHER, H.A SALMAN C.H. 1986. Reproductive biology of king mackeral, *Scomberomorus Cavalla*, from South eastern united states. *Fishery Bulletin*, vol 84, No.4, 1986

FUNUCANE, J.H and COLLINS, L.A. 1986. Reproduction of spanish mackeral, Scomberomorus Maculatus, from the South eastern united states. North east Gulf Science Vol. 8 No.2, November 1986 p. 97-106

\*GERAD, P. 1985. Organes reproducters. In : Traite de Zoologie "(p.p. Grasse ed.) Vol. 13 (2) 1565-1583.

GOMON, M.F., CLOVER, J.C.M. and KUITER, R.H. 1994. The Fishes of Australia's South coast. Adelaide : State print

GOPALAKRISHNAN. A. 1991. Studies on some aspects of the reproductive physiology of the female grey mullet *Mugil cephalus* L. *Ph.D Thesis. Cochin university of Science and Technology*. 214 pp

GRIMES, C.B; IDELBERGER, C.F; ABLE, K.W; and TURNER; S.C. 1988. The reproductive biology of tilefish *Lopholatilus chamaeleonticeps* Goode and bean, From the united states mid Atlantic bight and the effect of fishing on the breeding system. *Fishery Bulletin*, Vol 16, No, 1988

HAMSA, K.M.S.A. and M.N. KUTTY. 1972. Oxygen consumption in relation to Spontaneous activity and ambient oxygen in five marine teleosts. *Indian J. Fish.*, 19: 76-85

HARDER - JONES, F.R. 1968. Fish Migration Edward Arnold, London : 325pp.

HICKLING, C.F. 1940. The fecundity of the herring of the Southern North Sea. J. Mar. Biol. Association U.F. 24 : (2) 619-632.

HICKLING, C.F. and E. RUTHENBERG 1936. The ovary as an indicator of spawning of fishes. J. Mar. Biol. Association U.K., 21: 311-317

\*HILDE BRAND, S.F., and L.E. CABLE. 1938. Further notes on the development and Life history of Some teleosts at Beaufort, N.C.U.S. Bur. Fish. Bull., 48: 505-642

HOAR, W.S. 1957. The gonads and reproduction In : The Physiology of fishes (Ed. M.E. Brown) Academic press, New york, pp, 287-321

HOAR, W.S. 1959. Endocrme factors in the ecological adaptation of fishes. In "Comparitive Endocrinology" (A. Gorbman, ed) 1-23, Wiley, New York. HOAR, W.S. 1965. Comparative physiology : Hormones and reproduction in Fishes. A. Rev. Physiol., 27: 51-70

HOAR, W.S. 1966. Hormonal activities of the pars distalis in Cyclostomes fish and amphibia. In "Pitutary Gland" (G.W. Harris and B.T. Donovan, eds). Vol.1 : 242-294. Butter worth, London, Washington, D.C.

HOAR, W.S. and RANDALL, D.J. (ed); 1969. Fish Physiology -volume 111 Academic Press, New york, 485 p

HONSTON, A.H. 1982. Thermal effects upon Fishes. Nat. Res. Can. Can. Assoc. Comm. On Sci. Crit. For Enviorn. Qual. 200p

HOWARD, G.V. AND A. LANDA, 1958. A study of the age, growth, sexual maturity, and spwning of the androveta (Cetengsanlis mystiicelim) in the Gulf of Panama. *Inter-Am-Trop. Tuna Comm.*, Bull., **2** (9) : 384-467

HUNTER, J.R., and MACEWICZ, B.J. 1985b. Rates of atresia in the ovary of captive and wild northern Anchovy, *Engraulis Mordax*. U.S. National Marine Fisheries Service Fisheries Service Fishery Belletin **83**, 119-36

HUTECHINS, B. and SWAINSTON, R. 1986. Sea fishes of Southern Australia Perth : Swainston publishing

HYNDES, G.A., POTTER, I.C. and HESP, S.A. 1996 b. Relationship between the movements, growth, age structures and reproductive biology of teleosts Sillago burrus and Sillago vittata in temperate Marine waters. Marine biology 126, 549-558 In. G.W. Potts and R.J.Wootton (Eds). Fish reproductionstrategiesand tactics. Academicpress, London, pp. 187-205

JAMES, P.S.B.R. 1967. The ribbon fishes of the Family Trichiuridae of India : 226p. Memoir 1. Madpam Camp : *Marine Biological Association of India* 

JAYABALAN, N. 1988. Reproductive biology of the ponyfish, *Gazza minuta* (Bloch) from Porto Novo, East Cost of India. *Indian Journal of Marine Sciences* Vol .17. March 1998, pp 51-54.

JAYASANKAR, P., 1989. Studies on the reproduction of Indian whiting Sillago Sihama (Forskal) (Percoidei, Sillaginidae). Ph.D. Thesis, Cochin University of Science and Technology, 251 p

JHINGRAN, V.G. and NATARAJAN, A.V. 1969. A study of Fisheries and fish populations of Chilka lake during the period 1957-65 *Journal if Inland Fish Society India* **49**-126

JINGRAN, V.G. and A.V. NATARAJA (1996). Final report on fisheries of chilka lake (1957-65) Bull. Cont. Inl. Fish. Res. Inst. Barrackpore (8).

JOB, T.J. 1940 An investigation on the nutrition of the perches of Madras coast. Rec. Indian. Mus., 42: 289-364. JOHANNES, R.E. 1978. Reproductive strategies of coastal marine fishes in the tropics. *Enviiron. Biol. Fishes*, **3**(1): 65-84

JONES, S and K.H. SUJANSINGANI, 1954. Fish and Fisheries of Chilka lake with statistics of fish catches for the years 1948-1950 *Indian J. Fish.*, 1 (1 and 2) 256-344

JONES, S. And M. KUMARAN, 1980. Fishes of Laccadive Archipelago. The Nature Conservation and Aquatic Science Services, Trivandrum, 757pp

\*KALIYAMURTHY, M., 1982. Studies on the percoid Fishes of the Lake Pulicat, South east loast of India. *Ph.D Thesis Magadh University, Bodh. Gaya* 

KARLOU-RIGA, -C. : ECONOMIDIS, -P.S. 1996. Ovarian attretic rates and sexual maturity of European horse mackeral, Trachurus trachurus (L.), in the Saronika Gulf (Greece). *Fish. -Bull. 1996* Vol. **94**, No.1, pp. 66-76

KERSHNER, B.A., PETERSON, M.S. and GILMORE, R.G. 1985. Ecotopic and ontogenetic trophic variation in mojarras (Pisces. Gerreidae). *Estuaries* 8, 311-322

KESTEVEN, G.L. (Ed) 1960. Mannual of field methods in fisheries biology. FAO . Mammals in Fisheries Science 1, 1-152. KLIMA, E.F. 1959. Aspects of biology and Fishery for Spanish Mackeral, Scomberomorus maculatus (Mitchill) of Southern Florida. Fla. Brd. Conserv. Mar. Res. Lab., Tech. Ser. 39p

KOYA, -Y ; HAMATSU, -T; MATSUBARA, -T, 1995. Annual reproductive cycle and spawning charecteristics of the female kichiji rock fish. *Sebastolobus macrochir. Fish, -Sci. 1995* **61,** .2 pp. 203-208

KURUP, B.M. and SAMUEL, C.T. 1991. Spawning biology of Gerres filamentosus Cuvier in the Cochin estuary. Fishery Technology 28, 19-24.

LAM, T.J., 1983. Enviornmental influence on gonadl activity in fish In: Haar, et al. (Eds). Fish physiology, Vol.IX : Reproduction, Part B, pp 65-116. (Academic press, Newyork)

LAMBERT, T.C. and WARE, D.M. 1984. Reproductive strategies of dimersal and pelafic spawning fish. *Canadian Journal of Fisheries and Aquatic Science* **41**, 1564-1569

LAURENSON, L.J.B., NEIRA, F.J. and POTTER, I.C. 1993B. Reproductive biology and larval morphology of marine plotosid *enidoglanis macrocephalus* (Teleostei) in a seasonally closed Australian estuary. *Hydrobiologia* **268**, 179-192

LECREN, E.D. 1951. The length weight relationship and sesonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). Journal of Animal Ecology **20**; 201-19

LEMOS, PH DE B; CORREA, MFM; ABILHOA, V. 1993. Guide to the otoliths of Gerreidae (Osteichthys - Perciformes) of the coast of Parana state, Brazil

LIMA, -R.V.A; BERNARDINO, -G; Vol-SELLA -M.V; MORAES, -F.F; SCHEMY, -R.A; BORELLA, -M.J. 1991. Ovarian germinative tissue and reproductive cycle of pacu (*Piaractus mesopotamicus* Holmberg, 1887) reared in captivity. *Bol*, -*Tec.* -*Cepta 1991.* **4**, 1, pp. 1-46

LOPES, -R.A.; SALA, -M.A.; LEME - DOS - SANTOS, -H.S. ; NUT -SOBRINHO, -A. ; PAULA -LOPES, -O.V. 1991. Oocyte development in the piranha *pygolentius nattereri* Knerr 1860 (Pisces, characidae). *Rev, - Biol, -Trop. 1991.* **39**, 1, pp. 7-14. *Marine Ecology Progress Series.* 

MATHESON, R.E. Jr; GILMORE, R.G. Jr. Mojarras (Pisces Gerreidae) of Indian river Lagoon. Bull. Mar. Sci., 57, 1., pp 281-285, 1995

MC EVOY, L.A. and MC EVOY, J. 1992 . Multiple spawning in Several Commercial Fish species and its consequence for fisheries management, cultivation and experimentation. *Journal of Fish. Biol.* **41** (Suppl.B), 125-136

\*MEDOZA, N.A. 1968. Consideraciones sobre la biologica perquera de la sierra. Scomberomorus Maculatus (Mitchill) en el Estado de Veracruz. Bios 1 : 11-22.

MELO, -Y. C. 1994. Multiple spawning of the Anchovy Engraulis capensis. S. Afr. J. Mar. Sci. S. Afr. Typske, Seewet 1994. 14.

MICALE, -V.; GENOVESE, -L.; GRECO, -S., PARDICHIZZI, -F. 1993. Aspects of reproductive biology of the amberjack, *seriola dumerili* (Risso 1810)

MUNRO, I.S.R. 1955. The marine and Fresh water Fishes of Cylon. Halstead press Sydney. 349 pp.

MURAYAMA, -T; MITANI, -1.; AOKI, -1. 1995. Estimation of the spawning period of the Pacific mackerel *Scomber japonicus* based on the changes in gonad index and the ovarian histology. *Bull. - Jap. - Soc. - Fish, Eng., Ebidai, Kashima, Jbaraki* 314-04,

MUTHUKARUPPAN, S., 1987. Biochemical Aspects of Ovarian maturation in Liza Parsia (Hamilton- Buchanan). M.Sc. Thesis, Cochin Univ. Of Sci. And Technol., Cochin.

NAIR, R.V., 1952. Studies on some post larval fishes of the Madras plankton. Proc. Indian Acad. Sci., 35 (5) : 225-244 NATARAJAN, R, and P.S. REDDY, 1980. Reproduction of *Liza dussumiery* (Val.) in Porto Novo waters. *Proceedings of Indian National Science Academy*, **46** : 96-104

NELSON, J.S. 1994. Fishes of the world, New york : John Wiley

NIKOLSKII, G.V., 1963. The Ecology of Fishes, Academic press, Newyork, 352 p

PATNAIK, S. 1971. Observations on the Fishery and Biology of Chilka jagili, Gerres Setifer (Hamilton). J. Inland. Fish. Soc. India. 3, 25

PILLAI, T.V.R., 1954. The biology of grey mullet, *Mugil* trade forskal, with notes on the fishery in Bengal. *Proceedings of the National Institute of Science*, *India*, **20**: 201-257

PRABHU. M.S., 1956. Maturation of intra ovarian eggs and spawning periodicifies in some fishes. *Indian. J. Fish.*, **3**: 59-90

QUASIM, S.Z., 1966. Sex ratio in fish population as a function of sexual difference in growh rate. *Current Science*, **35**; 140-142

QUAZIM, S.Z. 1955. Time and duration of spawning season of teleosts in relation to theri distribution *J. Cons. Int. Explor. Mar*, **21** (2) : 144-155 RAO, AVP. 1968. Obsevations on the food and feeding habits of *Gerres Oyens* (Forskal) and *Gerres filamentosus* cuvier from the pulicat lake with notes on food of allied species.

ROBB, A.P. 1982. Histological observations on the reproductive biology of haddock *Melanogrammus aeglefinus* (L) *J. Fish. Biol. 1982* 20, 397-408

ROY, J.C. and SAHOO, N., 1962. Bulletin on the development of chilka lake (chilka fisheries, 1949-1955) Orissa Government Press, (5)

S. - Afr. - J. -Mar. - Sci. - S. - Afr, - Typske. - Seewet. 1994. VOL.14,

SAROJINI, K.K., 1957. Biology and Fishery of grey mullets of Bengal 1. Biology of Mugil Parsia (Hamilton) Indian Journal of Fisheries, 4: 160-207.

SARRE, G.A., HYNDES, G.A. and POTTER, J.E. 1997. Habital, reproductive biology and size Composition of *Parequala melbournensis* a gerreid with a temperate distribution. *Journal of fish Biology* 1997 **50**, 341-357

Seasonal and areal changes in standing stock of phytoplankton, Zooplankton and micronekton in the eastern tropical pacific. *Mar. Bio.*, **7** (1) : 14-31

SHAEFER, M.K. 1987. Reproductive biology of black skipjack. *Euthymnus lineatus* an Eastern Pacific Tuna. *Inter - An. Trop. Tuna. Comm. Bull.* **19** (2): (169-260)

SIMPSON, J.G. 1959. Identification of the egg, early life history and spawning areas of anchovata *Celengrauis Mysticetes* (Gunther) in the Gulf of Panama, *Inter* - *Am-Top. Tuna. Comm., Bull.*, 3 (10) : 437-580.

SMITH, J.L.B., 1953. Ses fishes of South Africa, Central News Agencies Ltd., S. Africa

SMITH, M.M. and HEEMSTRA, P.C. 1986. Smith's seafishes, Johannes burg: Macmillas

TAMPI., P.R.S. 1957. Some observation on the reproduction of milk fish, *Chanos chanos* (Forskal). *Proc. Indian. Acad. Sci.*, **46** : B: 154-273.

VARGHESE, T.J. 1961. Observations on the biology of Raconda. Indian. J. Fish. 8 96-106

VARGHESE, T.J. 1973. The Fecundity of rohu, Labeo rohita (Hamilton) Proc. Indian. Acad. Sci. 77: 214-224

VARGHESE, T.J. 1976. Studies on the fecundity of *Coilia ramcarati* (Ham.Buch). *Proc. Indian. Acad. Sci.* 83 : 47-54

VASUDEVAPPA, C. AND P.S.B.R. JAMES, 1978. Studies on the maturity and spawnng of the marine Cat-fish *Tachysaurus dussumeri* (Valenciennes) along the South Kanara coast. *Summer school an Fshery biology held at Magalore* 10-14 April 1978 WEBER, M. And L.F. de BEAFORT, 1931. Fishes of Indo Australian Archipelazo, Vol.6, Leiden

WEST, G. 1990. Methods of assessing ovarian development in fishes : a review Aus. Jau. Mar and F.W. Res. 41 : 199-222

WLSON, -C.A: NELAND, -D.L. 1994. Reproductve biology of red drun, Sciienops occelatus, from the nertic waters of the northern Gulf of Mexico Fish. -Bull. 1994.. 92, 4, pp 841-850

\* Not referred in orginal.