

Compound eye as a target organ of white spot baculovirus (WSBV) in *Penaeus indicus* H. Milne Edwards

JASMIN K. JOSE AND MARY K. MANISSERI

Central Marine Fisheries Research Institute, Cochin-682 014, India

ABSTRACT

Recently mass mortalities of penaeid prawns have been reported worldwide in the culture ponds due to the occurrence of White Spot Disease (WSD). Eventhough, several tissues of ectodermal and mesodermal origin have been confirmed to be the target organs for white spot baculovirus, not much importance has been given so far to the compound eyes of penaeid prawns affected with WSD. Histopathological and ultrastructural studies were conducted in the compound eye of *Penaeus indicus* H. Milne Edwards, affected with WSD. Eosinophilic to basophilic intranuclear inclusion bodies were observed in the narrow region of the reticular cell nuclei in the dioptic portion of the ommatidia. Electron microscopic examination of the compound eye revealed the presence of rod-shaped, enveloped, non-occluded virions and viral particles in the affected nuclei of reticular cells.

Introduction

White Spot Syndrome Baculovirus (WSBV) is a recently discovered group of crustacean viruses (Lightner, 1996; Nunan and Lightner, 1997). They can result in nearly 100% mortality within 4 to 6 days, because of their high pathogenicity. The rapid onset, lethality, extreme virulence and wide host range of WSBV are remarkable (Chou *et al.*, 1995, 1998; Chang *et al.*, 1996; Lo *et al.*, 1997). WSBV has been reported to infect many penaeid species and other arthropods (Lo *et al.*, 1996b; Momoyama *et al.*, 1997; Wang *et al.*, 1997; Kasornchandra *et al.*, 1998). The disease is characterized by the presence of

white spots on the exoskeleton (Chou *et al.*, 1995; Wang *et al.*, 1995; Lightner 1996). Histo-pathological studies have demonstrated that WSBV targeted various tissues of ectodermal and mesodermal origin (Momoyama *et al.*, 1994; Chou *et al.*, 1995; Wang *et al.*, 1995; Chang *et al.*, 1996; Durand *et al.*, 1997; Lo *et al.*, 1997; Lo and Kou, 1998). WSBV infections are characterized by hypertrophied nuclei with prominent intranuclear eosinophilic to basophilic inclusion bodies in the cells of cuticular epidermis (hypodermis) of the appendages such as pleopods and pereopods, eyestalk, general body, subcuticular epithelium of foregut and hindgut, epithelial cells of

gills, various connective tissues, antennal gland, lymphoidal organ, hematopoietic tissues, gonads, nervous tissues, hemocytes, heart muscles, striated muscles and in a variety of other cell types (Momoyama *et al.*, 1994, 1997; WongteerasupayaeJaZ., 1995; Durandei *al.*, 1997; Wang *et al.*, 1997; Kasornchandra *et al.*, 1998; Lightner *et al.*, 1998; Nunan *et al.*, 1998). Ultrastructural studies conducted in these tissues by the above authors and others such as, Nakano *et al.* (1994), Takahashi *et al.* (1994) and Lo *et al.* (1996 a) also confirmed the presence of rod-shaped, enveloped and non-occluded baculovirus. The present study describes the occurrence of WSBV, histopathological and ultrastructural changes in the compound eye of *Penaeus indicus* affected with WSD, on which no detailed study has been conducted.

Materials and methods

Live specimens of *Penaeus indicus*, 85 to 110 mm in length, affected with WSD were collected from the prawn culture ponds at Cochin and their compound eyes were processed for histopathological and ultrastructural studies. For histopathology, the material was fixed in Davidson's fixative, processed, embedded, sectioned (thickness 5-6µm) stained with haematoxylin and eosin according to the procedure outlined by Bell and Lightner (1988). The tissues were examined for histomorphological changes.

For transmission electron microscopic studies, compound eyes of live, WSD affected *P. indicus* were diced in 4%, buffered gluteraldehyde and post fixed in 1% osmium tetroxide. Fixed tissues were processed and embedded in Spurr's resin (Spurr, 1969). Ultrathin sections were cut with an ultratome of model Leica Ultracut R, stained with uranyl acetate and lead citrate (Reynolds, 1963). The sections were examined using a transmission

electron microscope (Hitachi H-600) at 50KV.

Results and discussion

P. indicus specimens, affected with White Spot Disease exhibited symptoms, such as small, obscure white spots of 0.1 to 3 mm size on the inner surface of the cuticle, especially in the cephalothoracic region, fifth and sixth abdominal segments. Areddish to pinkish colouration of the cephalothoracic region was the initial symptom and the infected prawns stopped feeding and moved to the edges of the ponds, became sluggish, lethargic and showed no escape reactions. In almost all of them, a part of the antennae was lost and their eyes became pale, loosing the natural glow. Similar symptoms were reported in *P. monodon* and *P. japonicus* by WongteerasupayaetaZ. (1995), Durand *et al.* (1997), Lo and Kou (1998) and Takahashi *etal.*, (1998).

Light microscopic observations of L.S. of the compound eye from healthy *P. indicus* revealed the ommatidium to be composed of two regions, the peripheral dioptic portion and the innermost ganglionic region. A major part of the dioptic region of the ommatidia is composed of crystalline tracts, which formed the distal lightly staining area, seen below the crystalline cone region as seen in Fig.1. A narrow layer of reticular cell nuclei could be observed lying in close contact with the crystalline tracts. The nuclei of reticular cells lie distal to the cell bodies and are concentrated in this narrow layer. In the nuclei of the reticular cells, nucleolus and nuclear membrane are very clearly distinguished as in a typical cell (Fig. 1). A basement membrane divide the dioptic portion of the ommatidium from the inner ganglionic region. Below this basement membrane is the fasciculated zone, which contain primary optic nerve fibres. Similar histological details are given by Bell and Lightner (1988) for the normal

compound eye of penaeid prawns.

gill epithelium, **cuticular** epidermis, stomach, **haemocytes**, hematopoietic tissues, lymphoid organ, connective tissues, muscles and nerves of *P. monodon*, *P. vannamei*, *P. stylirostris*, *P. japonicus* and *Metapenaeus ensis* affected with **WSD**. So far no work has been reported on the occurrence of WSBV in the compound eyes of **penaeid** prawn except the histopathological and **PCR observations** made by Chang *et al.* (1998), Lo *et al.* (1997) and Lo and **Kou** (1998) in the cuticular epidermis of the **eyestalks** of *P. monodon*.

The present investigation showed that both the nucleolus and the nuclear membrane of the affected nuclei were associated (Fig. 2). In some hypertrophied nuclei, chromatin was clear and the marginated, dense chromatin was seen as a ring like dark inclusion body in the nucleus. In the early stage of infection, small, eosinophilic inclusion bodies could be observed in the nuclei and a clear zone was seen in the inclusion body and chromatin. In the late stage of infection, the inclusion body was highly basophilic, occupying the nucleus and the transparent zone was no longer present. Similar observations were made by Wang *et al.* (1995) in the gills of *P. monodon* and Systemic Ectodermal and Hemolymphic **Baculovirus (SEMBV)**. Wang *et al.* (1998) opined that nuclear inclusions of the infected cells is due to the presence of **inclusion bodies** and accumulation of virions. The size of the hypertrophied nuclei ranged from 5 to 14 μ m in diameter. Wang *et al.* (1997b) reported a size range of 10 to 15 μ m for the infected nuclei of cuticular epidermis in *P. monodon* and *P. japonicus*. A few cells in the fasciculated zone and the innermost ganglionic region also exhibited signs of **WSD**. But the maximum concentration of

the hypertrophied nuclei of the reticular cells in the dioptric region of the ommatidia. Wongteerasupaya *et al.* (1995), Barmlet *et al.* (1997), Wang *et al.* (1997), Takahashi *et al.* (1998) and Rajendran *et al.* (1999) also reported such inclusion bodies in the

p 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000

i . e . h

i.:(KHtx

Acknowledgements

The authors are thankful to Dr. G. Sudhakara Rao, Principal Scientist, Dr. KRangarajan, Dr K.C. George, Senior Scientists, and Shri. N.K. Sanil, Scientist, CMFRI for their valuable help and encouragement.

References

- Bell, T.A. and D.V. Lightner 1988. *A handbook of normal penaeid shrimp histology*. WorldAquaculture Society, Baton Rouge, Louisiana, USA, 114 pp.
- Chang, P.S., C.F. Lo, Y.C. Wang and G.H. Kou 1996. Identification of White Spot Syndrome associated baculovirus (WSBV) target organs in the shrimp, *Penaeus monodon* by *in situ* hybridization. *Dis. Aquat. Org.*, 27(2): 131-139.
- Chou, H.Y., C.Y. Huang, C.H. Wang, H.C. Chiang and C.F. Lo 1995. Pathogenicity of a baculovirus infection causing White Spot Syndrome in cultured penaeid shrimp in Taiwan. *Dis. Aquat. Org.*, 23(3) : 165-173.
- Durand, S., D.V. Lightner, L.M. Nuan, R.M. Redman, J. Mari and J.R. Bonami 1996. Application of gene probes as diagnostic tools for White Spot Baculovirus (WSBV) of penaeid shrimp. *Dis. Aquat. Org.*, 27(1): 59-66.
- Durand, S., D.V. Lightner, R.M. Redman and J.R. Bonami 1997. Ultrastructure and morphogenesis of White Spot Syndrome Baculovirus (WSSBV). *Dis. Aquat. Org.*, 29(3) : 205-211.
- Kasornchandra, J., S. Boonyaratpalin and T. Itami 1998. Detection of White Spot Syndrome in cultured penaeid shrimp in Asia; Microscopic observation and polymerase chain reaction. *Aquacult.*, 164(1-4) : 243-251.
- Lightner, D.V, K.W. Hasson, B.L. White and R.M. Redman 1998. Experimental infection of western hemisphere penaeid shrimp with Asian White Spot Syndrome virus and Asian yellow head virus. *J. Aquat. Anim. Heal.*, 10(3): 271-281.
- Lo, C.F. and G.H. Kou 1998. Virus associated White Spot Syndrome of shrimp in Taiwan : A review. *Fish Pathol.*, 33(4): 365-371.
- Lo, C.F, J.H. Leu, C.H. Ho, C.H. Chen, S.E. Peng, Y.T. Chen, C.M. Chou, P.Y. Yeh, C.J. Huang, H.Y. Chou, C.H. Wang and G.H. Kou 1996a. Detection of Baculovirus associated with White Spot Syndrome (WSBV) in penaeid shrimps using polymerase chain reaction, *Dis. Aquat. Org.*, 25(1-2): 133-141.
- Lo, C. R, C.H. Ho, S.E. Peng, C.H. Chen, H.C. Hsu, Y.L. Chiu, C.F. Chang, K.F. Liu, M.S.Su, C.H. Wang and G.H. Kou 1996b. White Spot Syndrome Baculovirus (WSBV) detected in cultured and captured shrimp, crabs and other arthropods. *Dis. Aquat. Org.*, 27(3): 215-225.
- Lo, C.F, C.H. Ho, C.H. Chen, K.H. Liu, Y.L. Chiu, P.Y. Yeh, S.E. Peng, H.C. Hsu, H.C. Liu, C.F. Chang, M.S. Su, C.H. Wang and G.H. Kou 1997. Detection and tissue tropism of White Spot Syndrome Baculovirus (WSBV) in captured brooders of *Penaeus monodon* with a special emphasis on reproductive organs. *Dis. Aquat. Org.*, 30(1): 53-72.
- Momoyama, K., M. Hiraoka, H. Nakano, H. Koube, K. Inouye and N. Oseka 1994. Mass mortalities of cultured kuruma shrimp, *Penaeus japonicus*, in Japan in 1993: Histopathological study. *Fish pathol.*, 29: 141-148.
- Momoyama, K., M. Hiraoka, K. Inouye, T. Kimura, H. Nakano and M. Yasui 1997. Mass mortalities in the production of juvenile greasyback shrimp, *Metapenaeus ensis*, caused by Penaeid Acute Viremia (PAV). *Fish pathol.*, 32(1): 51-58.
- Nakano, H., H. Koube, S. Umezawa, K. Momoyama, M. Hiraoka, K. Inouye and N. Oseko 1994. Mass mortalities of cultured kuruma shrimp, *Penaeus japonicus* in Japan in 1993: Epizootiological survey and infection trials. *Fish pathol.*, 29 : 135-139.
- Nunan, L.M. and D.V. Lightner 1997. Development of a non radioactive gene

- probe by PCR for detection of White Spot Syndrome Virus (WSSV). *J. Virol. Meth.*, 63 : 193-201.
- Nunan, L.M., B.T.Poulos and D.V. Lightner 1998. The detection of White Spot Syndrome Virus (WSSV) and Yellow Head Virus (YHV) in imported commodity shrimp. *Aquacult.*, 160 (1-2) : 19-30.
- Rajendran, K.V., K.K. Vijayan, T.C. Santiago and R.M.Krol 1999. Experimental host range and histopathology of White Spot Syndrome Virus (WSSV) infection in shrimp, prawns, crabs and lobsters from India. *J. Fish Dis.*, 22 :183-191.
- Reynolds, E.S. 1963. The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *J. Cell Biol.*, 17 : 208-212.
- Shaw, S.R. and S. Stowe 1982. Photoreception. In : *The Biology of Crustacea. Neurobiology: Structure and function.* 2: 291-367, D.E. Bliss, H.L. Atwood and D.C. Sandeman (Eds.), Academic Press, New York, London.
- Squerr, A.S. 1969. A low-viscosity epoxy resin embedding medium for electron microscopy. *J. Ultrastruct. Res.*, 26 : 31-43.
- Takahashi, Y., T. Itami, M.Kondo, M. Maeda, R. Fijii, S.Tomonaga, K. Supamattaya and S. Boonyaratpalin 1994. Electron microscopic evidence of bacilliform virus infection in kuruma shrimp, *Penaeus japonicus*. *Fishpathol.*, 29(2): 121-125.
- Takahashi, Y, T. Itami, M. Maeda and M.Kondo 1998. Bacterial and viral diseases of kuruma shrimp, *Penaeus japonicus* in Japan. *Fish pathol.*, 33(4): 357-361.
- Wang, C.H., C.F. Lo, J.H. Leu, C.M.Chou, P.Y.Yeh, H.Y. Chou, M.C.Tung, C.F.Chang, M.S.Su and G.h.Kou 1995. Purification and genomic analysis of Baculovirus associated with White Spot Syndrome (WSBV) of *Penaeus monodon*. *Dis. Aquat. Org.*, 23(3): 239-242.
- Wang, C.S., K.F.J.Tang, G.H.Kou and S.N.Chen 1997. Light and electron microscopic evidence of White Spot Disease in the giant tiger shrimp, *Penaeus monodon* (Fabricius) and the kuruma shrimp, *Penaeus japonicus* (Bate), cultured in Taiwan. *J.Fish dis.*, 20 : 323-331.
- Wongteerasupaya, C, J.E.Vickers, S.Sriurairatana, G.L. Nash, A.Akarajamorn, V.Boonsaeng, S. Panyim, A.Tassanakajon, B.Withyachumnarnkul and T.W.Flegel 1995a. Anon occluded systemic baculovirus that occurs in cells of ectodermal and mesodermal origin and causes high mortality in the black tiger prawn, *Penaeus monodon*. *Dis. Aquat. Org.*, 21(1) : 69-77.