

Benthic communities of mesophotic coral ecosystem off Puducherry, east coast of India

P. Laxmilatha^{1,*}, S. Jasmine², Miriam Paul Sreeram³ and Periasamy Rengaiyan¹

¹ICAR-Madras Research Centre of the Central Marine Fisheries Research Institute, Chennai 600 028, India

²ICAR-Vizhinjam Research Centre of the Central Marine Fisheries Research Institute, Cochin 695 521, India

³ICAR-Central Marine Fisheries Research Institute, Cochin 682 018, India

The shallow coral reef ecosystems along the Indian coast are being threatened by anthropogenic global ocean warming and increased frequency of coral bleaching in the recent past. Identification and conservation of deeper reef habitats are essential as they serve as a source of larvae and livestock to replenish the shallow reef habitats. Information on the location and spatial extent of the mesophotic coral ecosystems (MCEs) and their biodiversity is poorly known in the continental shelf of the east coast of India. In this study, we have documented the species diversity of MCEs at a depth of 30–40 m off Puducherry along the east coast of India. In total, 12 species of corals, including 5 black corals and 16 octocorals, 4 species of sponges and 31 species of coral-associated benthopelagic fish species were recorded. *Subergorgia* sp. was the most dominant species of octocorals and found extensively as gorgonian forests. The MCEs reported in this study raise important questions about the origin and connectivity of the coral populations in this region to the other major coral reef ecosystems along the east coast. Understanding the physical processes and hydrographic features around the MCEs, off Puducherry will reveal more information about the distribution and colonization of coral communities and their vulnerability to changes in future.

Keywords: Benthic communities, gorgonian forest, mesophotic coral ecosystems, reef habitats.

MESOPHOTIC coral ecosystems (MCEs) are highly productive and stable, providing a variety of habitats to a wide range of benthic organisms. MCEs are characterized by low solar irradiance with blue spectral enrichment, low wave disturbance, reduced temperatures and rich nutrients-upwelled water^{1,2}. Light is the primary physical parameter that limits the distribution of coral reefs across depths and habitats. The sharp differences in the irradiance over depth gradient, on spatial scales of 10 m have major consequences for the distribution of corals. Photosynthetic corals are often found at depths ranging from 0

to 150 m in clear waters of the tropics³. However, several species of corals interface between the shallow and deep sea environments around the world^{1,2,4,5}. In general, MCEs occur at a depth of 30–150 m of the euphotic zone in tropical regions^{4,5}. The MCEs, situated off Puducherry the east coast of India are considered to be unique, showing all the features mentioned above.

Biotic assemblages in MCEs are considered to be extensions of shallow-water coral ecosystem assemblages due to their unique depth range⁴. In addition, a number of unique or depth-restricted species occur in these habitats. These diverse groups of benthic assemblage in MCEs hold a rich biodiversity and chemical reservoirs² which are of high, ecological and economic value⁶. Knowledge on MCEs is scarce, while little is known about their functional aspects. Advanced technologies may enable us to connect and overcome the logistic challenges to bridge this knowledge gap and allow resource managers to make informed decisions on conserving MCEs². The most important knowledge gap pertains to the species diversity and ecological role of organisms in MCEs. While intensive efforts were taken to study MCEs in different parts of the world, we do not have concerted efforts to study MCEs in India.

The potential importance of MCEs as refugia for shallow coral ecosystems and the rich source of novel compounds prompted scientists and managers to study them extensively. MCEs are well described in the western Atlantic and reported in the Pacific and Indian Oceans⁷. In India, MCEs and their associated fauna have been described in the Angria Bank along the west coast (pers. commun.). The shallow coral ecosystems are exposed to a range of threats, especially high sea temperature and solar radiation that cause coral bleaching and lead to mass mortality of corals⁸. In contrast, MCEs are protected from these threats due to greater depth of the overlying water column and reduced light irradiance⁹. Most importantly, MCEs serve as the source of coral larvae and other important marine organisms to the shallow reef environments, enabling them to recover post disturbances. Therefore, the conservation of MCEs is critical for the persistence of corals and their associated fauna under the

*For correspondence. (e-mail: laxmil@yahoo.com)

futuristic climate scenarios. Recent studies on MCEs in Japan have reviewed the knowledge on MCEs in different domains such as their distribution, biodiversity, occurrence in fossil records and future directions¹⁰. Considering the above factors we had taken up a short-term study of MCEs, off Puducherry, to understand the biodiversity of corals and associated fauna. Results of this study enrich the knowledge of biodiversity of MCEs in India and also serve as a baseline for evaluating the futuristic changes.

Materials and methods

Study area

The MCEs, situated off Puducherry in the east coast of India, are composed of a hard substratum with corals and associated biota. MCEs were located on a raised platform or wall which is 2.5 km long and 0.05 km wide, covering an area of ~12.5 ha (0.125 km²). The width of the wall was ~7 m in the depth range between ~30 and 40 m (Figure 1). The MCEs comprise patchy coral habitats dominated by gorgonians. *In situ* observations on species diversity were carried out on a random transect by scuba diving during February and March 2017 at ten stations along the wall (Table 1). The depth of the sampling stations ranged from 30 to 40 m. The organisms encountered under the transect were photographed using a mounted digital camera (Nikon D7000 DSLR with AF-S 18–105 mm VR Kit Lens) bearing a flash at a depth range of 33–40 m. Fishes were collected for identification using hooks and bait, and from the local fishermen. Nudi-branches were collected manually during *in situ* observations. Photographs taken *in situ* and *ex situ* were used to differentiate morphologically similar species.




Figure 1. Location of mesophotic coral ecosystems, including the present study site on the east coast of India.

Different species were identified using high-resolution photographs and videos following the keys described by Huang *et al.*¹¹ for coral species belonging to families Merulinidae, Montastraeidae and Diploastraeidae; Cairns¹² and Kitahara¹³ for azooxanthellate scleractinian; Laborel¹⁴ and Neves *et al.*¹⁵ for zooxanthellate scleractinian and Castro *et al.*¹⁶ and Pérez *et al.*¹⁷ for octocorals. Soft corals and sea fans were identified at the genus level following Fabricius and Alderslade¹⁸ and Antony Fernando¹⁹. Fishes were identified using the keys described by Smith²⁰ and FishBase²¹. The flatworms were identified following Newman and Cannon²², and sea snake by Rasmussen²³.

Results and discussion

Depth

The depth of MCEs observed in the present study ranged from 30 to 40 m. Comparatively, this is the lowest depth so far reported. Globally, the occurrence of MCEs below 50 m depth was reported from US Virgin Islands²⁴, Bermuda²⁵, the Caribbean (from 50 to 100)²⁶, Puerto Rico^{27,28}, Australia²⁹, Bahamas³⁰, Brazil³¹, Japan's Ryukyu archipelago (above 100 m)³² and Hawaii³³. The MCEs off Puducherry possess a variety of organisms composed of at least 77 species from 41 families. Table 2 provides detailed list of species recorded from visual surveys in MCEs.

Corals

The hard bottom of the shelf margin was dominated by slow-growing plate coral *Leptoseris explanata* Yabe & Sugiyama, 1941 and *Pachyseris speciosa* (Dana, 1846) (Figure 2). Rooney *et al.*³⁴ have reported *Leptoseris* spp. as restricted to extreme low-irradiance environments such as MCEs. Luck *et al.*³⁵ noted that the distribution of *Leptoseris papyracea* and *L. hawaiiensis* was restricted to deeper waters (more than 100 m) and habitats under shady conditions such as crevices, caves and overhangs^{36–38}.

Table 1. Details of scuba diving location

Station	Latitude	Longitude	Depth
S1	11°48.119'	79°55.403'	32.5
S2	11°48.202'	79°55.395'	32.5
S3	11°48.287'	79°55.389'	32.6
S4	11°48.362'	79°55.419'	33.0
S5	11°48.432'	79°55.426'	33.4
S6	11°48.470'	79°55.429'	34.1
S7	11°48.638'	79°55.428'	34.6
S8	11°48.967'	79°55.453'	34.7
S9	11°49.087'	79°55.431'	34.2
S10	11°49.404'	79°55.412'	34.4

Table 2. List of mesophotic corals and associated species

Family	Species	Family	Species
Algae		Others	
Schizymeniaceae	<i>Titanophora pikeana</i> (Dickie) Feldmann, 1942	Axinellidae	<i>Pipestela candelabra</i> Alvarez, Hooper & van Soest, 2008 <i>Axinella corrugate</i> (George & Wilson, 1919)
Hard coral		Dictyonellidae	<i>Acanthella acuta</i> Schmidt, 1862
Agariciidae	<i>Leptoseris explanata</i> Yabe & Sugiyama, 1941 <i>Pavona minuta</i> Wells, 1954 <i>Pavona maldivensis</i> (Gardiner, 1905)	Aplysiniidae	<i>Aplysica cauliformis</i> (Carter, 1882)
Dendrophylliidae	<i>Tubastraea micranthus</i> (Cairns and Zibrowius, 1997)	Pseudocerotidae	<i>Thysanozoon nigropapillosum</i> (Hyman, 1959)
Euphylliidae	<i>Tubastraea coccinea</i> Lesson, 1829 <i>Euphyllia ancora</i> Veron and Pichon, 1980	Phyllidiidae	<i>Phyllidia ocellata</i> Cuvier, 1804
Fungiidae	<i>Cycloseris</i> sp.	Gryphaeidae	<i>Hyotissa hyotis</i> (Linnaeus, 1758) <i>Pteria penguin</i> (Roding, 1798)
Merulinidae	<i>Hydnophora rigida</i> (Dana, 1846) <i>Goniastrea pectinata</i> (Ehrenberg, 1834) <i>Dipsastraea favus</i> (Forskål, 1775)	Trapeziidae	<i>Quadrella maculosa</i> Alcock, 1898
Psammocoridae	<i>Psammocora haimeana</i> Milne Edwards & Haime, 1851	Dromiidae	<i>Dromia dromia</i> (Linnaeus, 1763)
Scleractinia incertae sedis	<i>Pachyseris speciosa</i> (Dana, 1846)	Colobometridae	<i>Cenometra bella</i> (Hartlaub, 1890)
Black coral		Elapidae	<i>Hydrophis lapemoides</i> (Gray, 1849)
Antipathidae	<i>Cirripathes spiralis</i> (Linnaeus, 1758) <i>Pseudocirripathes mapia</i> Bo <i>et al.</i> , 2009 <i>Antipathes dendrochristos</i> Opresko, 2005 <i>Antipathes grandis</i> Verrill, 1928 <i>Cupressopathes abies</i> (Linnaeus, 1758)	Fish species	
Gorgonians		Torpedinidae	<i>Torpedo marmorata</i> Risso, 1810
Gorgoniidae	<i>Leptogorgia</i> sp. <i>Eugorgia</i> sp.	Acanthuridae	<i>Acanthurus nigricauda</i> Duncker & Mohr, 1929 <i>Acanthurus thompsoni</i> (Fowler, 1923) <i>Acanthurus xanthopterus</i> Valenciennes, 1835 <i>Pterocaesio chrysozona</i> (Cuvier, 1830) <i>Odonus niger</i> (Rüppell, 1836) <i>Caranx heberi</i> (Bennett, 1830) <i>Chaetodon decussatus</i> Cuvier, 1829 <i>Heniochus acuminatus</i> (Linnaeus, 1758) <i>Plectorhinchus vittatus</i> (Linnaeus, 1758)
Ellisellidae	<i>Ellisella</i> sp. 1 <i>Ellisella</i> sp. 2 <i>Ellisella</i> sp. 3 <i>Junceella</i> sp. <i>Dichotella gemmacea</i> <i>Viminella flagellum</i>	Holocentridae	<i>Myripristis kuntee</i> Valenciennes, 1831 <i>Myripristis botche</i> Cuvier, 1829 <i>Bodianus diana</i> (Lacepède, 1801) <i>Bodianus neilli</i> (Day, 1867) <i>Labroides dimidiatus</i> (Valenciennes, 1839) <i>Thalassoma lunare</i> (Linnaeus, 1758)
Acanthogorgiidae	<i>Alackagorgia</i> sp.	Lutjanidae	<i>Lutjanus bohar</i> (Forsskål, 1775) <i>Lutjanus russellii</i> (Bleeker, 1849)
Acanthogorgiidae	<i>Muricella</i> sp. <i>Acanthogorgia</i> sp.	Nemipteridae	<i>Scolopsis vosmeri</i> (Bloch, 1792)
Subergorgiidae	<i>Subergorgia</i> sp.	Pomacanthidae	<i>Apolemichthys xanthotis</i> (Fraser-Brunner, 1950) <i>Pomacanthus semicirculatus</i> (Cuvier, 1831) <i>Pomacanthus imperator</i> (Bloch, 1787) <i>Chromis dimidiata</i> (Klunzinger, 1871) <i>Chromis viridis</i> (Cuvier, 1830) <i>Chrysiptera rollandi</i> (Whitley, 1961) <i>Dascyllus aruanus</i> (Linnaeus, 1758) <i>Neopomacentrus cyanomos</i> (Bleeker, 1856) <i>Neopomacentrus filamentosus</i> (Macleay, 1882) <i>Epinephelus malabaricus</i> (Bloch & Schneider, 1801)
Plexauridae	<i>Menella</i> sp. <i>Muricea</i> sp. <i>Echinogorgia</i> sp.	Pomacentridae	<i>Pterois volitans</i> (Linnaeus, 1758) <i>Torpedo marmorata</i> Risso, 1810
Melithaeidae	<i>Acabaria</i> sp.	Serranidae	
		Scorpaenidae	
		Torpedinidae	

Leaf corals were dominated by medium-to-large colonies of *Pavona minuta* Wells, 1954, and *Pavona maldivensis* (Gardiner, 1905). Other hard corals such as *Tubastraea micranthus* (Cairns and Zibrowius, 1997) and *T. coccinea* Lesson, 1829, commonly known as the black sun coral, lack zooxanthellae (symbiotic algae) and rely on zooplankton captured with their tentacles for their nutrition. These azooxanthellate corals possess a slower growth rate compared to the zooxanthellate and they are usually adapted to the environments with strong current. The azooxanthellate corals were native to the tropical Indo-Pacific, ranging from the Red Sea and Madagascar to Japan, Hawaii and Tonga (Table 2).

Six black coral species with dense populations were observed in the study area (Figure 3). Roark *et al.*³⁹ observed that the black corals are sessile benthic suspension feeders, slow-growing species, potentially indicating the oldest living animals on earth, with colonies dated to over 4000 years old. They are important components of the mesophotic benthic communities, and are considered to be ecosystem engineers, providing valuable three-dimensional habitat for numerous associated fauna, fish and invertebrate species^{40,41}. A total of 16 species of octocorals were identified belonging to two sub-orders and seven families. Photographic identification of individuals was up to genus level, except for *Dichotella gemmacea*

(Milne Edwards & Haime, 1857). *Subergorgia* sp. was dominant among octocoral communities forming gorgonian forests (Figure 3).

Sponges

The sponges are major components in most coral reef ecosystems, including MCEs. In this study we observed limited number of species, with four distinct taxa, viz. Bob Marley sponge *Pipestela candelabra* Alvarez, Hooper & van Soest, 2008; *Axinella corrugate* (George & Wilson, 1919); orange lumpy sponge *Acanthella acuta* Schmidt, 1862, and row pore rope sponge *Aplysina cauliformis* (Carter, 1882) (Figure 4).

Fishes

A total of 31 fish species belonging to 16 families were identified in the study area. Malabar grouper *Epinephelus malabaricus* (Bloch & Schneider, 1801), emperor angel-fish *Pomacanthus imperator* (Bloch, 1787) and *Neopomacentrus filamentus* (Macleay, 1882) were the most common species. Fifteen reef fish species were observed, with the dominant species being *Odonus niger* (Rüppell, 1836), also known as red-toothed triggerfish (Figure 3), and the cleaner wrasse *Labroides dimidiatus* (Valenciennes, 1839). These fish species play key ecological




Figure 3. Black coral and gorgonian species in the study area: *a*, *Cirripathas spiralis*; *b*, *Cupressopathes abies*; *c*, *Antipathes dendrochritus*; *d*, *Tubastraea micranthus*; *e*, School of fish in the gorgonian forest; *f*, *Eugorgia* sp.; *g*, *Ellisella* sp.




Figure 2. Coral species in the study area: *a*, *Euphyllia ancora*; *b*, *Psammocora haimiana*; *c*, *d*, *Leptoseris explanata*; *e*, *Dipsastraea favus*; *f*, *Montipora* sp.




Figure 4. *a*, *Pterois volitans*; *b*, feather star *Cenometra bella* on *Eugorgia* sp.; *c*, row pore rope sponge *Aplysina cauliformis*; *d*, *Thysanozoon nigropapillosum*; *e*, a discarded fishing net was smothering the gorgonids species; *f*, sponge crab *Dromia dromia* on the fishing net.

roles, and show high species diversity within MCEs^{6,42–44}. The lionfish *Pterois volitans* (Linnaeus, 1758) originally native to the Indian and Pacific Oceans and the Red Sea, was also recorded. They have rapidly spread over the past decade from their few initial sightings to colonizing the mesophotic reef habitats across the Western Atlantic^{45,46} (Figure 4).

Other taxa

The polyclad flatworms *Thysanozoon nigropapillosum* (Hyman, 1959) collected from the study area has a characteristic white margin with colourful yellow-tipped papillae. These flatworms are obvious inhabitants of coral reefs throughout tropical and subtropical oceans, and are reported from various reef environments at Maldives, Sri Lanka, Indonesia, New Guinea and Solomon Islands^{47,48} (Figure 4). The varicose wart slug *Phyllidia varicose* Lamarck, 1801, is widely distributed throughout the Indo-West Pacific Ocean, the central Pacific and the Red Sea. The honeycomb oyster *Hytissa hyotis* Linnaeus 1758 has a wide native range in the Indian and Pacific Oceans. *Pteria penguin* (Röding, 1798), commonly known as the penguin's wing oyster and native to the western and central Indo-Pacific region, was also recorded (Figure 4). The crab *Quadrella maculosa* (Alcock, 1898) was found associated with black coral *Cupressopathes abies* (Linnaeus, 1758). Another crab *Dromia dromia* (Linnaeus, 1763) was found associated with the sponges and widely distributed across the Indo-Pacific region. The feather star *Cenometra bella* (Hartlaub, 1890) was found in the study area, which is a common inhabitant of deeper reef habitats. The Arabian Gulf sea snake *Hydrophis lapemoides* (Gray, 1849) was identified by its striking patterning, comprising 33–35 dark bands along its body length. It is well-adapted for an entire life cycle in the marine environment. It feeds on eels and other types of bony fish are the main source of prey it locates amongst crevices in rocks and coral reefs⁴⁹.

Threats to MCEs

Globally, MCEs are vulnerable to environmental disturbances such as overfishing, bottom fishing gear, capture of aquarium fish, precious coral trade, land-based pollution and invasive species. The MCEs off Puducherry are not an exemption, as we observed a discarded fishing net smothering the gorgonids (Figure 4e). Fishing boats operating trawl nets were observed in the study area and such trawl operations will undoubtedly cause severe destruction to the MCEs. Therefore, strong management responses are essential to mitigate the documented threats to prevent future destruction of MCEs. The stress from fishing activities is considered by marine experts to be the greatest threat to benthic habitats⁵⁰.

Conclusion

Results of this study provide a baseline on the overall diversity of MCEs off Puducherry. The high species diversity in the MCEs might play an important role in replenishing the biodiversity of shallow-water reef ecosystems and therefore they deserve suitable conservation strategies. Given the poor knowledge on the distribution and extension of these habitats, there is a need for detailed habitat mapping in the hard bottom of the east coast of India. The present study has documented the species diversity of MCEs within this offshore hard-bottom and provides important data for further studies on fish assemblages of MCE habitats.

1. Lesser, M. P., Slattery, M. and Leichter, J. J., Ecology of mesophotic coral reefs. *J. Exp. Mar. Biol. Ecol.*, 2009, **375**, 1–8.
2. Kahng, S. E. *et al.*, Community ecology of mesophotic coral reef ecosystems. *Coral Reefs*, 2010, **29**(2), 255–275.
3. Kleypas, J. A., McManus, J. W. and Meñez, L. A., Environmental limits to coral reef development: where do we draw the line?. *Am. Zool.*, 1999, **39**, 146–159.
4. Hinderstein, L. M. *et al.*, Mesophotic coral ecosystems: characterization, ecology, and management. *Coral Reefs*, 2010, **29**, 247–251.
5. Wagner, D., Pochon, X., Irwin, L., Toonen, R. J. and Gates, R. D., Azooxanthellate? Most Hawaiian black corals contain *Symbiodinium*. *Proc. R. Soc. London Ser. B*, 2011, **278**, 1323–1328.
6. Bejarano, I., Appeldoorn, R. S. and Nemeth, M., Fishes associated with mesophotic coral ecosystems in La Parguera, Puerto Rico. *Coral Reefs*, 2014, **33**(2), 313–328.
7. Baker, E. K., Puglise, K. A. and Harris, P. T. (eds), *Mesophotic Coral Ecosystems – A Lifeboat for Coral Reefs?* The United Nations Environment Programme and GRID-Arendal, Nairobi and Arendal, 2016, p. 98.
8. Jokiel, P. L., Temperature stress and coral bleaching. In *Coral Health and Diseases* (eds Rosenberg, E. and Loya, Y.), Springer-Verlag, 2004, pp. 401–428.
9. Harris, P. T., Bridge, T. C., Beaman, R. J., Webster, J. M., Nichol, S. L. and Brooke, B. P., Submerged banks in the Great Barrier Reef, Australia, greatly increase available coral reef habitat. *ICES J. Mar. Sci.*, 2012, **70**(2), 284–293.
10. Sinniger, F. and Harii, S., Studies on mesophotic coral ecosystems in Japan. In *Coral Reef Studies of Japan*, Springer, Singapore, 2018, pp. 149–162.
11. Huang, D., Benzoni, F., Fukami, H., Knowlton, N., Smith, N. D. and Budd, A. F., Taxonomic classification of the reef coral families Merulinidae, Montastraeidae, and Diploastraeidae (Cnidaria: Anthozoa: Scleractinia). *Zool. J. Linn. Soc.*, 2014, **171**(2), 277–355.
12. Cairns, S. D., A revision of the shallow-water azooxanthellate Scleractinia of the Western Atlantic. *Stud. Nat. Hist. Caribb.*, 2000, **75**, 1–231.
13. Kitahara, M. V., Species richness and distribution of azooxanthellate Scleractinia in Brazil. *Bull. Mar. Sci.*, 2007, **81**, 497–518.
14. Laborel, J., Les peuplements de madréporaires des cotes tropicales du Brésil. Abidjan: Ann. del'Université'Abidjan, 1970.
15. Neves, E., Silveira, F. L., Pinchon, M. and Johnsson, R., Cnidaria, Scleractinia, Siderastrea, *Siderastrea siderea* (Ellis and Solander, 1786): Hartt expedition and the first record of a Caribbean siderastroid in tropical southwestern Atlantic. *Check List*, 2010, **6**, 505–510.
16. Castro, C. B., Medeiros, M. S. and Loiola, L. L., Octocorallia (Cnidaria: Anthozoa) from Brazilian reefs. *J. Nat. Hist.*, 2010, **44**, 763–827.

17. Pérez, C. D., Neves, B. M. and Oliveira, D. H., New records of octocorals (Cnidaria: Anthozoa) from the Brazilian coast. *Aquat. Biol.*, 2011, **13**(3), 203–214.
18. Fabricius, K. and Alderslade, P., *Soft Corals and Sea Fans: A Comprehensive Guide to the Tropical Shallow-water Genera of the Central-West Pacific, the Indian Ocean and the Red Sea*, Australian Institute of Marine Science, Townsville, Australia, 2001, p. 264.
19. Antony Fernando, S., *Monograph on Gorgonids (Sea Fans) of India*, Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, 2011, pp. 1–145.
20. Smith, J. L. B., *Smiths' Sea Fishes*, Struik, South Africa, 2003.
21. Froese, R. and Pauly, D. (eds), FishBase. World Wide Web electronic publication, 2017; www.fishbase.org, version (06/2017).
22. Newman, L. J. and Cannon, L. R. G., *Marine Flatworms: The World of Polyclad Flatworms*, CSIRO Publishing, Collinwood, Australia, 2003, p. 97.
23. Rasmussen, A. R., Sea snakes. In *FAO Species Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Central Pacific* (eds Carpenter, K. E. and Niem, V. H.), 2001, vol. 6, pp. 3987–4008.
24. Weinstein, D. K., Smith, T. B. and Klaus, J. S., Mesophotic bioerosion: variability and structural impact on US Virgin Island deep reefs. *Geomorphology*, 2014, **222**, 14–24.
25. Reid, R. and Macintyre, I., Foraminiferal-algal nodules from the eastern Caribbean: growth history and implications on the value of nodules as paleoenvironmental indicators. *Palaos*, 1988, **3**, 424–435.
26. Focke, J. and Gebelein, C., Marine lithification of reef rock and rhodolites at a fore-reef slope locality off Bermuda. *Geol. Mijnbouw*, 1978, **57**, 163–171.
27. Ballantine, D. L., Ruiz, H. and Aponte, N. E., Changes to macroalgal species communities in shelf edge coral reef habitats in southwest Puerto Rico. In 11th International Coral Reef Symposiums, Abstr., Fort Lauderdale, Florida, USA, 7–11 July 2008, p. 424.
28. Rivero-Calle, S., Armstrong, R. A. and Soto-Santiago, F. J., Biological and physical characteristics of a mesophotic coral reef: Black Jack reef, Vieques, Puerto Rico. In Proceedings of the 11th International Coral Reef Symposium, Ft. Lauderdale, Florida, USA, 7–11 July 2008, pp. 574–578.
29. Bridge, T. C. L., Done, T. J., Beaman, R. J., Friedman, A., Williams, S. B., Pizarro, O. and Webster, J. M., Topography, substratum, and benthic macrofaunal relationships on a tropical mesophotic shelf margin, central Great Barrier Reef, Australia. *Coral Reefs*, 2011, **30**, 143–153.
30. Littler, M. M., Littler, D. S. and Hanisak, M. D., Deep-water rhodolith distribution, productivity, and growth history at sites of formation and subsequent degradation. *J. Exp. Mar. Biol. Ecol.*, 1991, **150**, 163–182.
31. Amado-Filho, G. M. *et al.*, Rhodolith beds are major CaCO₃ bio-factories in the tropical South West Atlantic. *PLoS ONE*, 2012, **7**, e35171.
32. Iryu, Y., Nakamori, T., Matsuda, S. and Abe, O., Distribution of marine organisms and its geological significance in the modern reef complex of the Ryukyu Islands. *Sediment. Geol.*, 1995, **99**, 243–258.
33. Agegian, C. R. and Abbott, I. A., Deep water macroalgal communities: a comparison between Penguin Bank (Hawaii) and Johnston Atoll. In Proceedings of the Fifth International Coral Reef Symposium, 1985, vol. 5, pp. 47–50.
34. Rooney, J. *et al.*, Mesophotic coral ecosystems in the Hawaiian Archipelago. *Coral Reefs*, 2010, **29**, 361–367.
35. Luck, D., Forsman, Z. and Toonen, R., Polyphyly and hidden species among Hawaii's dominant mesophotic coral genera, *Leptoseris* and *Pavona* (Scleractinia: Agariciidae). *Peer J.*, 2013, e132.
36. Dinesen, Z. D., A revision of the coral genus *Leptoseris* (Scleractinia: Fungiina: Agariciidae). *Mem. Queensl. Mus.*, 1980, **20**(1), 181–235.
37. Hoeksema, B. W., Dautova, T. N., Savinkin, O. V., Tuan, V. S., Ben, H. X., Hoang, P. K. and Du, H. T., The westernmost record of the coral *Leptoseris kalayaanensis* in the South China Sea. *Zool. Stud.*, 2010, **49**, 325.
38. Hoeksema, B. W., Forever in the dark: the cave-dwelling azooxanthellate reef coral *Leptoseris troglodyta* sp. n. (Scleractinia, Agariciidae). *ZooKeys*, 2012, **228**, 21–37.
39. Roark, E. B., Guilderson, T. P., Dunbar, R. B., Fallon, S. J. and Mucciarone, D. A., Extreme longevity in proteinaceous deep-sea corals. *Proc. Natl. Acad. Sci. USA*, 2009, **106**(13), 5204–5208.
40. Jones, C. G., Lawton, J. H. and Shachak, M., Organisms as ecosystem engineers. In *Ecosystem Management*, Springer, New York, USA, 1994, pp. 130–147.
41. Hourigan, T. F., Lumsden, S. E., Dorr, G., Bruckner, A. W., Brooke, S. and Stone, R. P., State of deep coral ecosystems of the United States: introduction and national overview. In *The State of Deep coral Ecosystems of the United States* (eds Lumsden, S. E. *et al.*), National Oceanic and Atmospheric Administration, USA, 2007, pp. 1–64.
42. Brokovich, E. *et al.*, Grazing pressure on coral reefs decreases across a wide depth gradient in the Gulf of Aqaba, Red Sea. *Mar. Ecol. Prog. Ser.*, 2010, **399**, 69–80.
43. Garcia-Sais, J. R., Reef habitats and associated sessile-benthic and fish assemblages across a euphotic-mesophotic depth gradient in Isla Desecheo, Puerto Rico. *Coral Reefs*, 2010, **29**(2), 277–288.
44. Lesser, M. P. and Slaterry, M., Phase shift to algal dominated communities at mesophotic depths associated with lionfish (*Pterois volitans*) invasion on a Bahamian coral reef. *Biol. Invasions*, 2011, **13**, 1855–1868.
45. Darling, E. S., Green, S. J., O'Leary, J. K. and Côté, I. M., Indo-Pacific lionfish are larger and more abundant on invaded reefs: a comparison of Kenyan and Bahamian lionfish populations. *Biol. Invasions*, 2011, **13**(9), 2045–2051.
46. Kulbicki, M., Beets, J., Chabanet, P., Cure, K., Darling, E., Floeter, S. R. and Letourneur, Y., Distributions of Indo-Pacific lionfishes *Pterois* spp. in their native ranges: implications for the Atlantic invasion. *Mar. Ecol. Prog. Ser.*, 2012, **446**, 189–205.
47. Gosliner, T., Behrens, D. W. and Williams, G. C., *Coral Reef Animals of the Indo-Pacific: Animal Life from Africa to Hawaii Exclusive of the Vertebrates*, Sea Challengers, Monterey, California, USA, 1996.
48. Jie, W. B., Kuo, S. C. and Mok, H. K., Re-description of *Thysanozoon nigropapillosum* (Polycladida: Pseudocerotidae) from the South China Sea, with observations on a novel pre-copulatory structure, sexual behaviour and diet. *Raff. Bull. Zool.*, 2014, **62**, 764–770.
49. Feulner, G. R., Neubert, E. and Green, S., Land snails. In *The Emirates: A Natural History* (eds Hellyer, P. and Aspinall, S.), Trident Press, London, 2005, pp. 222–227.
50. Baker, E. K. and Harris, P. T., Habitat mapping and marine management. In *Seafloor Geomorphology as Benthic Habitat: GeoHAB Atlas of Seafloor Geomorphic Features and Benthic Habitats*, Elsevier, London, 2011, pp. 23–38.

ACKNOWLEDGEMENTS. We thank the Director, ICAR-Central Marine Fisheries Research Institute, Cochin for encouragement and support. We also thank S. B. Aravind (Temple Adventures, Puducherry) for SCUBA diving support.

Received 10 August 2017; accepted 19 December 2018

doi: 10.18520/cs/v116/i6/982-987