

Community organization of reef fishes in the live coral sub-habitat of Kavaratti atoll, Lakshadweep, India

P.E. VIJAY ANAND* AND N.G.K. PILLAI

Central Marine Fisheries Research Institute, Cochin - 682 014, India

**Aquacare, Division of Tetragon Chemie Pvt. Ltd., IS-40, KHB Ind. Area, Yelahanka New Town, Bangalore-560 064*

ABSTRACT

Data on the community organization of coral reef fishes of Kavaratti atoll, Lakshadweep, India were collected during the period January 1991 to June 1992. Species were enumerated by visual census on the live coral sub-habitat for frequency of occurrence, abundance, composition, diversity, evenness and seasonal variation in community parameters. As the live coral zone was composed of a single species of ramose coral (*Acropora formosa*), only 14 families and 39 species of reef fishes made use of this zone, the community diversity being 1.84 and 3.16. Chaetodontids, labrids and pomacentrids were comparatively more abundant. Varying habitats of chaetodontids explain their restricted distribution. Live coral does not seem to be a preferred habitat of labrids. The occurrence of epinephelids was influenced by readily available prey. Most species among live coral were resident, variations resulted from factors affecting new recruits rather than habitat shifts.

Introduction

Discrete populations, small breeding pools, sedentary and non-migratory nature and association with specific habitats are striking characters of coral reef fishes. A variety of habitats found in coral reef areas may be rich or poor in species within the habitat or between habitats, supporting markedly different fish communities of which some may be cosmopolitan. Dependence on a particular food item, behavioral interactions and depths across reefs are other factors which limit reef fish distribution.

Although the study of the ecology of reef fishes is relatively recent, it is

developing rapidly, with many workers at present active in the tropical western Atlantic and Caribbean, the Gulf of California, several centres in the east Indo-West Pacific, and in the Red Sea (Sale, 1980). However, there is an overall paucity of knowledge on this subject from the Indian region, leaving gaps in reviews, which attempted to summarize a global picture on reef fish community organization. The present investigation provides a basis on coral reef fish community organization with reference to patterns of numerical densities, species richness, species composition and their occurrence on each of the five sub-habitats of Kavaratti atoll and the reef slope in general. This communication

deals with the live coral sub-habitat of Kavaratti atoll.

Materials and methods

Fish species enumeration and their distribution in the sub-habitats of Kavaratti atoll were studied using the visual census technique.

The visual census techniques include a considerable variety of procedures (Sale, 1980). This method is non-destructive and permits repeated sampling of an area without physical disturbances to the populations. In the present investigation due to limited facilities, a combination of/or modification of existing methods has been used to suit the sub-habitats. Belt transects of 30 m length (Goldman and Talbot, 1976) were used to record all visible fish species and numbers from 2 m on either side of a pre-laid line. Transect width was estimated visually (Roberts and Ormond, 1987). Due to shallow depths and absence of SCUBA facility, the fish were observed using a mask and snorkel apparatus (Sale, 1974; Goldman and Talbot, 1976; Bouchon-Navaro and Harmelin-Vivien, 1981; Hutomo and Adrim, 1986; Adrim and Hutomo, 1989; Bouchon-Navaro and Bouchon, 1989) with three intervals to

record the observations. The approximate area covered by a single census unit is 120 sq. m. for each sub-habitat. Censuses were conducted fortnightly for 18 months which resulted in 27 censuses for each sub-habitat. Censuses were not conducted when rough weather conditions prevailed. The period of study was from January 1991 to June 1992.

Data collected from each census was grouped by species. These were transformed into a data matrix for the sub-habitat (Fig. 1). Results obtained from the data matrix were used to study the community parameters and seasonal variation in community parameters.

1. *Community parameters*: Results of all censuses were combined to obtain the following aspects:

- a. Species composition of the particular region listed in the order of frequency of occurrence
- b. Frequency of occurrence
- c. Total abundance
- d. Percentage abundance
- e. Shannon-Weiner diversity index (H')

2. *Seasonal variation in community parameters*: Results of all censuses were treated separately and grouped into three distinct seasons prevailing in Lakshadweep region, namely, pre-monsoon (January to April), monsoon (May to August) and post-monsoon (September to December). These seasons have been classified based on a definite change in wind direction in the middle of May (south-westerly) and its subsequent termination in August and replacement by north-easterly winds. The following aspects were studied.

- a. Total number of individuals




Fig.1 Data matrix for community organisation studies on sub-habitats.

- b. Total number of species
- c. Shannon-Weiner diversity index (H') (Shannon-Weiner, 1949)
- d. Evenness index (J') (Pielou, 1966)

Results

Community parameters : Table 1 records the community parameters of 39 species recorded from subhabitat.

Variation in community parameters : The variation in community parameters is given in Table 2. Variation in total number of individuals during the pre-monsoon, was high (61 to 150) in 1991, while the same in 1992 was relatively low (57 to 86). January 1, March 4 and February 2 (1991) recorded 150, 115 and 108 individuals respectively. In the monsoon, June 3, 1991 recorded the least count (63), while in June 2, 1992 it was 240 counts. In general, May in both the years accounted for more fishes (106 to 126 nos). Counts during post-monsoon were also not consistent (72 to 141). Maximum count (141 nos) was recorded on October 3.

Total number of species recorded, ranged between 7 and 16 in pre-monsoon, with the maximum (16) during February 2 and April 6, 1991. Three samples during April 7 ('91), 5 and 6 ('92) accounted for least species counts of 8, 7 and 7 respectively. Species counts were stable (10-15) during monsoon, while in post-monsoon a minimum of 8 species in December 7 and a maximum of 14 species in September 1 were recorded.

During pre-monsoon, species diversity was high only in February 2 (3.39) and April 6 (3.35), while in all other samples it was below 3.00. May 2, June 3, July 4 in 1991 recorded diversities of 3.16, 3.13 and 3.28 respectively, while the least (1.77) was in June 2, 1992. Diversity during other monsoon months ranged between 2.81 and 2.90. In the post-monsoon, all samples taken in September 1, 2 and a single sample in November 6 had diversities of 3.02, 3.04 and 3.26 respectively, while October 3, 4 and December 7 registered values between 2.44 and 2.86.

TABLE 1. Frequency of occurrence (FO), total abundance (TA), percentage abundance (%) and diversity index (H') for each species recorded in the live coral sub-habitat of Kavaratti atoll

FAMILY / SPECIES	FO	TA	%	H'
ACANTHURIDAE				
<i>Acanthurus leucosternon</i>	1	1	0.04	0.00
BALISTIDAE				
<i>Balistoides viridescens</i>	14	35	1.31	3.44
<i>Rhinecanthus aculeatus</i>	4	5	0.19	1.91
CHAETODONTIDAE				
<i>Chaetodon auriga</i>	24	123	4.61	4.42
<i>Chaetodon xanthocephalus</i>	12	44	1.65	3.26
<i>Chaetodon trifascialis</i>	5	9	0.34	2.27
<i>Chaetodon trifasciatus</i>	5	19	0.71	2.10
<i>Chaetodon vagabundus</i>	5	13	0.49	2.27
<i>Chaetodon falcula</i>	4	7	0.26	1.96
<i>Chaetodon melannotus</i>	3	9	0.34	1.39

<i>Heniochus acuminatus</i>	3	14	0.53	1.50
<i>Chaetodon citrinellus</i>	2	7	0.26	0.98
CIRRHITIDAE				
<i>Paracirrhites forsteri</i>	7	21	0.79	2.68
FISTULARIDAE				
<i>Fistularia petimba</i>	1	1	0.04	0.00
GRAMMISTIDAE				
<i>Grammistes sexlineatus</i>	1	1	0.04	0.00
LABRIDAE				
<i>Halichoeres scapularis</i>	21	141	5.29	4.21
<i>Halichoeres centiquadrus</i>	6	25	0.94	1.74
<i>Thalassoma hardwicki</i>	4	40	0.15	1.85
<i>Coris gaimard</i>	2	3	0.11	0.92
<i>Labroides dimidiatus</i>	2	2	0.08	1.00
<i>Coris formosa</i>	1	1	0.04	0.00
<i>Cheilinus undulatus</i>	1	1	0.04	0.00
MURAENIDAE				
<i>Gymnothorax undulatus</i>	6	8	0.30	2.43
OSTRACIIDAE				
<i>Lactoria cornuta</i>	2	2	0.08	1.00
<i>Ostracion meleagris</i>	2	2	0.08	1.00
POMACANTHIDAE				
<i>Centropyge multispinis</i>	25	348	14.4	4.55
POMACENTRIDAE				
<i>Dascyllus aruanus</i>	27	657	24.64	4.46
<i>Chromis caerulea</i>	23	579	21.69	4.40
<i>Dascyllus trimaculatus</i>	17	172	6.45	3.80
<i>Dascyllus reticulatus</i>	14	133	4.99	3.62
<i>Abudefduf sexfasciatus</i>	10	89	3.34	3.16
<i>Amphiprion chrysogaster</i>	2	6	0.23	0.92
<i>Pomacentrus pavo</i>	2	4	0.15	1.00
<i>Chrysiptera biocellata</i>	1	3	0.11	0.00
SERRANIDAE				
<i>Epinephelus</i> spp.	17	59	2.21	3.82
<i>Cephalopholis argus</i>	4	4	0.15	2.00
SCORPAENIDAE				
<i>Dendrochirus zebra</i>	13	33	1.24	3.47
<i>Pterois radiata</i>	1	1	0.04	0.00
TETRAODONTIDAE				
<i>Tetraodon nigropunctatus</i>	5	23	0.86	1.19

Number of species recorded = 39

TABLE 2. Total number of individuals (*I*), number of species (*S*), diversity index (*H*ⁱ) and evenness (*J*) for each census in different seasons on the live coral sub-habitat of Kavarratti atoll

Sample No.		1	2	3	4	5	6	7
A PRE-MONSOON 1991		J	F	F	M	M	A	A
	I	150	108	81	115	61	99	86
	S	12	16	11	11	10	16	8
	H ⁱ	2.57	3.39	2.80	2.73	2.90	3.35	2.81
	J ⁱ	0.72	0.85	0.81	0.79	0.87	0.84	0.94
B MONSOON		MY	MY	JN	JY	AG		
	I	109	106	63	65	85		
	S	10	15	12	14	13		
	H ⁱ	2.81	3.16	3.13	3.28	2.90		
	J ⁱ	0.85	0.81	0.87	0.86	0.78		
C POST- MONSOON		S	S	O	O	N	N	D
	I	72	123	141	78	85	129	91
	S	14	12	10	9	11	12	8
	H ⁱ	3.02	3.04	2.86	2.44	2.84	3.26	2.79
	J ⁱ	0.79	0.85	0.86	0.71	0.82	0.91	0.93
D PRE-MONSOON 1992		J	F	M	M	A	A	
	I	84	57	82	86	83	61	
	S	10	9	11	11	7	7	
	H ⁱ	2.70	2.52	2.87	2.85	2.42	2.48	
	J ⁱ	0.81	0.79	0.83	0.77	0.65	0.67	
E MONSOON (2 samples)	I = 126, 240; H ⁱ = 2.83, 1.77; J ⁱ = 0.79, 0.51;							
May, June	S = 12, 11.							

J : January; F : February; M : March; A : April; MY : May; JN : June; JY : July; AG : August; S : September; O : October; N : November; D : December

Evenness in distribution of species was maximum (0.94) in April 7, 1991 and minimum (0.65 and 0.67) in two samples taken in April 5, 6 1992 during pre-monsoon. In general, a majority of samples registered values between 0.80 and 0.88. During monsoon of '91, May 1, 2, June 3 and July 4 showed relatively stable patterns of distribution (0.81 to 0.87), while in August 5 it was 0.78. Two monsoon samples taken in '92 however, showed uneven distribution (0.79 and 0.51). Distribution was not consistent during post-monsoon with a minimum (0.71) in October 4 and a maximum (0.93

and 0.91) in December 7 and November 6 respectively. Except in September 1 (0.79) other samples indicated a relatively stable distribution (0.82 to 0.86).

Discussion

Species richness of three families, namely Chaetodontidae, Labridae and Pomacentridae were high and comparable, each of them consisting of 7 to 8 species. Chaetodontids are known to spend their entire lives associated with a relatively small portion of the reef environment (Goldman and Talbot,

1976), while some species are relatively restricted in their distribution among zones and others widely distributed (Bouchon-Navaro, 1980, 1981; Foulter, 1990). Certain species (e.g., *Chaetodon trifasciatus* and *C. trifascialis*) are considered as obligatory coral feeders (Reese, 1981; Harmelin-Vivien and Bouchon-Navaro, 1983; Harmelin-Vivien, 1989; Sano, 1989), while certain species are considered as facultative coral feeders (Harmelin-Vivien, 1989; Sano, 1989). In the present study, both *C. trifascialis* and *C. trifasciatus* were found among *Acropora formosa* patch reefs but low H^i values (3.20 and 2.77) suggested that they were not frequent as *C. auriga*. *C. vagabundus* is also known to feed on corals (Sano, 1989). Sutton (1985) observed that *C. trifasciatus* had larger territories within lagoon habitats and interpreted this as a response to low population density and poor habitat quality. As patch reefs are not extensive on Kavaratti atoll, it is likely that obligate coral feeders are found in habitats with richer coral cover than on sparse coral patches. Bouchon-Navaro *et al.* (1985) found chaetodontid populations to decrease with degraded coral environment. Due to the facultative feeding habit, *C. auriga* (Hourigan, 1989) is more flexible in habitat choice, and thus the abundance. *C. citrinellus* is a non-coral feeder (Galzin, 1987), while *C. melannotus* is a soft coral feeder (Adrim and Hutomo, 1989) and *C. falcula* was found to feed predominately on polychaete tentacles. Low distribution of these species in the present study is probably due to varying habits mentioned above.

Among the seven labrids, *Halichoeres scapularis* was distinct from other species ($H^i = 4.21$) in abundance and occurrence. Though this species did not seem to depend on this zone directly,

it probably made use of interstitial sand patches for food procurement. This species was extensively associated with sand flats (Randall, 1983). *Thalassoma hardwicki* was observed on quiet reefs (Hiatt and Strasburg, 1960), while *Coris gaimard* was numerous where the reef was interspread with patches of sand (Hobson, 1974). In the present study, ramose corals perhaps did not suit *T. hardwicki* due to lack of shelter, while the presence of *C. gaimard* supports the observation of Hobson (1974). It is also likely that *C. formosa* also follows the same pattern as these species feed on benthic invertebrates on sand. However, rare occurrence of these species on live coral suggest that their preferred habitats lie elsewhere on the reef.

Of the eight pomacentrids, only *Dascyllus aruanus* and *Chromis caerulea* occurred frequently in high numbers. The fact is substantiated by H^i values (4.46 and 4.40 respectively) that weighted their occurrence in most census conducted. Relatively common species were *D. trimaculatus* and *D. reticulatus*. Shpigel (1982) observed that *D. aruanus* and *D. marginatus* are coral dwelling species and occur together frequently. Reese (1978) stated that mobility of planktivores is constrained for want of shelter or nesting sites. Due to intense site attachment, these species would have been included in most censuses. Pillai *et al.* (1986) reported co-existence of *D. aruanus* and *Chromis caerulea* on coral patches in Minicoy. Apart from this, few more interspecific associations, involving *D. trimaculatus*, *D. reticulatus* and *Amphiprion chrysogaster* were observed in the present study. The occurrence of *A. chrysogaster* was determined by the presence of small anemones on dead bases of *Acropora formosa*. Other rare species were accidental in occurrence.

The occurrence of *Epinephelus* spp. (probably *E. hexagonatus*) and *Dendrochirus zebra* and occasionally *Gymnothorax undulatus* is influenced by the readily available prey in the form of small pomacentrids and other invertebrates. *E. hexagonatus* were found to inhabit staghorn coral zones in Tutia reef (Talbot, 1965). The abundance and consistent occurrence of *Centropyge multispinis* is due to strong site attachment (as in pomacentrids) and were observed in most censuses. *Ostracion cubicus* was observed to be a weak swimmer, found around coral heads while, *Tetraodon nigropunctatus* fed on corals (Hiatt and Strasburg, 1960). In the present study, occurrence of *O. melegritis* and *T. nigropunctatus* could be possibly structured due to these habitats. A similar reasoning holds good for *Balistoides viridescens*. Though *Paracirrhites forsteri* is known to be associated with coral heads (Talbot, 1965; Galzin, 1987), they seem to prefer the tabloid *Acropora humilis* head over *A. formosa*. Other species that occurred only once could be considered as accidental.

Though the number of species did not vary much during monsoon, variation in total individuals indicated that each species supported varying number of individuals and the least during peak monsoon. This is supported by H' and J' values. As recruitment occurs during summer and the monsoon follows almost immediately in Lakshadweep, fish species count could possibly be influenced due to mortality or movements from affected areas. A similar reasoning was proposed by Jones (1991). A higher variation in community parameters during pre-monsoon could have resulted due to new recruits from the late post-monsoon months (November, December) coupled with

effects of summer months. Summer peaks in recruitment were recorded by Sale and Dybdahl (1975) and Coles and Tarr (1990). As seen earlier, most fish populations among live coral are resident and variations could result mainly through factors affecting new recruits rather than habitat shifts. Sale (1983) observed a highly variable community structure of fishes from one census to another on patch reefs and suggested that a variable pattern of settlement of new juvenile fish plays a major role in determining fish assemblages. Other than random settlement, predation pressure also influences assemblages (Shpigel, 1982).

Acknowledgments

The authors are grateful to Dr. P.S.B.R. James, Former Director, Central Marine Fisheries Research Institute, Cochin, for encouragement and for providing facilities. For the field facilities in Lakshadweep, they are indebted to the Administration. The first author is also grateful to the Indian Council of Agricultural Research, New Delhi for awarding a Senior Research Fellowship for conducting the study.

References

- Adrim, M. and M. Hutomo 1989. Species composition, distribution and abundance of Chaetodontidae along reef transects in the Flores Sea. *Neth. J. Sea Res.*, **23** : 85-93.
- Bouchon-Navaro, Y. 1980. Quantitative distribution of the Chaetodontidae on a fringing reef of the Jordanian coast (Gulf of Aqaba, Red Sea). *Tethys*, **9** : 247-251.
- Bouchon-Navaro, Y. 1981. Quantitative distribution of Chaetodontidae on a reef of Moorea Island (French Polynesia). *J. Exp. Mar. Biol. Ecol.*,

- 55 : 145-157.
- Bouchon-Navaro, Y. and C. Bouchon 1989. Correlations between chaetodontid fishes and coral communities of the Gulf of Aqaba (Red Sea). *Environ. Biol. Fishes*, **25**: 47-60.
- Bouchon-Navaro, Y. and M.L. Harmelin-Vivien 1981. Quantitative distribution of herbivorous reef fishes in the Gulf of Aqaba (Red Sea). *Mar. Biol.*, **63**: 79-86.
- Bouchon-Navaro, Y., C. Bouchon and Harmelin-Vivien 1985. Impact of coral degradation on a chaetodontid fish assemblage. *Proc. 5th Int. Coral Reef Congr.*, P. 427-432. Moorea, French Polynesia
- Coles, S.L. and A.B. Tarr 1990. Reef fish assemblages in the western Arabian Gulf : a geographically isolated population in an extreme environment. *Bull. Mar. Sci.*, **47**: 696-720.
- Fouler, A.J. 1990. Spatial and temporal patterns of distribution and abundance of chaetodontid fishes at One Tree Reef, southern GBR. *Mar. Ecol. Prog. Ser.*, **64**: 39-53.
- Galzin, R. 1987. Structure of fish communities of French Polynesia coral reefs. II. Temporal scales. *Mar. Ecol. Progr. Ser.*, **41** : 137-145.
- Goldman, B. and F.H. Talbot 1976. Distribution of inshore fishes at Canton Atoll. *Atoll Res. Bull.*, **221** : 99-157.
- Harmelin-Vivien, M.L. 1989. Implications of feeding specialization on the recruitment process and community structure of butterflyfishes. *Environ. Biol. Fishes*, **25** : 101-110.
- Harmelin-Vivien, M.L. and Y. Bouchon-Navaro 1983. Feeding diets and significance of coral feeding among chaetodontid fish in Moorea (French Polynesia). *Coral Reefs*, **2** : 119-127.
- Hiatt, R.W. and D.W. Strasburg 1960. Ecological relationships of the fish fauna on coral reefs of the Marshall Islands. *Ecol. Monogr.*, **30** : 65-127.
- Hobson, E.S. 1974. Feeding relationships of teleostean fishes on coral reefs in Kona, Hawaii. *Fish. Bull.*, **72** (4) : 915-1031.
- Hourigan, T.F. 1989. Environmental determinants of butterflyfish social systems. *Environ. Biol. Fishes*, **25** : 61-78.
- Hutomo, M. and M. Adrim 1986. Distribution of reef fish among transects in Bay of Jakarta and Kepulauan Seribu. *UNESCO Rep. Mar. Sci.*, **40** : 135-156.
- Jones, G.P. 1991. Post recruitment processes in the ecology of coral reef fish populations : a multifunctional perspective. In : P.F. Sale (Ed.), *The Ecology of Fishes on Coral Reefs*. P. 294-328. Academic Press Inc., San Diego, California.
- Pielou, E.C. 1966. Measurement of diversity in different types of biological collections. *Theoretical Biol.*, **13**: 131-144.
- Pillai, C.S.G., Madan Mohan and K.K. Kunhikoya 1986. Functional mechanism of co-existence in *Chromis caeruleus* and *Dascyllus aruanus* (Pomacentridae : Pisces). *Indian J. Fish.*, **32** : 18-26.
- Randall, J.E. 1983. *Red Sea Reef Fishes*. IMMEL Publ. London
- Reese, E.S. 1978. The study of space related behaviour in aquatic animals

- : special problems and selected examples. In : E.S. Reese, F.S. Lighter (Eds.) *Contrasts in behaviour*. P. 347-374. John Wiley, New York.
- Reese, E.S. 1981. Predation on corals by fishes of the family Chaetodontidae : implications for conservation and management of coral reef ecosystems. *Bull. Mar. Sci.*, **31** : 594-604.
- Roberts, C.M. and R.F.G. Ormond 1987. Habitat complexity and coral reef fish diversity and abundance on Red Sea fringing reefs. *Mar. Ecol. Prog. Ser.*, **41** : 1-8.
- Sale, P.F. 1974. Mechanisms of co-existence in a guild of territorial fishes at Heron Island. *Proc. 2nd Int. Coral Reef Symp.*, **1** : 193-206.
- Sale, P.F. 1980. The ecology of fishes on coral reefs. *Oceanogr. Mar. Biol.*, **18** : 367-421.
- Sale, P.F. 1983. Temporal variability in the structure of reef fish communities. *Proc. Great Barrier Reef Conf.*, P. 239-244.
- Sale, P.F. and R. Dybdahl 1975. Determinants of community structure for coral reef fishes in an experimental habitat. *Ecology*, **56** : 1343-1355.
- Sano, M. 1989. Feeding habits of Japanese butterflyfishes (Chaetodontidae). *Environ. Biol. Fishes.*, **25** : 195-203.
- Shannon, C.E. and W. Weiner 1949. *The Mathematical Theory of Communication*. University of Illinois Press, Urban.
- Shpigel, M. 1982 Niche overlap among two species of coral dwelling fishes of the genus, *Dascyllus* (Pomacentridae). *Environ. Biol. Fishes*, **7** : 65-68.
- Sutton, M. 1985. Patterns of spacing in a coral reef fish in two habitats on the Great Barrier Reef. *Anim. Behav.*, **33** (4) : 1332-1337.
- Talbot, F.H. 1965. A description of the coral structure of Tutia Reef (Tanganyika Territory, East Africa) and its fish fauna. *Proc. Zool. Soc. London*, **145** : 431-470.