Avian fauna sighting

hirty five species ofcoastal birds were identified and photographed from 10 surveys conducted in the coastal regions of Ullal, Someshwara, Mangalore Bunder, Bengre, Kudroli, Hegemadikodi, Mulki, Kaup, Uppunda, and Murudeshwar. This included the groups of Egrets, Herons, Cormorants, Kites, Shanks, Sandpipers, Lapwingsand Plovers. Among seabirds, the major birds observed were the gulls, brown head Larusbrunnicephalus and black headed Larusridibundus followed by the terns. Oceanic birds Pomarine Skua Stercorarius pomarinus 4 no's were observed off Mangalore coast on 8.11.12. At Uppunda, the fishermen had caught a vagrant Blue-footed Booby Sula nebouxii. The bird belongs

to the Order Pelecaniformes and FamilySulidae. This has not been reported from the coast of Karnataka. The natural breeding habitat of the Bluefooted Booby is tropical and subtropical islands off the Pacific Ocean especially found in the Galapagos Islands, Ecuador

where they are a protected species. The IUCN status of the bird is leastconcern. Sardine is the main food for the bird and recent studies in Galapagos Islands suggest that, there is a shortage of food for the birds hence the birds are failing to breed. (Reported by Mangalore RC.)

Little cormorantMicrocarboniger, Large EgretCasmerodiusalbus, Western reef EgretEgrettagularis, Cattle EgretBubulcus ibis and Little EgretEgrettagarzetta observed at Bunder 21.11.12 min: CMFRI Newsletter No. 135 | 11