

## NOTES

### ON *STERECHINUS ANTARCTICUS* KOEHLER (CAMERODONTA: ECHINOIDEA: ECHINODERMATA) COLLECTED FROM THE COASTAL WATERS OF ANTARCTICA OFF QUEEN MAUD LAND

#### ABSTRACT

*Sterechinus antarcticus* Koehler 1901, an echinoderm of the family Echinoidea and order Camerodonta collected from the coastal waters of Antarctica off Queen Maud Land is reported giving some of its morphological features and remarks on its affinity to two closely related species namely *S. diadema* and *S. agassizii*.

*STERECHINUS ANTARCTICUS* Koehler 1901 is a species of sea urchin found distributed around the Antarctic continent with its northernmost geographical limit going as far north as south Georgia (Mortensen 1943). A specimen of this species (Plate I) was collected during the Third Indian Scientific Expedition to Antarctica in 1983-'84 period by the second author. The specimen was collected using a snapper from a depth of 150 m at 70°02'S and 12°36'E about 1 km away from the Antarctic ice edge at Queen Maud Land. As the snapper brought no sedimentary material it is assumed that the substratum at the Station is hard. Since the material would form one more record to the geographical distribution of the species, a brief description of the specimen is presented with remarks on its close relationship with two other species namely *S. diadema* and *S. agassizii*.

The authors wish to thank Dr. P.S.B.R. James, former Director, Central Marine Fisheries Research Institute, Cochin for his interest in this work. The second author is thankful to the Department of Ocean Development for providing all facilities during the Expedition.

**Material** : A single specimen of h.d. 24 mm and v.d. 14 mm (Plate I).

**Description** : The test is globular. The apical system is 25% of the diameter of the test. The ocular plates are all insert. The anus is excentric in position. Sections of the spines show that the spokes are round and broad.

Peristome is large and forms 40% of the diameter of the disc. The peristomial membrane is thin.

The Aristotle's lantern (Plate I) has well developed epiphysis. The rotule are also well developed. The pyramids are well formed. The teeth are somewhat weak.

Primary spines are short, pointed and grooved. Secondary spines are long and needle-like. The primary tubercles are arranged in distinct rows, those at the ambitus being the largest. The tubercles are noncrenulate.

Globiferous and triphyllous pedicellariae are distinctly seen. In globiferous pedicellariae the basal part of the valves are not produced. The valves of the triphyllous pedicellariae are broad and rounded.

Remarks: Whereas Koehler (1901) while establishing the species thought it to be quite

separate from *S. diadema*. Finally in his Discovery Report, Mortensen (1936) considered




PLATE I *Sterechinus antarcticus* Koehler 1. Dorsal view, 2. Ventral view, 3. Lateral view, 4. Aristotle's Lantern (Lateral view)

unique differing sharply from all other species of the order Camerodonta by the characters of its apical system and the smaller number of its conical plates. Mortensen (1903) came to the conclusion that it could not be distinguished from *S. diadema* and *S. agassizii*. In 1909, Mortensen (1909) while placing *S. antarcticus* as a separate species opined that if more intermediate forms are collected, this species could not be kept

*S. antarcticus* as separate species though very close to *S. diadema* and *S. agassizii*. In the comparative account made for these three related species Mortensen (1943) examined the various characters critically to conclude that they represented three distinct species eventhough there could be intermediate hybrid forms between *S. antarcticus* and *S. agassizii*, their area of distribution overlapping in the South Georgia region.

*Distribution:* Distributed all around the Antarctic continent. In the South American region it goes

as far north as S. Georgia. Bathymetrically the species is distributed between 100 and 1080 m.

*Central Marine Fisheries Research Institute,  
Kochi-682 014*

D. B. JAMES  
K. J. MATHEW

## REFERENCES

KOEHLER, R. 1901. Echinoides et Ophiurus. *Res. Voyage 'Belgica'*, 8.

MORTENSEN, T. 1903. Echinodermata-1. *'Ingolf' Exp.* 102.

————— 1909. Echnoiden. *Deuteschen Sudpolar Exp.*, 75

MORTENSEN, T. 1936. Echinoidea and Ophiuroidea. *Discovery Rep.*, 12 : 218.

————— 1943. Echinoidea and Ophiuroidea. *Monograph of the Echinoidea-III*, 1-446.