

CMFRI
bulletin 40



MAY 1987

**MARINE CATFISH
RESOURCES OF INDIA**
EXPLOITATION AND PROSPECTS

CENTRAL MARINE FISHERIES RESEARCH INSTITUTE
(Indian Council of Agricultural Research)
P. B. No. 2704, E. R. G. Road, Cochin-682 031, India

CHAPTER SIX

ASSESSMENT OF THE RESOURCES OF IMPORTANT SPECIES OF CATFISHES

— K. ALAGARAJA AND M. SRINATH

DATA BASE

The centres Waltair and Mandapam, in the east coast and Cochin, Mangalore and Veraval, in the west coast, where biological data on catfishes have been collected, are considered for the assessment of resources of *Tachysurus thalassinus*, *T. tenuispinis*, *T. serratus*, *T. dussumieri* and *Osteogeneiosus militaris*. Length-frequency data of *T. thalassinus* collected during 1974-76 at Waltair from trawl catches, during 1972-76 at Mandapam from trawl catches, during 1981 at Cochin Fisheries Harbour from the catches of trawls, purse-seines and gill nets and during 1981 and 1982 at Veraval from trawl catches are taken up for this study. For *T. tenuispinis*, the data collected during 1974-76 at Waltair from trawl catches, during 1981 at Cochin Fisheries Harbour and during 1982-83 at Mangalore both from trawls, purse-seines and gill nets are considered. For *T. serratus*, the length-frequency data collected from the catches of trawls, purse-seines and gill nets at Cochin Fisheries Harbour during 1981 and for *T. dussumieri* and *O. militaris* those collected from the catches of trawls at Veraval during 1981 and 1982 are also considered.

ESTIMATION OF GROWTH PARAMETERS AND THE TECHNIQUES USED

i. *Estimation of 'L_∞' and 'K'*: Assuming that the growth of catfish is isometric and following Von Bertalanffy's growth pattern, namely

$$L_t = L_\infty [1 - e^{-k(t - t_0)}] \dots\dots\dots (1)$$

in the usual notation, estimates for L_∞ and K are obtained using ELEFAN-I (Pauly et al, 1981). Since the estimates for each species did not show much variation between centres, sets of estimates (L_∞ and K) one for each

species is obtained and is presented along with the corresponding estimate for W_∞ in Table 1.

TABLE 1
Estimation of growth parameters

Species	Parameters	L_∞ (mm)	K(annual)	W_∞ (gm)
<i>T. thalassinus</i>		755	0.36	4030
<i>T. tenuispinis</i>		560	0.78	3230
<i>T. serratus</i>		1100	0.25	8000
<i>T. dussumieri</i>		850	0.25	6000
<i>O. militaris</i>		540	0.78	3200

It can be seen from the above table that estimates of 'K' decrease while the corresponding values of L_∞ increase. This is quite consistent with the growth model under consideration.

ii. *Estimates of 'Z' the Instantaneous Rate of Total Mortality*

Age-frequency distribution was found to be very difficult to obtain from the available data. Using length-frequency data and following the method of Alagaraja (1984) estimates of 'Z' for each year at each centre for every species mentioned above have been obtained along with their error estimates. In some cases three point moving averages have been taken for this purpose. That portion of the length frequency distribution which resembled the right limb of catch curve alone was considered for estimation of 'Z'. The steps taken for this purpose as well as marking the portion considered for the estimation of 'Z' are indicated in the work sheets enclosed.

The formula used for this purpose is:

$$\log (N_t + \Delta_t / N_t) = \frac{Z}{K} \log \frac{L_\infty - L_t + \Delta_t}{L_\infty - L_t} \dots\dots\dots (2)$$

Estimates of L_∞ and k are available from Table 1 and L_t and $L_t + \Delta_t$ are the successive mid values of the length classes whose frequencies are N_t and $N_t + \Delta_t$. Since constant 'Z' for the entire size range of fishing is considered, catches in numbers at successive ages C_t and, $C_t + \Delta_t$ are proportional to N_t and $N_t + \Delta_t$. Hence

$$N_t + \Delta_t / N_t = C_t + \Delta_t / C_t \dots \dots \dots (3)$$

This is made use of in the above formula (2) and length frequency data are used for estimation of 'Z'. The procedure for obtaining the estimates of 'Z' is as follows. The deviations of L_t from $L_\infty (= L_\infty - L_t)$ are to be taken. Converting them to log values (common log will do), the values $\log(L_\infty - L_t)$ are obtained. After tabulating these values, their successive differences $\log(L_\infty - L_t) - \log(L_\infty - L_t + \Delta_t)$ are calculated. In the same way, the successive differences of $\log(N_t)$ are to be taken and tabulated as follows :

L_t				$L_\infty - L_t$	$\log(L_\infty - L_t)$	$\frac{\Delta \log(L_\infty - L_t)}{A}$
1	2	3	4			
					C_t	$\frac{\Delta \log C_t}{B}$
5	6	7	8			Z/K B/A

Where $\Delta \log(L_\infty - L_t)$ and $\Delta \log C_t$ are the respective successive differences.

Thus for each row of successive differences an estimate of \bar{z}/k is available. If there are ' $n+1$ ' length groups then there will be ' n ' estimates of \bar{z}/k . If the first estimate of \bar{z}/k is termed as x_1 , the second as x_2 and so on with the last one as x_n then

$$\bar{z}/k = 1/n \sum_{i=1}^n x_i \dots \dots \dots (4)$$

and

$$s^2 \bar{z}/k = \frac{1}{n-1} \left[\sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2 / n \right] \dots \dots \dots (5)$$

Hence $s\bar{z}/k$ can be obtained as the square root of the above expression. The standard error of $\bar{z}/k (= s\bar{z}/k)$ is $\frac{1}{\sqrt{n}} s\bar{z}/k$. Multiplying \bar{z}/k and $s\bar{z}/k$ by the already available estimate of 'k' the estimate of \bar{z} and $s\bar{z}$ are derived. The

detail procedure is given in the worksheet II. The estimates thus obtained are given in tables 2-4 below along with the sample size (n), I_c and I_r , where I_c indicates the size at first capture of the fully recruited phase and I_r is the size at entry to the fishery.

TABLE 2

Values of the estimates of \bar{z} and its standard error and 'n' I_c and I_r for *T. thalassinus*

Centre	Year	Values				
		\bar{z}	$s\bar{z}$	n	I_c	I_r
		(mm)	(mm)			
1. Waltair (Trawls)	1974	2.34	0.41	4		
	1975	2.24	0.68	7	180	100
	1976	1.16	0.54	4		
	Combined	1.98	0.37	15	180	100
2. Mandapam (Trawls)	1972	2.32	0.62	6	170	
	1973	1.54	0.27	5	230	
	1974	1.16	0.36	7	190	
	1975	2.06	0.40	11	230	60
	1976	3.72	0.55	8	270	
	Combined	2.22	0.25	37	220	60
3. Cochin Fisheries Harbour (Trawl & gill net)	1981	2.06	0.52	9	360	100
4. Veraval (Trawls)	1981	1.71	0.88	4		
	1982	1.50	0.63	4	380	100
	Combined	1.60	0.50	8	380	100

TABLE 3

Values of estimates of \bar{z} , and its standard error and 'n', I_c and I_r for *T. tenuispinis*

Centre	Year	Values				
		\bar{z}	$s\bar{z}$	n	I_c	I_r
		(mm)	(mm)			
1. Waltair (Trawls)	1974	2.14	0.64	3	220	80
	1975	2.57	0.84	3	220	120
	1976	2.95	0.73	4	180	160
	Combined	2.59	0.40	10	200	120
2. Cochin Fisheries Harbour (Purse seine, trawls & gill nets)	1981	3.04	1.29	5	280	100
	1982	2.76	0.92	9	260	40
	Combined	2.85	0.95	14	280	100
3. Manga- lore (Purse seine trawl & gill nets)	1982-83	2.76	0.92	9	260	40

TABLE 4

Values of estimates of \bar{z} and its standard error and 'n', I_c and I_r for *T. serratus*, *T. dussumieri* and *O. militaris*

Species	Centre	year	Values				
			\bar{z}	$s\bar{z}$	n	I_c (mm)	I_r (mm)
i. <i>T. serratus</i>	Cochin Fisheries Harbour (Purse-seine)	1981	2.83	0.58	5	310	200
ii. <i>T. dussumieri</i>	Veraval (Trawls & gill nets)	1981	2.64	0.68	6	250	180
	Combined	1982	1.98	0.74	5	260	120
iii. <i>O. militaris</i>	Veraval (Trawl & gill nets)	1981	2.34	0.49	11	250	120
	Combined	1982	2.80	0.50	6	350	200
			1.74	0.37	10	270	200
			2.14	0.32	16	300	200

It may be noted from the above tables that the estimates of \bar{z} and $s\bar{z}$ are not differing very much between years. The differences between centres are also not very high.

iii. Estimation of 'M' the instantaneous natural mortality rate :

Effort data available, did not lead to estimation of effective effort particularly when data for more than one gear was considered. In multi-species fishery operated on by multi-gears the usual approach of

$$Z = M + qf \quad \dots \dots (6)$$

may not be possible. Hence a different approach is taken here, following Alagaraja (1984) to estimate 'M' directly from the length frequency data. Assuming one present survival of fish after they attain a length of $L_\infty - 0.5$ cm, the age T at which $L_\infty - 0.5$ is attained is obtained using

$$-\frac{1}{k} \log e \left(1 - \frac{L_\infty - 0.5}{L_\infty} \right) = T' - t_0 \approx T. \quad \dots \dots (7)$$

and using

$N_t / N_0 = 0.01 = e^{-MT} \dots \dots (8)$
an estimate of 'M' is arrived at. For example in the case of *T. thalassinus* $L_\infty = 755$ mm and $K = 0.36$. Hence

$$-\frac{1}{0.36} \log e \left(1 - \frac{750}{755} \right) \approx 14$$

$$N_{14} / N_0 = 0.01 = e^{-14M}$$

and $M = 0.33$

at one percent level of survival.

In this way 'M' for other species also have been estimated both at five and one percent levels of survival and the estimates are given below in table 5.

TABLE - 5
Estimates of 'M' instantaneous rate of natural mortality

Level of survival

Species	5%	1%
<i>T. thalassinus</i>	0.21	0.33
<i>T. tenuispinis</i>	0.50	0.76
<i>T. dussumieri</i>	0.15	0.22
<i>T. serratus</i>	0.15	0.21
<i>O. militaris</i>	0.50	0.77

For the present, estimates of 'M' at one percent level alone are considered since the values at five percent level appear to be low and at $L_\infty - 0.5$ cm length 'One percent survival' would not be far from the truth.

iv. Construction of yield isopleths

Considering the relatively long life span of cat fish and assuming isometric growth the yield equation

$$Y = FR W_\infty e^{-M(t_c - t_r)} \quad \dots \dots (8)$$

$$\sum_{n=0}^3 \frac{U_n e^{-nk(t_c - t_o)}}{F + M + nk} \quad \dots \dots (9)$$

(Gulland, 1969) has been considered here. Referring to the yield tables (Gulland 1969) for $M/k = 1.0$ yield isopleths have been drawn for $W_\infty = 4030$ gm and $t_r = 9$ mm. Eumetric fishing line BB' and the line AA' joining the maxima of yield-mesh curves are also indicated in fig. 1. Since for all the five species considered here, M/k remained more or less equal to unity, the same fig. 1 can be used for these species with varying multiplying factors according to their W_∞ and t_r values. These multiplying factors are given below in table 6.




Fig. 1. Yield isopleths for the five species of cat fishes

TABLE 6

Multiplying factors to obtain actual Y/R in grams from fig. 1

Centres Species	Walair	Mandapam	Cochin Fisheries Harbour	Mangalore	Veraval
<i>T. thalassinus</i>	1.14	1.08	1.14	—	1.14
<i>T. tenuispinis</i>	1.01	—	0.97	0.85	—
<i>T. serratus</i>	—	—	2.40	—	—
<i>T. dussumieri</i>	—	—	—	—	1.71
<i>O. militaris</i>	—	—	—	—	1.24

Yield-effort curves and yield-mesh curves have also been drawn (figs. 2-21) to get a clear picture of the status of fishery in each of these centres at the existing level of mesh size and effort.

From the yield isopleths (fig. 1) it can be

seen the maximum sustainable yield (MSY) of about 400 gm per recruit could be obtained at the level of $E=0.71$ and $C=0.66$ where $E=F/Z$ and $C=I_c / L_\infty$. The values of E and C for each species at each centre at the existing level of fishing are given in table 7.

TABLE 7

Levels of 'E' and 'C' at the existing level of fishing

Centre Species	Walair	Mandapam	Cochin Fisheries Harbour	Mangalore	Veraval
<i>T. thalassinus</i>	E 0.83 C 0.24	0.85 0.30	0.84 0.48	—	0.79 0.50
<i>T. tenuispinis</i>	E 0.71 C 0.36	—	0.76 0.50	0.72 0.46	—
<i>T. serratus</i>	E — C —	—	0.93 0.28	—	—
<i>T. dussumieri</i>	E — C —	—	—	—	0.91 0.30
<i>O. militaris</i>	E — C —	—	—	—	0.64 0.56

For *T. thalassinus* in all the four centres namely Waltair, Mandapam, Cochin Fisheries Harbour and Veraval the range for 'E' is 0.79 to 0.83 and for 'C' is 0.24 to 0.50. The values required for obtaining a MSY per recruit of about 400 gm are no where within the range. In other words to attain a MSY per recruit of 400 gm a reduction in effort and increase in mesh size are required.

In the case of *T. tenuispinis* 'E' values are nearer to the required level for 400 gm of MSY per recruit. However, I_c values are much lower than the required level indicating that for this species also mesh size has to be increased to attain 400 gm of MSY per recruit.

So far as *T. serratus* is concerned the rate of first capture is too low. Similar is the case with *T. dussumieri*. Hence for these two species reduction in effort and increase in mesh size are required to reach 400 gm MSY per recruit.

Only in the case of *O. militaris* the level of exploitation both for effort and size at first capture is nearer to the required level for obtaining MSY per recruit of about 400 gm.

It is hence clear that the level of exploitation in general was not favourable to the fishery of all the species except *O. militaris*. In order to see the effect of fishing at the existing level of effort on these stocks yield-effort curves have been drawn. Similarly to find out the impact of mesh size used in the fishery on these stocks, yield-yield-mesh curves have been drawn (figs. 2-21).

Effects of Fishing on *T. thalassinus*

At Waltair the yield-effort curve for the existing $C=0.24$ indicated that MSY per recruit




Fig. 2 Yield-effort curve for *T. thalassinus* at Waltair




Fig. 3. Yield-effort curve for *T. thalassinus* at Mandapam




Fig. 4. Yield-effort curve for *T. Thalassinus* at Cochin




Fig. 5 Yield-effort curve for *T. thalassinus* at Veraval

could be attained at $E=0.50$ which is far below the present level of exploitation where $E=0.83$ and the yield at $E=0.83$ is below half of that at $E=0.50$. Hence effort pressure should be considerably reduced to increase the returns from this stock at the present mesh size (Fig.2). For Mandapam at $C=0.30$ and the existing level of exploitation $E=0.85$ the yield per recruit is just above half of the MSY that could be obtained at $E=0.55$. Here also reduction in effort is suggested to increase the returns




Fig. 6. Yield-mesh curve for *T. thalassinus* at Walteir




Fig. 8. Yield-mesh curve for *T. thalassinus* at Cochin




Fig. 7. Yield-mesh curve for *T. thalassinus* at Mandapam

from this species (fig.3). Though similar is the trend for *T. thalassinus* at Cochin Fisheries Harbour and Veraval the existing level of




Fig. 9. Yield-mesh curve for *T. thalassinus* at Veraval

exploitation is not far away from the required level to obtain MSY and relatively less reduction of effort will improve the landings at these centres (fig. 4 and 5). Yield-mesh curve for this fishery at Waltair (fig. 6) indicates that the existing mesh size ($C = 0.24$) is far below the required one ($C = 0.70$) and this mesh has to be increased considerably to gain in returns from the fishery. Similar trend is seen at Mandapam (fig. 7). Regarding Cochin Fisheries Harbour (fig. 8) and Veraval (fig. 9), relatively less increase in mesh size will improve the landings.

Effects of Fishing on T. tenuispinis

The existing levels of effort at Waltair (fig. 10) Cochin Fisheries Harbour (fig. 11) and Mangalore (fig. 12) are not far above the required level for obtaining MSY. This is quite in contrast to the fishery of *T. thalassinus* where considerable reduction in effort is recommended to achieve MSY. This is due to higher levels of 'C' for *T. tenuispinis*. Yield-mesh curves indicate




Fig. 10 Yield-effort curve for *T. tenuispinis* at Waltair




Fig. 11 Yield-effort curve for *T. tenuispinis* at Cochin Fisheries Harbour




Fig. 12. Yield-effort curve for *T. tenuispinis* at Mangalore




Fig. 13. Yield-mesh curve for *T. tenuispinis* at Waltair

that size at first capture at the present levels of exploitation have to be increased so as to attain MSY at Waltair (fig. 13), Cochin Fisheries Harbour (fig. 14) and Mangalore (fig. 15).

Effects of Fishing on T. serratus, T. dussumieri and O. militaris

The fishery of *T. serratus* as observed at Cochin Fisheries Harbour is facing high fishing




Fig. 14. Yield-mesh curve for *T. tenuispinis*
at Cochin Fisheries Harbour




Fig. 15. Yield-mesh curve for *T. tenuispinis* at Mangalore




Fig. 16. Yield-effort curve for *T. serratus*
at Cochin Fisheries Harbour




Fig. 17. Yield-mesh curve for *T. serratus*
at Cochin Fisheries Harbour




Fig. 18. Yield-effort curve for *T. dussumieri* at Veraval




Fig. 19. Yield-mesh curve for *T. dussumieri* at Veraval




Fig. 21. Yield-mesh curve for *O. militaris* at Veraval




Fig. 20. Yield-effort curve for *O. militaris* at Veraval

pressure (fig. 16) and at this level the yield per recruit is only just above one third of MSY that could be obtained had the effort been reduced considerably. Yield-mesh curve also

indicates (fig. 17) that at this level of fishing pressure the size at first capture should be increased two and half times so as to get MSY without effecting the stocks.

The fishery of *T. dussumieri* at Veraval is very similar to that of *T. serratus* as seen above. The yield-effort curve (fig. 18) and yield-mesh curve (fig. 19) indicate the same trends as in those of *T. serratus*. Hence heavy reduction of fishing pressure at the present level of 'C' or steep increase in the level of 'C' for the existing fishing pressure along will lead to MSY.

However, in the case of *O. militaris* the present level of exploitation at Veraval appears to be ideal both in terms of effort (fig. 20) and the size at first capture (fig. 21).

CONCLUSIONS

The stocks of the four species of catfish considered here other than *O. militaris* were under heavy fishing pressure. It is hence indicated that in order to get MSY from these stocks either the fishing pressure is to be reduced at the existing level of 'C' (the index of the size at first capture) or the present level of 'C' is to be increased considerably at the existing level of fishing pressure.

Suggestion to increase mesh size so as to increase 'C' may not be appreciated as the trawl fishery is mainly aimed at shrimp fishing and shrimp fishery may not be profitably exploited at the increased level of mesh size. However, effort pressure may be brought down so as to attain MSY from these stocks.

Instead of studying catfish fishery from trawl landings in isolation, it would be better to study this fishery along with other stocks particularly the shrimps to arrive at final conclusion on the suitable levels of mesh size and effort pressure. As indicated above so far

as *O. militaris* is concerned present level of exploitation at Veraval appears to be ideal.

Annual catch estimates (Y in tonnes) for Waltair are based on the years 1978-80, for Mandapam on 1972-'76; for Cochin on 1981, Mangalore on 1982-'83, and for Veraval on 1981 and 1982. But for Mandapam, at other centres the estimates on average annual stock and average standing stock are comparable as these are based on the recent years.

For *T. thalassinus* Waltair region appears to be better when compared to other areas. However, for *T. tenuispinus* Managalore region indicates the maximum average annual stock. Regarding other species though region-wise comparision is not possible, from the present data base it can be said that the Veraval region hosts *T. duossumieri* and *O. militaris* more in abundance than *T. thalassinus*. Similarly Cochin region appears to be more favourable to *T. tenuispinus* than to *T. thalassinus* and *T. serratus* (Table 8).

TABLE 8

*Estimates of annual catch (Y in tonnes), average standing stock (Y/F in tonnes)
and average annual stock (Y/U in tonnes)*

Species	Centre	F	$\left(\frac{F}{Z} \frac{U}{(1 - e^{-Z})} \right)$	Y		Y/F (tonnes)	Y/U
I. <i>T. thalassinus</i>							
1.	Waltair	1.65	0.72	151	92	210	
2.	Mandapam	1.89	0.76	40	21	53	
3.	Cochin	1.73	0.73	116	67	159	
4.	Veraval	1.27	0.63	99	78	157	
II. <i>T. tenuispinus</i>							
1.	Waltair	1.83	0.65	158	86	243	
2.	Cochin	2.28	0.71	209	92	294	
3.	Mangalore	2.00	0.68	1008	504	1482	
III. <i>T. serratus</i>							
	Cochin	2.62	0.87	165	63	190	
IV. <i>T. duossumieri</i>							
	Veraval	2.12	0.82	438	207	534	
V. <i>O. militaris</i>							
	Veraval	1.37	0.57	275	200	482	

WORK SHEET I-A

A. *T. thalassinus* (Catch in numbers)

i. *Mandapam*

Mid point (mm)	1972	1973	1974	1975	1976	1977
	M. A.*	M. A.*	M. A.	M. A.*	M. A.*	M. A.*
70	6673	1721	448	4396	5982	
90	17629	16505	4816	4220	776	1568
110	25214	22383	6125	6886	3479	3875
130	24305	24760	10018	11036	7371	7563
150	24761	25223	16965	15914	11839	10817
170	26604	26961	20758	20130	13242	12961
190	29519	26766	22666	22658	13801	13624
210	21446	24056	24550	24206	13830	13283
230	18496	18995	25403	24299	12217	12805
250	14344	13738	22944	22781	12367	10704
270	8385	9560	19997	19640	7637	8709
290	5662	5802	15978	15680	6123	6389
310	3069	6434	11004	12198	5406	6315
330	10272	...	9611	9689	7415	5367
350	15931	...	8452	10175	3280	5580
370	19574	...	12463	...	6046	
390	17968		12285		1911	
410	9689		3388		1139	
430	1719		732		501	
450	460		90		1606	
470	789		0		31	
490	108		146		—	

ii. *Waltair* (Catch in numbers)

Mid point	1974	1975	M.A.*	1976
100	—			7936
140	—	4928		18681
180	21805	35333	16213	16407
220	13071	8379	15726	18907
260	7790	3465	4865	16917
300	3684	2750	2841	11617
340	2693	2308	2527	3229
380	527	2523	1964	808
420	—	1060	1321	
460		381	489	
500	—	25		

iii. *Veraval* (Catch in numbers)

Mid point	1981	1982
380	23765	43358
420	21144	36595
460	14038	18939
500	9394	5359
540	1226	4293

Note:-

1. M. A. * indicates three point moving average.
2. Bracketed portion alone is considered for the estimation of 'Z'.

T. thalassinus (Catch in numbers) contd.

iv Cochin

Mid point	Trawl	Gill net	1981	Total	M. A.*
110	994	—	994	994	
130	35790	—	35790	35790	13633
150	4114	—	4114	4114	13821
170	1517	42	1559	1559	2822
190	2511	281	2792	2792	4639
210	9189	377	9566	9566	8157
230	12034	78	12112	12112	12392
250	15121	377	15498	15498	26716
270	51949	588	52537	52537	34844
290	35159	1340	36499	36499	39608
310	26297	3492	29789	29789	35536
330	35735	4586	40321	40321	30233
350	15692	4997	20589	20589	30331
370	22282	7801	30083	30083	22204
390	6017	9923	15940	15940	20916
410	9052	7670	16722	16722	13315
430	3034	4248	7282	7282	8779
450	—	2334	2334	2334	3736
470	—	1593	1593	1593	1890
490	—	1744	1744	1744	1620
510	—	1522	1522	1522	1460
530	—	1115	1115	1115	1190
550	—	935	935	935	1235
570	—	1655	1655	1655	1418
590	—	1664	1664	1664	1620
610	—	1542	1542	1542	1526
630	—	1374	1374	1374	1398
650	—	1277	1277	1277	955
670	—	213	213	213	564
690	—	201	201	201	149
710	—	33	33	33	81
730	—	10	10	10	

Note: 1. M. A.* indicate three point moving average.

2. Bracketed portion alone is considered for estimation of 'Z'

WORK SHEET I - B

B. *T. tenuispinis* (Catch in numbers)

i. *Waltair*

Mid point (mm)	1974	M. A*	1975	M. A*	1976
100	4295	...			
140	26646	10961	2152	8734	11402
180	1943	21916	13858	12150	8534
220	37160	19322	10193	9366	6809
260	18865	27692	12399	6665	3370
300	27053	20617	5505	2682	1279
340	15933	15298	3091		310
380	2909	6349	450		
420	204				

ii. *Cochin*

Mid point	Purse seine	Trawl	Gill net	Total	M. A*
110	—	2238	—	2238	
130	—	21958	—	21958	11092
150	—	9080	—	9080	13366
170	—	9061	—	9061	11349
190	—	15906	—	15906	22387
210	—	42192	40	42233	27822
230	295	25367	141	25803	31430
250	1182	26730	876	28688	42630
270	5908	75795	1381	83086	58451
290	103304	72828	3262	86394	73683
310	8272	72426	7644	88342	70809
330	2954	67174	18327	88455	66327
350	591	59380	29515	89486	49061
370	295	20630	23559	24484	28389
390	—	5157	8830	13987	9942
410	—	4038	1080	5118	—
430	—	—	74	74	—
450	—	—	128	128	—
470	—	—	—	—	—
490	—	—	—	—	—
510	—	—	—	—	—
530	—	—	—	—	—
550	—	—	160	160	—

Note: 1. M. A. * indicates three point moving average.

2. Bracketed portion alone is considered for the estimation of 'Z'..

T. tenuispinis (Catch in numbers)

Mangalore		1982-83			
Mid point (mm)	Purse- seine	Trawl	Gill net	Total	M. A*
60	—	18482	—	18482	
70	—	255058	—	255058	124448
90	—	99805	—	99805	122897
110	13827	—	—	13827	40260
130	1887	5262	—	7149	70139
150	—	189440	—	189440	426926
170	—	1084189	—	1084189	784129
190	—	1078758	—	1078768	1259842
210	—	1616579	—	1616579	1111906
230	37	630345	—	640382	1001129
250	37	746389	—	746426	850307
270	9853	1153764	497	1164114	927855
290	29632	841002	2391	873025	787685
310	45392	271878	8645	325915	543195
330	273013	127460	30173	430646	415453
350	275856	179044	34897	489797	409271
370	189728	76763	40880	307371	319951
390	118360	10992	33333	162685	246431
410	155292	57946	56000	269238	173910
430	42477	10992	36339	29808	148035
450	—	—	12183	12183	95698
470	—	—	5672	5672	6335
490	—	—	1150	1150	—

Note: 1. M. A* indicates three point moving average.
 2. Bracketed portion alone is considered for the estimation of 'Z'.

WORK SHEET I — C

T. serratus (Catches in numbers)

Cochin Fisheries Harbour

Mid point (mm)	Purse seine	Trawl	Gill net	1981		
				I _t (mm)	M. A. ^{**} (Nos)	
210	—	3427	—			
230	20	—	10			
250	82	—	44			
270	61	—	41			
290	204	3427	35			
310	204	3427	159			
330	163	—	41			
350	122	3427	297			
370	61	—	570			
390	41	—	421			
410	41	—	197			
430	41	—	64			
450	—	—	112			
470	—	—	285			
490	—	—	1350			
510	—	3427	1150			
530	—	—	1790	250	54	
550	—	—	1762	270	116	
570	—	—	1309	290	156	
590	—	—	1243	310	190	
610	—	—	2019	330	163	
630	—	—	3216	350	115	
650	—	—	4226	370	75	
670	—	—	4139	390	48	
690	—	—	3541	410	41	
710	—	—	2132	430	—	
730	—	—	1681			
750	—	—	2674			
770	—	—	2468			
790	—	—	3194			
810	—	—	3252			
830	—	—	2728			
850	—	—	3115			
870	—	—	2667			
890	—	—	1709			
910	—	—	1665			
930	—	—	829			
950	—	—	774			
970	—	—	230			
990	—	—	26			

^{**} (M. A.) the bracketed portion shown below is taken for the estimation of 'Z'.

WORK SHEET I-D

T. dussumieri (Catch in numbers)

Veraval:

Mid point (mm)	Trawl	1981 gill net	M.A.1 (trawl only)	1982 Trawl	Gill net	Mid point (mm)	M. A.2 (Trawl only)
130	—	—	—	273	—	—	140
150	—	—	—	—	—	180	14174
170	—	—	—	273	—	220	32559
190	3273	—	—	289	—	260	48520
210	7102	—	9837	13727	—	300	40387
230	19137	—	12165	28657	—	340	22690
250	10267	—	13413	16516	—	380	7216
270	10844	—	10053	38913	—	420	1876
290	9097	112	9519	13952	—	460	1837
310	8658	—	7486	34392	—	—	—
330	4743	—	5035	16811	—	—	—
350	1705	575	2269	578	—	—	—
370	359	463	1185	969	—	—	—
390	1490	1842	1414	1369	—	—	—
410	2392	635	611	926	—	—	—
430	1894	911	1310	1184	—	—	—
450	868	1560	1027	1028	—	—	—
470	1455	1242	344	1075	—	—	—
490	85	1620	1226	3216	—	—	—
510	601	2050	995	3849	—	—	—
530	247	2384	1258	4464	—	—	—
550	—	1831	3259	4914	—	—	—
570	—	610	958	3537	—	—	—
590	—	2077	1769	6096	—	—	—
610	—	2538	2058	3689	—	—	—
630	—	3808	395	3802	—	—	—
650	—	3592	647	5107	—	—	—
670	—	2448	711	3105	—	—	—
690	—	4227	857	3390	—	—	—
710	—	2727	857	3579	—	—	—
730	—	1447	459	3090	—	—	—
750	—	620	84	1526	—	—	—
770	—	—	—	18	—	—	—
790	—	68	—	—	—	—	—
810	—	211	—	—	—	—	—

Note: 1. Data for gill nets do not appear to be the representative of gill net landings. Hence catch in numbers of three point moving average for trawls alone are considered for estimating 'Z'.

2. In this case successive classes have been merged forming a single wider class and for such wider classes three point moving average is taken.

3. Bracketed portion alone is considered for the estimation of 'Z'.

WORK SHEET 1-E

O. militaris (Catch in numbers)

Veraval

Mid point (mm)	1981			M. A.*	1982			M.A.*
	Trawl	Gill net	Total		Trawl	Gill net	Total	
190	—	—	—	—	—	—	—	—
210	1543	53	1596	—	5226	895	6121	—
230	13806	311	14117	18619	10429	958	11387	16090
250	38825	1318	40143	33980	29580	1181	39761	26154
270	46361	1319	47680	48571	32516	3799	36315	37600
290	56074	1816	57890	52365	41640	4085	48725	35973
310	49976	1549	51525	52157	23908	1970	25878	32127
330	44824	2233	47057	44979	22051	2727	24778	24414
350	35036	1319	36355	45682	16958	2929	22587	24226
370	52860	773	53633	35950	23125	2190	25315	21406
390	17532	357	17889	26749	13699	2617	16316	17648
410	8583	143	8726	10577	9461	1751	11312	11536
430	5022	94	5116	5445	6900	3079	6970	6776
450	2461	31	2492	2842	1884	153	2037	3263
470	917	—	917	1228	—	773	773	979
490	276	—	276	—	126	—	126	—

Note:

1. M. A.* indicates three point moving averages.
2. Bracketed portion alone is considered for the estimation of 'Z'.

CHAPTER NINE

REFERENCES

- ALAGARAJA, K. 1984. Simple methods for estimation of parameters for assessing exploited fish stocks. *Indian J. Fish.*, 31 (2) : 177-208.
- ALEXANDER, K. M. 1970. A study on the lipid distribution pattern in the muscles of two teleosts, *Arius duosumieri* and *Ophiocephalus striatus*. *Fishery Technology* 4 : 81-85.
- ANONYMOUS, 1976. *Pelagic Fishery Project, Prog. Rept No. 13* : 1-22.
- BALON, E. K. 1975. Reproductive guild of fishes: A proposal and definitions. *J. Fish. Res. Bd. Canada* 32 (6) : 821-864.
- BAPAT, S. V., V. M. DESHMUKH, B. KRISHNA-MOORTHI, C. MUTHIAH, P. V. KAGWADE, C. P. RAMAMRITHAM, K. J. MATHEW, S. KRISHNA PILLAI AND C. MUKUNDAN 1982. Fishery resources of Exclusive Economic Zone of the northwest coast of India. *Bulletin No. 33 CMFRI*.
- BLEEKER, P. 1847. *Siluroideorum bataviensium conspectus diagnosticus*. *Verh. Batav. Genootsch*, 22 : 9-375.
- CHACKO, P. I. 1949. Food and feeding habits of the fishes of the Gulf of Mannar. *Proc. Indian Acad. Sci* 29. B: 83-97.
- CHANDY, M. 1953. A key for the identification of the catfishes of the genus *Tachysurus* Lacepede with a catalogue of the specimens in the collection of the Indian Museum (Zool. Surv.) *Rec. Indian Mus.* 51(1) : 1-18.
- CHIDAMBARAM, K. 1942. Observations on the development of *Arius jelle* (Day). *Proc. Indian. Acad. Sci.*, 14 (B) : 502-508.
- DAN, S. S. 1977. Maturity, spawning and fecundity in the catfish, *Tachysurus tenuispinis* (Day). *Indian. J. Fish.* 24 : 96-106.
- DAN, S. S. 1980. Age and growth in the catfish. *Tachysurus tenuispinis* (Day). *Indian J. Fish.* 27 (1 & 2) : 220-235.
- DAN, S. S. AND P. MOJUMDER 1978. Length-weight relationship in catfish, *Tachysurus tenuispinis* (Day). *Indian J. Fish.* 25 : 23-28.
- DAY, F. 1878. *The fishes of India*, London, Wm. Dawson and sons. 778 PP.
- DEVANESAN D. W. AND K. CHIDAMBARAM. 1953. *The common food fishes of the Madras Presidency*. Madras Govt. Publication PP. 207.
- FISCHER, W. AND P. J. P. WHITEHEAD 1974. *FAQ Species Identification Sheets for fishery purposes*. Eastern Indian Ocean and Western Central Pacific (Fishing area 57 & 71).
- FISCHER, W. AND G. BIANCHI, 1984. *FAO Species Identification Sheets for fishery purposes*. Western Indian Ocean (Fishing area 51).
- FOWLER, H. W. 1941. The fishes of the Philippine and adjacent seas. *Bull. U. S. Nat. Mus.*, 13 (100) : 753-775.
- GOPAKUMAR, K AND V. SHENOY. 1977. Comparative evaluation of fish protein concentrate and functional fish protein concentrate from catfish. *Fish. Technolgy* 14 (1) : 84-88.
- GULLAND, J. A. 1969. Manual of methods for fish stock assessment. Part I. Fishery population analysis. *FAO Man. Fish. Sic. A*, 154 PP.

- GUNTER, G. 1947. Observations on breeding of the marine catfish, *Galeichthys felis*. *Copeia*, 4 : 217-223.
- GUNTHER, A. 1864. *Catalogue of the Fishes in the British Museum*. London, 5: 138-182.
- HARDENBERG, J. D. F. 1935. Miscellaneous notes on Indian fishes, IV. On the propagation of *Arius maculatus* (Thunb.) *Naturk. Tijdsch. Nedel Ind.*, 95 : 54-57.
- HERRE, A W. C. T. 1953. Check list of Philippine fishes. *Res. Rep. U. S. Wildl. Serv.*, 20 : 289—295.
- JAYARAM, K. C. AND G. R. DHANZE, 1978 a. Siluroid fishes of India, Burma and Ceylon. 22. A preliminary review of the genera of the family ariidae (Pisces : Siluroidea). *Matsya*, 4 : 42-51.
- JAYARAM, K. C. AND J. R. DHANZE 1978b. Siluroid fishes of India, Burma and Ceylon. 21. A note on the systematic position of *Tachysurus serratus* (Day) (Ariidae). *Bull Zool. Surv. India* 1 (2) : 203-205.
- JORDAN, D. S. 1963. *The genera of fishes and a classification of fishes*. Stanford University, California, 800 pp.
- KARTHA, K. N., S. D. DESHPANDE AND S. V. S. RAMA RAO 1973. On the results of bottom long lines operated off Veraval with particular reference to the selective action of bait and hooks used. *Fish. Tech.* 10 (2) : 105—109.
- KRISHNAMOORTHI, B. 1974. An assessment of the demersal fishery resources off the Andhra-Orissa coast based on exploratory trawling. *Indian J. Fish.* 21 (2) : 557—565.
- KRISHNAMOORTHI, B. 1978. A note on the trends of catfishes, *Tachysurus thalessinus* and *T. tenuispinis* based on the exploratory data for the period from 1966 to 1976. *Indian J. Fish.* 25 (1 & 2) : 268—270.
- KUTHALINGAM, M. D. K., P. MOJUMDER AND A. K. CHATTERJI 1973. Offshore fisheries resources of the Bay of Bengal-Sand Heads to Gopalpur. *Proc. Symp. Living Resources. Seas around India*. CMFR Institute. 338-364.
- KUTTY, M. P. AYYAPPA, V. SHENOY AND K. GOPALAKRISHNAN 1976. Chemical composition of *Tachysurus* spp. *Fish. Tech.* 13 (2) : 153-155.
- LEE, G. 1937. Oral gestation in the marine catfish, *Galeichthys felis*. *Copeia*, 1: 49-56.
- MENON, N. G. 1975. On buccal papilloma of *Tachysurus platystomus* (Day). *J. Mar. Biol. Ass. India*. 16 (1) (1974) : 317-320.
- MENON, N.G. 1979. Studies on the biology and fishery of the giant marine catfish, *Tachysurus thalessinus* (Ruppell). Ph. D. thesis to University of Cochin (unpublished).
- MENON, N. G. 1984 a. Observations on the intraovarian ova of a few tachysurids from Indian waters. *Indian J. Fish.* 31 (2) : 250-256.
- MENON, N. G. 1984 b. On the biology of the flat-mouthed catfish, *Tachysurus platystomus* (Day) from Mandapam. *Indian J. Fish.* 31 (?) : 293-308.
- MENON, N. G. 1984 c. On a new myxosporidian *Henneguya tachysuri* sp. nov. from the marine cat fish *Tachysurus thalessinus* (Ruppell) from the Gulf of Mannar. *J. Mar. Biol. Ass. India*, 21 (1 & 2) (1975) : 196-199.
- MOJUMDER, P. 1969. Food of *Tachysurus thalessinus* (Ruppell). *Indian J. Fish.* 16 : 161-169.
- MOJUMDER, P. 1971. Length-weight relationship in the catfish, *Tachysurus thalessinus* (Ruppell). *Indian J. Fish.* 18 : 179-182.

- MOJUMDER, P. 1977. Length frequency studies in the catfish, *Tachysurus thalassinus* (Ruppell) at Waltair during the years 1964-65 to 1960-70. *Indian J. Fish.*, 24 : 90-95.
- MOJUMDER, P. 1978. Maturity and spawning of the catfish, *Tachysurus thalassinus* (Ruppell) off Waltair coast. *Indian J. Fish.* 25 (1 & 2) : 109-121.
- MOJUMDER, P. AND S. S. DAN 1981. Studies on food and feeding habits of catfish, *Tachysurus tenuispinis* (Day). *Indian J. Fish.* 26. (1 & 2) (1979) : 115 - 224.
- MUNRO I. S. R. 1955. *The marine and freshwater fishes of Ceylon*. Canberra, Australia.
- NAGABHUSHANAM, A. K. 1966. A survey of the offshore demersal fisheries of the Andhra and Orissa coast with special references to the biological data collected during 1960. *Indian J. Fish.* 13 (1) : 359-376.
- NATARAJAN, S., K. C. GEORGE AND V. N. BANDE 1980. Echo location of fish. *Mar. Fish. Infor. Serv. T & E Ser.* 17 : 1-10, March 1980.
- OPPENHEIMER, J. R. 1970. Mouth breeding in fishes. *Anim. Behav.* 18 : 493-503.
- PANTULU, V. R. 1963. Studies on the age and growth, fecundity and spawning of *Osteogeneiosus militaris* (Linn.). *J. cons. int. Explor Mer.*, 28 (2) : 295-315.
- PAULY, D., N. DAVID AND J. INGLES 1981. ELEFAN II : users instruction and programme listing. *Mimeo. pag. war.*
- PILLAI, N. K. 1962. Copepods parasitic on south Indian fishes, 1. Caligidae. *Bull. Res. Inst. Univ. Kerala* 8 : 87-130.
- PILLAI, N. K. 1962. A review of the general *Hermilius*. *J. Zool. Soc. India* 14 : 179-187.
- PILLAI, T. V. R. AND K. K. GHOSH 1962. The bag-net fishery of the Hooghly-matla estuarine system. *Indian J. Fish.* 9 (1) : 71—99.
- RAO, K. SRINIVASA 1964. Food and feeding habits of fishes from trawl catches in the Bay of Bengal with observations on diurnal variation in the nature of feed. *Indian J. Fish.* 11 (2) : 227 — 314.
- RAO, K. V. 1969. Distribution pattern of the major exploited marine fishery resources of India. *Bulletin No. 6 CMFR*, pp 69.
- RAO, K. V. AND K. DORAIRAJ, 1968. Exploratory trawling off Goa by the Govt. of India fishing vessels. *Indian J. Fish.* 15 (1) : 1—14
- RAO, K. V., K. DORAIRAJ, P. V. KAGWADE AND D. M. PUNWANI, 1972. Results of the exploratory fishing operation of the Govt. of India vessels at Bombay base for the period 1961-67. *Proc. IPFC. Symposium on demersal fishes, 13th season III* : 402-430.
- RAO, K. V. N., M. KUMARAN AND J. SANKARASUBRAMANIAM, 1977. Resources of ribbon fishes and catfishes off the southwest coast of India. *Seafood Export J.* 9 (II): 9-26.
- SEKHARAN, K. V. 1973 a. The depth distribution off the catfishes *Tachysurus thalassinus* (Rupp.) and *Tachysurus tenuispinis* (Day) in the south western Bay of Bengal. *Indian J. Fish.* 20 (1) : 191-202.
- SEKHARAN, K. V. 1973 b. On the catfish resources of the coasts of Andhra Pradesh, Orissa and West Bengal. *Proc. Symp. Living Resources Seas around India*, CMFR Institute : 517-536.
- SEKHARAN, K. V. AND P. MOJUMDER. 1973. On the size of eggs found in the month of two males of catfish, *Tachysurus caelatus* (Val.). *J. mar. biol. Ass. India.* 15 (1) : 431-433.

- SEKHARAN, K. V., M. S. MUTHU, K. VENKATA-SUBBA RAO, V. RAMAMOHANA RAO, P. MOJUMDER AND S. REUBEN, 1973. Exploratory trawling on the continental shelf along the north-western Bay of Bengal. In *proceedings of the Symposium on the living resources of the seas around India*. *Spl. publi. C.M.F.R-Institute. Cochin* 280-337.
- SELVARAJ, G. S. D., K. GOPAKUMAR AND M. RAJAGOPALAN 1973. On the occurrence of Osteochondroma and osteoma in the marine catfish, *Tachysurus Jelte* (Day). *J. mar. biol. Ass. India*. 15 (2) : 571-576.
- SILAS, E. G., S. K. DHARMARAJA AND K. RENGARAJAN 1976. Exploited marine fishery resources of India—a synoptic survey with comments on potential resources, *Central Marine Fisheries Research Institute, Bull.* 27.
- SILAS E. G., P. PARAMESWARAN PILLAI, M. H. DHULKED, C. MUTHIAH AND G. SYDA RAO, 1980. Purse seine fishery - imperative need for regulation *Mar. Fish. Infor. Serv. T & E Ser. No. 24* : 1-9.
- SINGH, V. D. AND M. S. REGE. 1964. On the utilisation of catfish liver as a source of Vitamin A. *Curr. Sci.* 33 (12) : 371-72.
- SINGH, V. D. AND M. S. REGE. 1968. Observations on age and growth of *Tachysurus sona* (Ham.). *J. Bombay nat. Hist. Soc.*, 65 (1) : 75-87.
- SUSEELAN, C. AND K. V. SOMASEKHARAN NAIR. 1969. Food and Feeding habits of demersal fishes off Bombay. *Indian J. Fish.* 16 : 56-74.
- TILAK, R. 1965. The comparative morphology of the osteocranum and the Weberian apparatus of Tachysuride (Pisces: Siluroidea). *J. Zool.*, 146 : 150-174.
- THOLASILINGAM, T., K. C. GEORGE, M. G. DAYANANDAN, P. KARUNAKARAN NAIR, AND K. NANDAKUMARAN, 1973. Exploratory trawl fishing and ground fish resources along the Kerala coast and adjacent seas. *Proc. Symp. Living Resources. Seas. around India* CMFR Institute: 241-257.
- VALENCIENNES, M. A. 1840. IN : CUVIER, M AND VALENCIENNES, M. A. 1840. *Histoire naturelle des Poissons*. Paris, 15 : 28-206.
- VENKATARAMAN, G. 1960. Studies on the food and feeding relationships of the inshore fishes off Calicut on the Malabar coast. *Indian J. Fish.* 7 (1) : 275-303.
- WEBBER, M AND L. F. DE. BEAUFORT, 1913. *The fishes of the Indo-Australian archipelago*. Leiden, E. J. Brill Ltd. 2

WORK SHEET II

Estimation of 'Z' for A. thalassinus at Mandapam from the bracketed portions of work sheet I. A for the year 1972.

$l_\infty = 755 \text{ mm}$ and $k = 0.36$.

l_t	$l_\infty - l_t$	$\log(l_\infty - l_t)$	Their difference (A)	C_t	$\log C_t$	Their difference (B)	B/A	x_i
170	585	2.7672	0.0152	26961	4.4307	0.0033	0.22	
190	565	2.7520	0.0156	26756	4.4274	0.0462	2.96	
210	545	2.7364	0.0162	24054	4.3812	0.1026	6.33	
230	525	2.7202	0.0169	18985	4.2786	0.1407	8.32	
250	505	2.7033	0.0176	13738	4.1379	0.1574	8.94	
270	485	2.6857	0.0182	9560	3.9805	0.2169	11.92	
290	465	2.6675		5802	3.7636			
		$x = 6.45$	$n = 6$.					

$$\begin{aligned}
 s^2_x &= \frac{1}{n-1} \left[\sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2 / n \right] \\
 &= \frac{1}{5} [340.11 - (38.69)^2 / 6] \\
 &= \frac{1}{5} [340.11 - 249.49] \\
 &= 18.12
 \end{aligned}$$

Hence $s_x = \sqrt{18.12} = 4.26$ and $s_{\bar{x}} = 4.26/\sqrt{6} = 1.74$

Now $\bar{x} = \bar{z}/k$. Hence $\bar{z} = k\bar{x} = 0.36 \times 6.45 = 2.32$

$s_{\bar{x}} = s_{\bar{z}}/k$. Hence $s_{\bar{z}} = k s_{\bar{x}} = 0.36 \times 1.74 = 0.62$

Thus $\bar{z} = 2.32$ and $s_{\bar{z}} = 0.62$