



MARINE FISHERIES INFORMATION SERVICE

TECHNICAL AND
EXTENSION SERIES

No.11
September 1979

CENTRAL MARINE FISHERIES RESEARCH INSTITUTE
COCHIN, INDIA

INDIAN COUNCIL OF AGRICULTURAL RESEARCH

MARINE FISH PRODUCTION IN INDIA DURING JANUARY TO JUNE 1979*

The total marine fish production in India (excluding Andamans and Lakshadweep) during the half year ending June 1979 was provisionally estimated at 546,876 tonnes as against 530,056 tonnes recorded during the corresponding period in 1978, showing an increase of about 17,000 tonnes (3.17%). While the landings in West Bengal, Orissa, Pondicherry, Kerala, Karnataka, Goa and Maharashtra increased, Andhra Pradesh, Tamil Nadu and Gujarat recorded lower landings. The monthwise total landings of marine fish in the various maritime states of India and the specieswise catch details for the first half of 1979 are shown in Tables 1 and 2. In general, the monthly production during the half year showed a decreasing trend from the first month, recording the maximum in January and the minimum in June.

Pelagic and demersal groups of fishes

The pelagic and demersal groups of species contribute to the total landings. The pelagic group consists of *Chirocentrus*, sardines, *Hilsa* spp., anchovies and white baits, other clupeids, Bombay duck, *Hemiramphus* & *Belone*, flying fish, ribbon fish, carangids, mackerel, seer fish, tunnies, *Sphyraena*, mullets and *Bregmaceros* spp. The elasmobranchs, eels, cat fishes, lizard fishes, perches, red mullets, polynemids, sciaenids, silver bellies, *Lactarius*, pomfrets, soles, prawns, lobsters and cephalopods form the demersal group. The statewise distribution of pelagic and demersal groups of fishes is shown in Fig. 1.

Kerala, Tamil Nadu, Karnataka, Maharashtra, Gujarat and Andhra Pradesh recorded higher catch of pelagic fishes. The demersal fishes are predominantly caught in the states of Maharashtra, Tamil Nadu, Gujarat, Kerala, Andhra Pradesh and Karnataka. While Kerala accounted for the highest catch of pelagic fishes, Maharashtra contributed the maximum catch of demersal fishes.

Statewise production

West Bengal

The total marine fish production in West Bengal increased by about 3,000 tonnes as compared to the corresponding period in 1978 (Table 1). While the landings of sciaenids, other clupeids, *Harpodon nehereus*, pomfrets, *Chirocentrus*, *Hilsa ilisha* and perches showed significant increase, *Leiognathus* and non-penaeid prawns recorded poor landings (Table 3). While the maximum landings were recorded in the month of January, minimum catch was seen in March.




Fig. 1. Statewise pelagic and demersal catches during January to June 1979

Orissa

An increase of about 6,600 tonnes in the total landings was noticed in Orissa over that of the corresponding half year in 1978 (Table 1). The catch of *Hilsa ilisha*, other sardines, elasmobranchs, pomfrets, seer fish, *Chirocentrus*, cat fishes, other clupeids and *Leiognathus* increased substantially. The landings of *Anchoviella*, tunnies, perches and penaeid prawns, however, were comparatively poor. The monthwise and specieswise landings of marine fish in Orissa during the first half of

*Prepared by the Fishery Resources Assessment Division.

Table 1. Statewise and monthwise total marine fish production in India (excluding Andamans and Lakshadweep) during the half year ending June 1979* (in tonnes)

| Sl. No. | Name of State | January | February | March | April | May | June | Total | Total for January to June 1978 |
|--------------|----------------|-----------------|---------------|-----------------|---------------|---------------|---------------|-----------------|--------------------------------------|
| 1. | West Bengal | 2,572 | 402 | 250 | 329 | 316 | 387 | 4,256 | 1,372 |
| 2. | Orissa | 4,876 | 2,893 | 1,892 | 1,674 | 1,250 | 865 | 13,450 | 6,826 |
| 3. | Andhra Pradesh | 11,110 | 6,839 | 9,357 | 7,584 | 3,452 | 2,507 | 40,849 | 43,123 |
| 4. | Tamil Nadu | 19,274 | 19,041 | 23,899 | 16,348 | 14,529 | 19,153 | 1,12,244 | 1,12,947 |
| 5. | Pondicherry | 214 | 621 | 214 | 710 | 509 | 1,422 | 3,690 | 2,760 |
| 6. | Kerala | 20,368 | 18,691 | 22,844 | 20,907 | 31,281 | 8,407 | 1,22,498 | 1,14,913 |
| 7. | Karnataka | 10,478 | 9,414 | 10,610 | 8,254 | 6,990 | 2,742 | 48,488 | 47,018 |
| 8. | Goa | 2,488 | 1,297 | 3,186 | 1,911 | 717 | 37 | 9,636 | 5,784 |
| 9. | Maharashtra | 24,668 | 23,254 | 19,130 | 25,426 | 22,609 | 2,589 | 1,17,676 | 1,02,066 |
| 10. | Gujarat | 28,016 | 9,260 | 18,525 | 12,896 | 3,961 | 1,431 | 74,089 | 93,247 |
| TOTAL | | 1,24,064 | 91,712 | 1,09,907 | 96,039 | 85,614 | 39,540 | 5,46,876 | 5,30,056 |

* Provisional

1979 is shown in Table 4. The maximum and minimum landings were recorded in January and June respectively.

Andhra Pradesh

The total landings in Andhra Pradesh during the first half of 1979 showed a decline of about 2,000 tonnes (Table 1). While the landings of cat fishes, *Caranx*, *Leiognathus* spp, seer fish, *Thrissocles*, other clupeids, perches, *Saurida* & *Saurus* and non-penaeid prawns increased significantly, the catch of *Anchoviella*, other sardines, elasmobranchs, eels, other *Hilsa* and mackerel recorded poor landings (Table 5). While the maximum landings were recorded in the month of January, the minimum were seen in June.

Tamil Nadu

A minor decline of about 700 tonnes in the total landings was noticed in Tamil Nadu as compared to the half year of 1978 (Table 1). A significant increase in the landings of *Leiognathus* spp, other sardines, *Anchoviella*, sciaenids, mackerel, flying fish, other *Hilsa* and seer fish was noticed. The catch of elasmobranchs, perches, ribbon fish, crabs and other crustaceans and cat fishes, however, was comparatively poor. Table 6 shows the details of catch for the half year. While the maximum catch was recorded in March 1979, in the month of May minimum catch was seen.

Pondicherry

The total landings increased by about 900 tonnes in Pondicherry as compared to the first half of 1978. (Table 1). While the catch of flying fish, perches, *Leiognathus* spp, mackerel, penaeid prawns, *Chirocentrus* and other clupeids increased substantially, *Anchoviella*, other *Hilsa*, other sardines and sciaenids recorded poor catch (Table 7). The maximum landings were recorded in June 1979. The minimum catch was recorded in the months of January and March 1979.

Kerala

An increase of about 7,600 tonnes was noticed in the total landings in Kerala as compared to the corresponding period in 1978 (Table 1). A significant increase in the landings of oil sardines, mackerel, perches, *Leiognathus* spp, tunnies, crabs and other crustaceans was noticed. The landings of penaeid prawns, soles, sciaenids, ribbon fish, *Anchoviella*, *Chirocentrus*, cat fishes, elasmobranchs, *Saurida* & *Saurus* and cephalopods, however, were poor. Table 8 gives the monthwise and specieswise estimates of marine fish production in Kerala for the half year ending June 1979. While the total landings were maximum in the month of May, the same decreased to minimum in June.

Table 2. Statewise composition of marine fish landings in India (excluding Andamans & Lakshadweep) during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | West Bengal | Orissa | Andhra Pradesh | Tamil Nadu | Pondi- cherry | Kerala | Karnataka | Goa | Maha- rashtra | Gujarat | Total |
|------------|----------------------------------|----------------|--------|-------------------|---------------|------------------|----------|-----------|-------|------------------|---------|----------|
| 1. | Elasmobranchs | 62 | 1,300 | 3,489 | 6,688 | 97 | 2,854 | 571 | 534 | 5,081 | 2,407 | 23,083 |
| 2. | Eels | — | — | 146 | 87 | 70 | 3 | — | 4 | 2,191 | 1,719 | 4,220 |
| 3. | Cat fishes | 94 | 501 | 2,568 | 2,074 | 27 | 4,084 | 1,314 | 441 | 4,634 | 2,926 | 18,663 |
| 4. | Chirocentrus | 195 | 557 | 424 | 860 | 95 | 650 | 53 | 58 | 478 | 1,091 | 4,461 |
| 5. | (a) Oil sardine | — | — | 743 | — | 37,676 | 9,656 | 658 | — | — | — | 48,733 |
| | (b) Other sardines | — | 1,494 | 3,940 | 17,227 | 515 | 4,017 | 3,557 | 485 | 186 | — | 31,421 |
| | (c) <i>Hilsa ilisha</i> | 190 | 3,623 | — | 5 | — | — | 2 | — | 345 | 186 | 4,351 |
| | (d) Other <i>Hilsa</i> | 32 | 293 | 628 | 2,255 | 4 | 34 | 26 | 14 | 109 | 1,440 | 4,835 |
| | (e) <i>Anchoviella</i> | 8 | 16 | 3,046 | 5,062 | 81 | 3,677 | 851 | — | 42 | — | 12,783 |
| | (f) <i>Thrissocles</i> | 126 | 120 | 1,020 | 3,636 | 212 | 434 | 210 | 815 | 1,115 | 316 | 8,004 |
| | (g) Other clupeids | 347 | 497 | 1,600 | 2,608 | 139 | 368 | 740 | 289 | 5,974 | 2,015 | 14,577 |
| 6. | (a) <i>Harpodon nehereus</i> | 321 | 96 | 72 | 1 | — | 1 | 1 | — | 14,333 | 10,762 | 25,587 |
| | (b) <i>Saurida & Saurus</i> | — | 50 | 867 | 667 | 100 | 1,124 | 156 | .96 | 1,515 | 5 | 4,580 |
| 7. | <i>Hemirhamphus & Belone</i> | — | 28 | 31 | 395 | — | 144 | 9 | 11 | 11 | — | 629 |
| 8. | Flying fish | — | 4 | 47 | 1,402 | 495 | — | — | 1 | — | — | 1,949 |
| 9. | Perches | 186 | 39 | 2,009 | 2,669 | 333 | 2,332 | 103 | 181 | 1,292 | 690 | 9,834 |
| 10. | Red mullets | — | — | 170 | 824 | 94 | 126 | 29 | — | 535 | — | 1,798 |
| 11. | Polynemids | 55 | 242 | 996 | 181 | — | 26 | 2 | — | 836 | 136 | 2,474 |
| 12. | Sciaenids | 610 | 281 | 2,500 | 9,392 | 89 | 2,829 | 1,040 | 765 | 10,138 | 13,523 | 41,167 |
| 13. | Ribbon fish | 175 | 176 | 2,380 | 1,626 | 47 | 2,623 | 432 | 454 | 5,528 | 2,048 | 15,489 |
| 14. | (a) <i>Caranx</i> | 70 | 216 | 2,253 | 1,465 | 65 | 2,290 | 284 | 640 | 1,205 | 79 | 8,567 |
| | (b) <i>Chorinemus</i> | 43 | 260 | 233 | 570 | 3 | 44 | 6 | 49 | 91 | 82 | 1,381 |
| | (c) <i>Trachynotus</i> | — | — | — | 152 | — | — | 13 | — | — | — | 165 |
| | (d) Other carangids | — | — | 18 | 14 | — | — | — | — | 41 | — | 73 |
| | (e) <i>Coryphaena</i> | — | — | — | 33 | 2 | 48 | — | — | 11 | — | 94 |
| | (f) <i>Elacate</i> | — | — | — | 280 | — | 23 | 9 | 1 | — | — | 313 |
| 15. | (a) <i>Leiognathus</i> | 22 | 496 | 2,142 | 25,207 | 236 | 2,056 | 672 | 583 | 309 | — | 31,723 |
| | (b) <i>Gazza</i> | — | — | — | 130 | — | — | — | — | — | — | 130 |
| 16. | <i>Lactarius</i> | — | — | 378 | 314 | 5 | 143 | 140 | 146 | 299 | 209 | 1,634 |
| 17. | Pomfrets | 287 | 1,194 | 1,112 | 518 | 12 | 774 | 98 | 74 | 4,381 | 2,897 | 11,347 |
| 18. | Mackerel | — | 165 | 1,031 | 2,137 | 227 | 16,703 | 10,766 | 205 | 683 | 41 | 31,958 |
| 19. | Seer fish | 69 | 690 | 2,112 | 2,241 | 53 | 1,209 | 404 | 295 | 446 | 1,475 | 8,994 |
| 20. | Tunnyes | — | 18 | 218 | 446 | 1 | 11,409 | 676 | 5 | 414 | 98 | 13,285 |
| 21. | <i>Sphyraena</i> | — | 2 | 17 | 753 | 9 | 276 | 30 | 7 | 30 | — | 1,124 |
| 22. | <i>Mugil</i> | — | 22 | 137 | 109 | 18 | 30 | 1 | 5 | 19 | 225 | 566 |
| 23. | <i>Bregmaceros</i> | — | — | — | — | — | — | — | — | 1 | 254 | 255 |
| 24. | Soles | — | 8 | .247 | 1,098 | 78 | 1,767 | 215 | 235 | 787 | 123 | 4,558 |
| 25. | (a) Penaeid prawns | 124 | 69 | 2,482 | 6,223 | 220 | 12,668 | 4,208 | 1,276 | 18,741 | 4,394 | 50,405 |
| | (b) Non-penaeid prawns | 102 | 32 | 589 | 183 | 56 | 24 | 2 | 55 | 31,003 | 799 | 32,845 |
| | (c) Lobsters | — | — | 84 | 143 | — | 11 | 15 | — | 496 | 113 | 862 |
| | (d) Other crustaceans | — | 4 | 207 | 2,203 | 72 | 3,198 | 2,483 | 824 | 86 | 500 | 9,577 |
| 26. | Cephalopods | — | 1 | 203 | 561 | 20 | 286 | 52 | 106 | 1,558 | 4,518 | 7,305 |
| 27. | Miscellaneous | 1,138 | 956 | 1,453 | 9,062 | 215 | 6,537 | 9,662 | 324 | 2,712 | 19,018 | 51,077 |
| TOTAL | | 4,256 | 13,450 | 40,849 | 1,12,244 | 3,690 | 1,22,498 | 48,488 | 9,636 | 1,17,676 | 74,089 | 5,46,876 |

Karnataka

In Karnataka, the total landings increased by about 1,500 tonnes when compared to the landings recorded during the first half of 1978 (Table 1). While the landings of mackerel, other sardines, crabs and other crusta-

ceans, tunnies, *Thrissocles* spp, other clupeids and ribbon fish were comparatively higher, the catch of oil sardine, penaeid prawns, pomfrets and elasmobranchs was poor (Table 9). The maximum catch was recorded in the months of January and March whereas the minimum catch was accounted for in June.

Table 3. Composition of marine fish landings in West Bengal during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of Fish | Jan. | Feb. | March | April | May | June | Total |
|--------------|----------------------------------|--------------|------------|------------|------------|------------|------------|--------------|
| 1. | Elasmo-brancls | 22 | — | — | 26 | — | 14 | 62 |
| 2. | Eels | — | — | — | — | — | — | — |
| 3. | Cat fishes | 88 | — | — | 6 | — | — | 94 |
| 4. | <i>Chirocentrus</i> | 86 | 30 | 6 | 24 | 24 | 25 | 195 |
| 5. a) | Oil sardine | — | — | — | — | — | — | — |
| b) | Other sardines | — | — | — | — | — | — | — |
| c) | <i>Hilsa ilisha</i> | 112 | — | 6 | 18 | — | 54 | 190 |
| d) | Other <i>Hilsa</i> | — | — | — | 12 | — | 20 | 32 |
| e) | <i>Ancho-viella</i> | — | — | 8 | — | — | — | 8 |
| f) | <i>Thris-socles</i> | 98 | 6 | 4 | 6 | 12 | — | 126 |
| g) | Other clupeids | 150 | 38 | 42 | 34 | 51 | 32 | 347 |
| 6. a) | <i>Harpodon nehereus</i> | 190 | 36 | 34 | 6 | 32 | 23 | 321 |
| b) | <i>Saurida & Saurus</i> | — | — | — | — | — | — | — |
| 7. | <i>Hemirhamphus & Belone</i> | — | — | — | — | — | — | — |
| 8. | Flying fish | — | — | — | — | — | — | — |
| 9. | Perches | 160 | — | — | 18 | 4 | 4 | 186 |
| 10. | Red mullets | — | — | — | — | — | — | — |
| 11. | Polynemids | — | — | 4 | 24 | 8 | 19 | 55 |
| 12. | Sciaenids | 492 | 32 | 20 | 22 | 24 | 20 | 610 |
| 13. | Ribbon fish | 78 | 36 | 16 | 4 | 26 | 15 | 175 |
| 14. a) | <i>Carenx</i> | 62 | — | 4 | 4 | — | — | 70 |
| b) | <i>Chorinemus</i> | — | — | — | 17 | 8 | 18 | 43 |
| c) | <i>Trachy-notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | — | — | — | — | — | — | — |
| e) | <i>Coryphaena</i> | — | — | — | — | — | — | — |
| f) | <i>Elacate</i> | — | — | — | — | — | — | — |
| 15. a) | <i>Leiognathus</i> | — | — | 4 | — | 18 | — | 22 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | — | — | — | — | — | — | — |
| 17. | Pomfrets | 12 | 56 | 48 | 56 | 34 | 81 | 287 |
| 18. | Mackerel | — | — | — | — | — | — | — |
| 19. | Seer fish | — | — | — | 24 | 17 | 28 | 69 |
| 20. | Tunnies | — | — | — | — | — | — | — |
| 21. | <i>Sphyraena</i> | — | — | — | — | — | — | — |
| 22. | <i>Mugil</i> | — | — | — | — | — | — | — |
| 23. | <i>Breg-maceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | — | — | — | — | — | — | — |
| 25. a) | Penaeid prawns | 68 | 50 | 6 | — | — | — | 124 |
| b) | Non-penaeid prawns | 80 | 22 | — | — | — | — | 102 |
| c) | Lobsters | — | — | — | — | — | — | — |
| d) | Other crustaceans | — | — | — | — | — | — | — |
| 26. | Cephalopods | — | — | — | — | — | — | — |
| 27. | Miscellaneous | 874 | 96 | 48 | 28 | 58 | 34 | 1,138 |
| TOTAL | | 2,572 | 402 | 250 | 329 | 316 | 387 | 4,256 |

Table 4. Composition of marine fish landings in Orissa during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Jan. | Feb. | March | April | May | June | Total |
|--------------|----------------------------------|--------------|--------------|--------------|--------------|--------------|------------|---------------|
| 1. | Elasmo-brancls | 286 | 314 | 268 | 179 | 148 | 105 | 1,300 |
| 2. | Eels | — | — | — | — | — | — | — |
| 3. | Cat fishes | 185 | 196 | 57 | 20 | 41 | 2 | 501 |
| 4. | <i>Chirocentrus</i> | 57 | 102 | 105 | 106 | 120 | 67 | 557 |
| 5. a) | Oil sardine | — | — | — | — | — | — | — |
| b) | Other sardines | 386 | 587 | 170 | 265 | 84 | 2 | 1,494 |
| c) | <i>Hilsa ilisha</i> | 2,386 | 556 | 124 | 288 | 147 | 122 | 3,623 |
| d) | Other <i>Hilsa</i> | 72 | 94 | 12 | 12 | 62 | 41 | 293 |
| e) | <i>Ancho-viella</i> | — | 16 | — | — | — | — | 16 |
| f) | <i>Thris-socles</i> | 44 | 4 | 39 | — | 20 | 13 | 120 |
| g) | Other clupeids | 120 | 124 | 37 | 97 | 65 | 54 | 497 |
| 6. a) | <i>Harpodon nehereus</i> | 70 | 2 | 4 | 6 | 12 | 2 | 96 |
| b) | <i>Saurida & Saurus</i> | 49 | 1 | — | — | — | — | 50 |
| 7. | <i>Hemirhamphus & Belone</i> | 25 | 3 | — | — | — | — | 28 |
| 8. | Flying fish | 1 | 3 | — | — | — | — | 4 |
| 9. | Perches | 16 | 1 | 3 | 18 | — | 1 | 39 |
| 10. | Red mullets | — | 78 | 20 | 24 | 44 | 38 | 242 |
| 11. | Polynemids | 38 | 17 | 102 | 50 | 22 | 46 | 281 |
| 12. | Sciaenids | 44 | — | — | — | — | — | — |
| 13. | Ribbon fish | 26 | 8 | 31 | 54 | 33 | 24 | 176 |
| 14. a) | <i>Caranx</i> | 10 | 169 | 8 | 14 | — | 15 | 216 |
| b) | <i>Chorinemus</i> | 56 | 79 | 22 | 35 | 48 | 20 | 260 |
| c) | <i>Trachy-notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | — | — | — | — | — | — | — |
| e) | <i>Coryphaena</i> | — | — | — | — | — | — | — |
| f) | <i>Elacate</i> | — | — | — | — | — | — | — |
| 15. a) | <i>Leiognathus</i> | 16 | 25 | 332 | 53 | 34 | 36 | 496 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | — | 233 | 147 | 218 | 186 | 120 | 1,194 |
| 17. | Pomfrets | 290 | — | — | — | — | — | 165 |
| 18. | Mackerel | 3 | 8 | 154 | — | — | — | 690 |
| 19. | Seer fish | 46 | 187 | 187 | 105 | 101 | 64 | 18 |
| 20. | Tunnies | 10 | 8 | — | — | — | — | 2 |
| 21. | <i>Sphyraena</i> | — | 2 | — | — | — | — | 22 |
| 22. | <i>Mugil</i> | 22 | — | — | — | — | — | — |
| 23. | <i>Breg-maceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | — | — | — | — | 8 | — | 8 |
| 25. a) | Penaeid prawns | 28 | — | — | 12 | 1 | 28 | 69 |
| b) | Non-penaeid prawns | 30 | — | 2 | — | — | — | 32 |
| c) | Lobsters | — | — | — | — | — | — | — |
| d) | Other crustaceans | — | — | — | — | 2 | 2 | 4 |
| 26. | Cephalopods | — | 1 | — | — | — | — | 1 |
| 27. | Miscellaneous | 559 | 76 | 68 | 118 | 72 | 63 | 956 |
| TOTAL | | 4,876 | 2,893 | 1,892 | 1,674 | 1,250 | 865 | 13,450 |

Table 5. Composition of marine fish landings in Andhra Pradesh during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|--------------|--|--------|-------|-------|-------|-------|-------|--------|
| 1. | Elasmo-bran- | 1,620 | 493 | 563 | 511 | 109 | 193 | 3,489 |
| 2. | Eels | 26 | 14 | 22 | 53 | 20 | 11 | 146 |
| 3. | Cat fishes | 698 | 230 | 1,335 | 173 | 60 | 72 | 2,568 |
| 4. | <i>Chiro-</i> <i>centrus</i> | 198 | 57 | 126 | 26 | 2 | 15 | 424 |
| 5. a) | Oil sardine | — | — | — | — | — | — | — |
| b) | Other sardines | 1,142 | 785 | 1,044 | 551 | 402 | 16 | 3,940 |
| c) | <i>Hilsa</i> <i>ilisha</i> | — | — | — | — | — | — | — |
| d) | Other <i>Hilsa</i> | 477 | 94 | 20 | 8 | 29 | — | 628 |
| e) | <i>Ancho-</i> <i>viella</i> | 875 | 404 | 641 | 1,000 | 85 | 41 | 3,046 |
| f) | <i>Thris-</i> <i>socles</i> | 234 | 315 | 177 | 108 | 64 | 122 | 1,020 |
| g) | Other clupeids | 602 | 456 | 293 | 98 | 57 | 94 | 1,600 |
| 6. a) | <i>Harpodon</i> <i>nehereus</i> | 3 | 26 | 15 | 26 | 2 | — | 72 |
| b) | <i>Saurida</i> & <i>Saurus</i> | 500 | 42 | 74 | 85 | 99 | 67 | 867 |
| 7. | <i>Hemir-</i> <i>hamphus</i> & <i>Belone</i> | 11 | — | 1 | — | — | 19 | 31 |
| 8. | Flying fish | 47 | — | — | — | — | — | 47 |
| 9. | Perches | 693 | 366 | 336 | 467 | 112 | 35 | 2,009 |
| 10. | Red mullets | 32 | 35 | 24 | 39 | 30 | 10 | 170 |
| 11. | Polynemids | 99 | 90 | 86 | 608 | 98 | 15 | 996 |
| 12. | Sciaenids | 463 | 442 | 528 | 563 | 226 | 278 | 2,500 |
| 13. | Ribbon fish | 550 | 208 | 168 | 1,081 | 166 | 207 | 2,380 |
| 14. a) | <i>Caranx</i> | 89 | 653 | 1,106 | 290 | 25 | 90 | 2,253 |
| b) | <i>Chori-</i> <i>nemus</i> | 64 | 5 | 135 | 24 | 2 | 3 | 233 |
| c) | <i>Trachy-</i> <i>notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | 3 | — | 2 | 5 | 5 | 3 | 18 |
| e) | <i>Cory-</i> <i>phaena</i> | — | — | — | — | — | — | — |
| f) | <i>Elacate</i> | — | — | — | — | — | — | — |
| 15. a) | <i>Leiog-</i> <i>nathus</i> | 181 | 390 | 426 | 691 | 237 | 217 | 2,142 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | 71 | 27 | 91 | 102 | 8 | 79 | 378 |
| 17. | Pomfrets | 159 | 89 | 568 | 124 | 45 | 127 | 1,112 |
| 18. | Mackerel | 58 | 25 | 592 | 290 | 65 | 1 | 1,031 |
| 19. | Seer fish | 938 | 135 | 320 | 54 | 521 | 144 | 2,112 |
| 20. | Tunnies | 91 | 11 | — | 19 | 97 | — | 218 |
| 21. | <i>Sphyraena</i> | — | 4 | 8 | 4 | 1 | — | 17 |
| 22. | <i>Mugil</i> | 6 | 131 | — | — | — | — | 137 |
| 23. | <i>Breg-</i> <i>maceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | 86 | 33 | 41 | 22 | 39 | 26 | 247 |
| 25. a) | Penaeid prawns | 360 | 993 | 178 | 238 | 381 | 332 | 2,482 |
| b) | Non- penaeid prawns | 158 | 1 | 31 | 79 | 247 | 73 | 589 |
| c) | Lobsters | 2 | 1 | 22 | 17 | 32 | 10 | 84 |
| d) | Other crusta- | — | — | — | — | — | — | — |
| 26. | Cepha- | 30 | 6 | 19 | 14 | 27 | 111 | 207 |
| 27. | lopods | 21 | 20 | 45 | 44 | 54 | 19 | 203 |
| Miscel- | lanous | 523 | 258 | 320 | 170 | 105 | 77 | 1,453 |
| TOTAL | | 11,110 | 6,839 | 9,357 | 7,584 | 3,452 | 2,507 | 40,849 |

Table 6. Composition of marine fish landings in Tamil Nadu during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|--------------|--|--------|--------|--------|--------|--------|--------|----------|
| 1. | Elasmo- branchs | 1,011 | 817 | 1,803 | 1,541 | 868 | 648 | 6,688 |
| 2. | Eels | 55 | 7 | 4 | 5 | 14 | 2 | 87 |
| 3. | Cat fishes | 394 | 379 | 694 | 186 | 290 | 131 | 2,074 |
| 4. | <i>Chiro-</i> <i>centrus</i> | 376 | 109 | 72 | 74 | 102 | 127 | 860 |
| 5. a) | Oil sardine | 71 | 111 | 24 | 5 | 532 | — | 743 |
| b) | Other sardines | 4,401 | 2,549 | 3,759 | 2,539 | 2,120 | 1,859 | 17,227 |
| (c) | <i>Hilsa</i> <i>ilisha</i> | — | — | — | — | 5 | — | 5 |
| d) | Other <i>Hilsa</i> | 459 | 650 | 675 | 42 | 294 | 135 | 2,255 |
| e) | <i>Ancho-</i> <i>viella</i> | 279 | 1,539 | 1,108 | 538 | 758 | 840 | 5,062 |
| f) | <i>Thris-</i> <i>socles</i> | 516 | 422 | 1,261 | 474 | 616 | 347 | 3,636 |
| g) | Other clupeids | 778 | 289 | 974 | 246 | 156 | 165 | 2,608 |
| 6. a) | <i>Harpodon</i> <i>nehereus</i> | 1 | — | — | — | — | — | 1 |
| b) | <i>Saurida</i> & <i>Saurus</i> | 275 | 59 | 73 | 93 | 102 | 65 | 667 |
| 7. | <i>Hemir-</i> <i>hamphus</i> & <i>Belone</i> | 115 | 105 | 44 | 58 | 60 | 13 | 395 |
| 8. | Flying fish | — | 2 | 2 | 7 | 491 | 900 | 1,402 |
| 9. | Perches | 342 | 549 | 738 | 324 | 309 | 407 | 2,669 |
| 10. | Red mullets | 149 | 176 | 164 | 178 | 107 | 50 | 824 |
| 11. | Polynemids | 85 | 49 | 24 | 19 | 2 | 2 | 181 |
| 12. | Sciaenids | 1,464 | 2,028 | 1,998 | 1,485 | 1,055 | 1,362 | 9,392 |
| 13. | Ribbon fish | 58 | 150 | 58 | 17 | 58 | 1,285 | 1,626 |
| 14. a) | <i>Caranx</i> | 201 | 139 | 223 | 301 | 463 | 138 | 1,465 |
| b) | <i>Chori-</i> <i>nemus</i> | 87 | 40 | 199 | 44 | 133 | 67 | 570 |
| c) | <i>Trachy-</i> <i>notus</i> | 12 | — | 109 | 6 | 25 | — | 152 |
| d) | Other carangids | — | 12 | — | — | 2 | — | 14 |
| e) | <i>Cory-</i> <i>phaena</i> | 10 | — | 3 | 10 | 8 | 2 | 33 |
| f) | <i>Elacate</i> | 37 | 70 | 61 | 50 | 60 | 2 | 280 |
| 15. a) | <i>Leiog-</i> <i>nathus</i> | 3,307 | 4,976 | 5,343 | 3,403 | 2,336 | 5,842 | 25,207 |
| b) | <i>Gazza</i> | 33 | 70 | 22 | 2 | 3 | — | 130 |
| 16. | <i>Lactarius</i> | 135 | 2 | 42 | 6 | 66 | 63 | 314 |
| 17. | Pomfrets | 63 | 40 | 12 | 288 | 12 | 103 | 518 |
| 18. | Mackerel | 313 | 148 | 464 | 555 | 611 | 46 | 2,137 |
| 19. | Seer fish | 956 | 151 | 294 | 307 | 233 | 300 | 2,241 |
| 20. | Tunnies | 15 | 11 | 16 | 145 | 216 | 43 | 446 |
| 21. | <i>Sphyraena</i> | 246 | 158 | 52 | 201 | 87 | 9 | 753 |
| 22. | <i>Mugil</i> | 13 | 22 | 28 | 5 | 25 | 16 | 109 |
| 23. | <i>Breg-</i> <i>maceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | 130 | 218 | 152 | 220 | 173 | 205 | 1,098 |
| 25. a) | Penaeid prawns | 819 | 793 | 680 | 1,492 | 605 | 1,834 | 6,223 |
| b) | Non- penaeid prawns | 33 | 16 | 2 | 6 | 38 | 88 | 183 |
| c) | Lobsters | 47 | 32 | 38 | 3 | 9 | 14 | 143 |
| d) | Other crust- | 252 | 262 | 294 | 527 | 319 | 549 | 2,203 |
| 26. | Cepha- | 92 | 109 | 90 | 134 | 55 | 81 | 561 |
| 27. | lopods | 1,644 | 1,782 | 2,300 | 812 | 1,111 | 1,413 | 9,062 |
| TOTAL | | 19,274 | 19,041 | 23,899 | 16,348 | 14,529 | 19,153 | 1,12,244 |

Table 7. Composition of marine fish landings in Pondicherry during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|--------------|---------------------------------|-------|------|-------|-------|-----|-------|-------|
| 1. | Elasmobranchs | 2 | 20 | 5 | 25 | 11 | 34 | 97 |
| 2. | Eels | — | 20 | — | 46 | — | 4 | 70 |
| 3. | Cat fishes | — | 3 | — | 11 | — | 13 | 27 |
| 4. | <i>Chirocentrus</i> | 2 | 8 | 8 | 6 | 9 | 62 | 95 |
| 5. a) | Oil sardine | — | — | — | — | — | — | — |
| b) | Other sardines | 119 | 114 | 17 | 24 | 107 | 134 | 515 |
| c) | <i>Hilsa ilisha</i> | — | — | — | — | — | — | — |
| d) | Other <i>Hilsa</i> | 1 | 1 | 2 | — | — | — | 4 |
| e) | <i>Ancho-viella</i> | — | 32 | 10 | 30 | 2 | 7 | 81 |
| f) | <i>Thris-socies</i> | 7 | 26 | 47 | 24 | 43 | 65 | 212 |
| g) | Other clupeoids | 5 | 24 | 20 | 17 | 55 | 18 | 139 |
| 6. a) | <i>Harpodon nehereus</i> | — | — | — | — | — | — | — |
| b) | <i>Saurida & Saurus</i> | 3 | 2 | 1 | 35 | 3 | 56 | 100 |
| 7. | <i>Hemiramphus & Belone</i> | — | — | — | — | — | — | — |
| 8. | Flying fish | — | — | — | — | 495 | 495 | — |
| 9. | Perches | 6 | 54 | 14 | 98 | 27 | 134 | 333 |
| 10. | Red mullets | 1 | 1 | 3 | 48 | 15 | 26 | 94 |
| 11. | Polynemids | — | — | — | — | — | — | — |
| 12. | Sciaenids | 7 | 19 | 11 | 14 | 9 | 29 | 89 |
| 13. | Ribbon fish | 15 | 5 | 8 | 8 | — | 11 | 47 |
| 14. a) | <i>Caranx</i> | 2 | 6 | 5 | 19 | 30 | 3 | 65 |
| b) | <i>Chorinemus</i> | — | — | 1 | — | — | 2 | 3 |
| c) | <i>Trachy-notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | — | — | — | — | — | — | — |
| e) | <i>Coryphaena</i> | — | — | — | 2 | — | — | 2 |
| f) | <i>Elacate</i> | — | — | — | — | — | — | — |
| 15. a) | <i>Leiognathus</i> | 12 | 49 | 7 | 48 | 31 | 89 | 236 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | — | 1 | — | — | 3 | 1 | 5 |
| 17. | Pomfrets | 2 | — | — | 10 | — | — | 12 |
| 18. | Mackerel | 4 | 134 | 25 | 42 | 22 | — | 227 |
| 19. | Seer fish | 3 | 1 | — | 2 | 8 | 39 | 53 |
| 20. | Tunnies | — | — | — | — | — | 1 | 1 |
| 21. | <i>Sphyraena</i> | — | — | 1 | 1 | 7 | — | 9 |
| 22. | <i>Mugil</i> | — | 4 | — | 9 | — | 5 | 18 |
| 23. | <i>Breg-maceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | 2 | 8 | 5 | 14 | 22 | 27 | 78 |
| 25. a) | Penaeid prawns | 7 | 51 | 6 | 60 | 37 | 59 | 220 |
| b) | Non-penaeid prawns | — | — | — | 27 | — | 29 | 56 |
| c) | Other crustaceans | 4 | 1 | 5 | 3 | 31 | 28 | 72 |
| 26. | Cephalopods | — | — | 1 | 15 | — | 4 | 20 |
| 27. | Miscellaneous | 10 | 37 | 12 | 72 | 37 | 47 | 215 |
| TOTAL | | 214 | 621 | 214 | 710 | 509 | 1,422 | 3,690 |

Table 8. Composition of marine fish landings in Kerala during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|--------------|---------------------------------|--------|--------|--------|--------|--------|-------|---------|
| 1. | Elamo-branches | 633 | 402 | 542 | 255 | 847 | 175 | 2,854 |
| 2. | Eels | 1 | — | — | 2 | — | — | 3 |
| 3. | Cat fishes | 210 | 114 | 194 | 1,332 | 1,766 | 468 | 4,084 |
| 4. | <i>Chirocentrus</i> | 22 | 115 | 30 | 381 | 102 | — | 650 |
| 5. a) | Oil sardine | 10,341 | 9,530 | 7,221 | 2,168 | 6,737 | 1,679 | 37,676 |
| b) | Other sardines | 188 | 774 | 322 | 1,314 | 1,410 | 9 | 4,017 |
| c) | <i>Hilsa ilisha</i> | — | — | — | — | — | — | — |
| d) | Other <i>Hilsa</i> | — | 1 | — | 15 | 18 | — | 34 |
| e) | <i>Ancho-viella</i> | 34 | 338 | 1,776 | 551 | 410 | 568 | 3,677 |
| f) | <i>Thris-socies</i> | 89 | 12 | 41 | 214 | 45 | 33 | 434 |
| g) | Other clupeoids | 16 | 31 | 5 | 202 | 103 | 11 | 368 |
| 6. a) | <i>Harpodon nehereus</i> | — | — | 1 | — | — | — | 1 |
| b) | <i>Saurida & Saurus</i> | 51 | 48 | 13 | 142 | 686 | 184 | 1,124 |
| 7. | <i>Hemiramphus & Belone</i> | 11 | 107 | 7 | 16 | 3 | — | 144 |
| 8. | Flying fish | — | — | — | — | — | — | — |
| 9. | Perches | 1,477 | 171 | 153 | 279 | 227 | 25 | 2,332 |
| 10. | Red mullets | — | 21 | 105 | — | — | — | 126 |
| 11. | Polynemids | 1 | — | 24 | — | 1 | — | 26 |
| 12. | Sciaenids | 632 | 361 | 433 | 494 | 638 | 271 | 2,829 |
| 13. | Ribbon fish | 35 | 32 | 19 | 189 | 101 | 2,247 | 2,623 |
| 14. a) | <i>Caranx</i> | 447 | 572 | 200 | 527 | 461 | 83 | 2,290 |
| b) | <i>Chorinemus</i> | 3 | 21 | 4 | 1 | 15 | — | 44 |
| c) | <i>Trachy-notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | — | — | — | — | — | — | — |
| e) | <i>Coryphaena</i> | — | 10 | 31 | 5 | 2 | — | 48 |
| f) | <i>Elacate</i> | — | 1 | 16 | 1 | 5 | — | 23 |
| 15. a) | <i>Leiognathus</i> | 231 | 449 | 146 | 745 | 422 | 63 | 2,056 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | 1 | — | 1 | 1 | 61 | 79 | 143 |
| 17. | Pomfrets | 95 | 156 | 255 | 45 | 216 | 7 | 774 |
| 18. | Mackerel | 1,472 | 2,361 | 3,017 | 4,902 | 4,759 | 192 | 16,703 |
| 19. | Seer fish | 542 | 398 | 95 | 62 | 95 | 17 | 1,209 |
| 20. | Tunnies | 322 | 208 | 537 | 2,527 | 7,374 | 441 | 11,409 |
| 21. | <i>Sphyraena</i> | 67 | 30 | 38 | 60 | 74 | 7 | 276 |
| 22. | <i>Mugil</i> | 1 | 1 | 28 | — | — | — | 30 |
| 23. | <i>Breg-maceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | 398 | 184 | 493 | 353 | 258 | 81 | 1,767 |
| 25. a) | Penaeid prawns | 1,274 | 971 | 4,580 | 1,665 | 2,741 | 1,437 | 12,668 |
| b) | Non-penaeid prawns | — | 9 | 7 | 8 | — | — | 24 |
| c) | Lobsters | 7 | 2 | 1 | — | 1 | — | 11 |
| d) | Other crustaceans | 974 | 565 | 558 | 783 | 239 | 79 | 3,198 |
| 26. | Cephalopods | 68 | 41 | 31 | 36 | 81 | 29 | 286 |
| 27. | Miscellaneous | 725 | 655 | 1,920 | 1,632 | 1,383 | 222 | 6,537 |
| TOTAL | | 20,368 | 18,691 | 22,844 | 20,907 | 31,281 | 8,407 | 122,498 |

Table 9. Composition of marine fish landings in Karnataka during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|---------|----------------------------------|--------|-------|--------|-------|-------|-------|--------|
| 1. | Elasmo-brancls | 108 | 59 | 244 | 72 | 62 | 26 | 571 |
| 2. | Eels | — | — | — | — | — | — | — |
| 3. | Cat fishes | 296 | 137 | 313 | 502 | 60 | 6 | 1,314 |
| 4. | <i>Chirocentrus</i> | 6 | 8 | 4 | 4 | — | 31 | 53 |
| 5. a) | Oil sardine | 598 | 5,065 | 1,247 | 1,357 | 953 | 436 | 9,656 |
| b) | Other sardines | 293 | 276 | 599 | 821 | 1,546 | 22 | 3,557 |
| c) | <i>Hilsa ilisha</i> | 2 | — | — | — | — | — | 2 |
| d) | Other <i>Hilsa</i> | 1 | 1 | 24 | — | — | — | 26 |
| e) | <i>Ancho- viella</i> | 443 | 45 | 285 | 7 | 71 | — | 851 |
| f) | <i>Thris-socles</i> | 73 | 42 | 41 | 28 | 18 | 8 | 210 |
| g) | Other clupeids | 385 | 119 | 140 | 53 | 29 | 14 | 740 |
| 6. a) | <i>Harpodon nehereus</i> | — | 1 | — | — | — | — | 1 |
| b) | <i>Saurida & Saurus</i> | 24 | 18 | 31 | 71 | 12 | — | 156 |
| 7. | <i>Hemirhamphus & Belone</i> | — | — | 9 | — | — | — | 9 |
| 8. | Flying fish | — | — | — | — | — | — | — |
| 9. | Perches | 1 | 7 | 25 | 46 | 23 | 1 | 103 |
| 10. | Red mullets | — | — | 14 | 10 | 5 | — | 29 |
| 11. | Polynemids | 2 | — | — | — | — | — | 2 |
| 12. | Sciaenids | 195 | 182 | 401 | 64 | 28 | 170 | 1,040 |
| 13. | Ribbon fish | 51 | 50 | 61 | 43 | 225 | 2 | 432 |
| 14. a) | <i>Caranx</i> | 113 | 47 | 12 | 57 | 43 | 12 | 284 |
| b) | <i>Chorinemus</i> | 1 | 4 | — | — | 1 | — | 6 |
| c) | <i>Trachy-notus</i> | — | 13 | — | — | — | — | 13 |
| d) | Other carangids | — | — | — | — | — | — | — |
| e) | <i>Coryphaena</i> | — | — | — | — | — | — | — |
| f) | <i>Elacate</i> | 4 | 1 | 3 | — | — | 1 | 9 |
| 15. a) | <i>Letognathus</i> | 110 | 118 | 132 | 173 | 120 | 19 | 672 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | 45 | 16 | 28 | 8 | 13 | 30 | 140 |
| 17. | Pomfrets | 48 | 20 | 23 | 5 | 1 | 1 | 98 |
| 18. | Mackerel | 3,699 | 990 | 2,253 | 1,535 | 568 | 1,721 | 10,766 |
| 19. | Seer fish | 132 | 170 | 92 | 7 | 3 | — | 404 |
| 20. | Tunnies | — | — | 1 | 21 | 626 | 28 | 676 |
| 21. | <i>Sphyraena</i> | 8 | 2 | 18 | 2 | — | — | 30 |
| 22. | <i>Mugil</i> | — | — | — | — | 1 | — | 1 |
| 23. | <i>Bregmaceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | 35 | 26 | 58 | 78 | 18 | — | 215 |
| 25. a) | Penaeid prawns | 942 | 882 | 886 | 787 | 651 | 60 | 4,208 |
| b) | Non-penaeid prawns | — | 2 | — | — | — | — | 2 |
| c) | Lobsters | — | — | 15 | — | — | — | 15 |
| d) | Other crustaceans | 747 | 350 | 1,185 | 162 | 39 | — | 2,483 |
| 26. | Cephalopods | 32 | 5 | 11 | 2 | 2 | — | 52 |
| 27. | Miscellaneous | 2,084 | 758 | 2,455 | 2,339 | 1,872 | 154 | 9,662 |
| | TOTAL | 10,478 | 9,414 | 10,610 | 8,254 | 6,990 | 2,742 | 48,488 |

Table 10. Composition of marine fish landings in Goa during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|---------|----------------------------------|-------|-------|-------|-------|-----|------|-------|
| 1. | Elasmo-brancls | 138 | 77 | 133 | 144 | 42 | — | 534 |
| 2. | Eels | — | — | — | — | 4 | — | 4 |
| 3. | Cat fishes | 72 | 54 | 141 | 158 | 13 | 3 | 441 |
| 4. | <i>Chirocentrus</i> | 25 | 5 | 7 | 21 | — | — | 58 |
| 5. a) | Oil sardine | 36 | 79 | 417 | 113 | 13 | — | 658 |
| b) | Other sardines | 145 | 92 | 186 | 62 | — | — | 485 |
| c) | <i>Hilsa ilisha</i> | — | — | — | — | — | — | — |
| d) | Other <i>Hilsa</i> | — | — | — | 14 | — | — | 14 |
| e) | <i>Ancho- viella</i> | — | — | — | — | — | — | — |
| f) | <i>Thris-socles</i> | 475 | 104 | 120 | 58 | 58 | — | 815 |
| g) | Other clupeids | 120 | 25 | 109 | 9 | 26 | — | 289 |
| 6. a) | <i>Harpodon nehereus</i> | — | — | — | — | — | — | — |
| b) | <i>Saurida & Saurus</i> | 1 | 5 | 88 | 2 | — | — | 96 |
| 7. | <i>Hemirhamphus & Belone</i> | 2 | — | — | 9 | — | — | 11 |
| 8. | Flying fish | 1 | — | — | — | — | — | 1 |
| 9. | Perches | 24 | 25 | 128 | 2 | 2 | — | 181 |
| 10. | Red mullets | — | — | — | — | — | — | — |
| 11. | Poly-nemids | — | — | — | — | — | — | — |
| 12. | Sciaenids | 254 | 133 | 190 | 133 | 49 | 6 | 765 |
| 13. | Ribbon fish | 27 | 37 | 176 | 145 | 69 | — | 454 |
| 14. a) | <i>Caranx</i> | 295 | 116 | 132 | 40 | 54 | 3 | 640 |
| b) | <i>Chorinemus</i> | 3 | — | 33 | 9 | 4 | — | 49 |
| c) | <i>Trachy-notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | — | — | — | — | — | — | — |
| e) | <i>Coryphaena</i> | — | — | — | — | — | — | — |
| f) | <i>Elacate</i> | — | 1 | — | — | — | — | 1 |
| 15. a) | <i>Letognathus</i> | 127 | 101 | 195 | 94 | 65 | 1 | 583 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | 35 | 22 | 23 | 55 | 11 | — | 146 |
| 17. | Pomfrets | 29 | 16 | 12 | 14 | 3 | — | 74 |
| 18. | Mackerel | 137 | 29 | 39 | — | — | — | 205 |
| 19. | Seer fish | 173 | 52 | 64 | 6 | — | — | 295 |
| 20. | Tunnies | 3 | 2 | — | — | — | — | 5 |
| 21. | <i>Sphyraena</i> | — | — | 7 | — | — | — | 7 |
| 22. | <i>Mugil</i> | 1 | 1 | — | — | — | 3 | 5 |
| 23. | <i>Bregmaceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | 53 | 28 | 35 | 76 | 43 | — | 235 |
| 25. a) | Penaeid prawns | 128 | 99 | 357 | 518 | 166 | 8 | 1,276 |
| b) | Non-penaeid prawns | 55 | — | — | — | — | — | 55 |
| c) | Lobsters | — | — | — | — | — | — | — |
| d) | Other crustaceans | 18 | 114 | 478 | 142 | 69 | 3 | 824 |
| 26. | Cephalopods | 37 | 21 | 13 | 23 | 12 | — | 106 |
| 27. | Miscellaneous | 74 | 59 | 103 | 64 | 14 | 10 | 324 |
| | TOTAL | 2,488 | 1,297 | 3,186 | 1,911 | 717 | 37 | 9,636 |

Table 11. Composition of marine fish landings in Maharashtra during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|--------------|----------------------------------|--------|--------|--------|--------|--------|-------|---------|
| 1. | Elasmo-branches | 1,576 | 510 | 847 | 609 | 1,338 | 201 | 5,081 |
| 2. | Eels | 677 | 407 | 493 | 565 | 41 | 8 | 2,191 |
| 3. | Cat fishes | 923 | 935 | 990 | 1,177 | 510 | 99 | 4,634 |
| 4. | <i>Chirocentrus</i> | 68 | 148 | 70 | 23 | 169 | — | 478 |
| 5. a) | Oil sardine | — | — | — | — | — | — | — |
| b) | Other sardines | 87 | 6 | 12 | 7 | 74 | — | 186 |
| c) | <i>Hilsa ilisha</i> | 49 | 170 | 38 | 9 | 79 | — | 345 |
| d) | Other <i>Hilsa</i> | 11 | 15 | 49 | 14 | 16 | 4 | 109 |
| e) | <i>Ancho- viella</i> | — | 9 | — | 1 | 22 | 10 | 42 |
| f) | <i>Thri- soles</i> | 207 | 230 | 339 | 39 | 297 | 3 | 1,115 |
| g) | Other clupeids | 1,117 | 1,488 | 988 | 1,301 | 918 | 162 | 5,974 |
| 6. a) | <i>Harpodon nehereus</i> | 3,768 | 1,920 | 1,461 | 2,832 | 4,063 | 289 | 14,333 |
| b) | <i>Saurida & Saurus</i> | 607 | 383 | 348 | 114 | 62 | 1 | 1,515 |
| 7. | <i>Hemirhamphus & Belone</i> | 9 | — | — | 1 | — | 1 | 11 |
| 8. | Flying fish | — | — | — | — | — | — | — |
| 9. | Perches | 278 | 300 | 258 | 122 | 249 | 85 | 1,292 |
| 10. | Red mullets | 117 | 132 | 281 | 21 | 3 | 1 | 555 |
| 11. | Poly- nemids | 178 | 144 | 242 | 207 | 53 | 12 | 836 |
| 12. | Sciaenids | 2,735 | 1,803 | 2,069 | 1,849 | 1,546 | 136 | 10,138 |
| 13. | Ribbon fish | 1,527 | 1,191 | 869 | 874 | 924 | 143 | 5,528 |
| 14. a) | <i>Caranx</i> | 466 | 122 | 268 | 61 | 258 | 30 | 1,205 |
| b) | <i>Chori- nemus</i> | 9 | 2 | — | 52 | 28 | — | 91 |
| c) | <i>Trachy- notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | — | 27 | — | 4 | 1 | 9 | 41 |
| e) | <i>Cory- phaena</i> | — | — | — | — | 9 | 2 | 11 |
| f) | <i>Elaeate</i> | — | — | — | — | — | — | — |
| 15. a) | <i>Letog- nathus</i> | 97 | 12 | 18 | 52 | 107 | 23 | 309 |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | 58 | 229 | 2 | 9 | 1 | — | 299 |
| 17. | Pomfrets | 573 | 903 | 583 | 1,573 | 671 | 78 | 4,381 |
| 18. | Mackerel | 556 | 88 | 6 | 21 | 12 | — | 683 |
| 19. | Seer fish | 93 | 43 | 74 | 114 | 114 | 8 | 446 |
| 20. | Tunnies | 139 | 28 | 10 | 43 | 94 | 100 | 414 |
| 21. | <i>Sphyraena</i> | 25 | — | 3 | — | 2 | — | 30 |
| 22. | <i>Mugil</i> | — | — | — | 1 | 8 | 10 | 19 |
| 23. | <i>Breg- maceros</i> | — | — | — | — | — | — | — |
| 24. | Soles | 11 | 20 | 220 | 109 | 413 | 14 | 787 |
| 25. a) | Penaeid prawns | 2,947 | 2,651 | 4,283 | 4,948 | 3,421 | 491 | 18,741 |
| b) | Non-penaeid prawns | 4,623 | 8,358 | 3,503 | 7,797 | 6,204 | 518 | 31,003 |
| c) | Lobsters | 58 | 274 | 46 | 65 | 36 | 17 | 496 |
| d) | Other crustaceans | — | 1 | 13 | 39 | 20 | 13 | 86 |
| 26. | Cephalopods | 314 | 235 | 271 | 387 | 285 | 66 | 1,558 |
| 27. | Miscellaneous | 765 | 470 | 476 | 386 | 560 | 55 | 2,712 |
| TOTAL | | 24,668 | 23,254 | 19,130 | 25,426 | 22,609 | 2,589 | 117,676 |

Table 12. Composition of marine fish landings in Gujarat during the half year ending June 1979 (in tonnes)

| Sl. No. | Name of fish | Janu. | Feb. | March | April | May | June | Total |
|--------------|----------------------------------|--------|-------|--------|--------|-------|-------|--------|
| 1. | Elasmo-branches | 534 | 293 | 308 | 517 | 521 | 234 | 2,407 |
| 2. | Eels | 1,347 | 88 | 72 | 210 | 2 | — | 1,719 |
| 3. | Cat fishes | 257 | 158 | 1,671 | 632 | 139 | 69 | 2,926 |
| 4. | <i>Chirocentrus</i> | 332 | 238 | 324 | 128 | 53 | 16 | 1,091 |
| 5. a) | Oil sardine | — | — | — | — | — | — | — |
| b) | Other sardines | — | — | — | — | — | — | — |
| c) | <i>Hilsa ilisha</i> | — | 161 | 12 | 5 | 2 | 6 | 186 |
| d) | Other <i>Hilsa</i> | 210 | 333 | 462 | 269 | 121 | 45 | 1,440 |
| e) | <i>Ancho- viella</i> | — | — | — | — | — | — | — |
| f) | <i>Thri- soles</i> | 87 | 81 | 33 | 86 | 22 | 7 | 316 |
| g) | Other clupeids | 1,039 | 29 | 484 | 324 | 101 | 38 | 2,015 |
| 6. a) | <i>Harpodon nehereus</i> | 6,564 | 474 | 837 | 2,159 | 668 | 60 | 10,762 |
| b) | <i>Saurida & Saurus</i> | — | 5 | — | — | — | — | 5 |
| 7. | <i>Hemirhamphus & Belone</i> | — | — | — | — | — | — | — |
| 8. | Flying fish | — | — | — | — | — | — | — |
| 9. | Perches | 461 | 11 | 169 | 40 | 4 | 5 | 690 |
| 10. | Red mullets | — | — | — | — | — | — | — |
| 11. | Poly- nemids | — | — | — | — | — | — | — |
| 12. | Sciaenids | 4,821 | 2,284 | 4,329 | 1,868 | 172 | 49 | 13,523 |
| 13. | Ribbon fish | 649 | 344 | 272 | 344 | 439 | — | 2,048 |
| 14. a) | <i>Caranx</i> | 9 | 44 | 2 | 5 | 4 | 15 | 79 |
| b) | <i>Chori- nemus</i> | 48 | 9 | 3 | 18 | 1 | 3 | 82 |
| c) | <i>Trachy- notus</i> | — | — | — | — | — | — | — |
| d) | Other carangids | — | — | — | — | — | — | — |
| e) | <i>Cory- phaena</i> | — | — | — | — | — | — | — |
| f) | <i>Elaeate</i> | — | — | — | — | — | — | — |
| 15. a) | <i>Letog- nathus</i> | — | — | — | — | — | — | — |
| b) | <i>Gazza</i> | — | — | — | — | — | — | — |
| 16. | <i>Lactarius</i> | 161 | 37 | 11 | — | — | — | 209 |
| 17. | Pomfrets | 474 | 133 | 369 | 383 | 925 | 613 | 2,897 |
| 18. | Mackerel | — | — | — | 6 | 35 | — | 41 |
| 19. | Seer fish | 557 | 372 | 485 | 42 | 12 | 7 | 1,475 |
| 20. | Tunnies | 41 | 31 | 25 | 1 | — | — | 98 |
| 21. | <i>Sphyraena</i> | — | — | — | — | — | — | — |
| 22. | <i>Mugil</i> | 52 | 29 | 55 | 23 | 18 | 48 | 225 |
| 23. | <i>Breg- maceros</i> | 85 | 34 | 135 | — | — | — | 254 |
| 24. | Soles | 21 | 10 | 70 | 2 | — | 20 | 123 |
| 25. a) | Penaeid prawns | 3,256 | 149 | 470 | 392 | 97 | 30 | 4,394 |
| b) | Non-penaeid prawns | 65 | 33 | 414 | 206 | 76 | 5 | 799 |
| c) | Lobsters | 43 | 2 | 43 | 20 | 5 | — | 113 |
| d) | Other crustaceans | — | 75 | 359 | 48 | 10 | 8 | 500 |
| 26. | Cephalopods | 194 | 544 | 2,924 | 827 | 29 | — | 4,518 |
| 27. | Miscellaneous | 6,709 | 3,256 | 4,186 | 4,318 | 408 | 141 | 19,018 |
| TOTAL | | 28,016 | 9,260 | 18,525 | 12,896 | 3,961 | 1,431 | 74,089 |

Goa

The total landings in Goa showed an increase of about 4,000 tonnes (Table 1). The landings of elasmobranchs, penaeid prawns, crabs and other crustaceans, *Leiognathus* spp, sciaenids, *Caranx*, ribbon fish, oil sardine and other clupeids increased substantially. The catch of other sardines, *Saurida* & *Saurus*, perches and *Lactarius*, however, was poor. Table 10 gives the details of catch. While the maximum landings were recorded in the month of March, the minimum were observed in June.

Maharashtra

The total catch in Maharashtra increased by about 15,600 tonnes as compared to the corresponding half year ending June 1978 (Table 1). While the landings of penaeid prawns, non penaeid prawns, sciaenids, *Harpodon nehereus*, ribbon fish, *Caranx*, pomfrets,

mackerel, other clupeids and elasmobranchs, showed substantial increase, the catch of cat fishes, perches, eels, *Chirocentrus*, *Hilsa ilisha*, cephalopods and soles was comparatively poor. The catch details are shown in Table 11. The maximum catch was noticed in April 1979 and the minimum in June 1979.

Gujarat

A decline of about 19,000 tonnes in the total landings was noticed in Gujarat in comparison to the first half of 1978 (Table 1). The catch of *Harpodon nehereus*, penaeid prawns and cephalopods showed significant increase. But the landings of many varieties of fishes such as elasmobranchs, sciaenids, perches, ribbon fish, *Lactarius*, pomfrets, other *Hilsa* and other clupeids were significantly lower (Table 12). The maximum and minimum catch was recorded in the months of January and June respectively.

