


Cephalopholis igarashiensis, Garish Hind

Assessment by: Rhodes, K., Samoilys, M., Choat, J.H., Myers, R., To, A., Ma, K., Nair, R., Suharti, S., Law, C., Amorim, P. & Russell, B.



View on www.iucnredlist.org

Citation: Rhodes, K., Samoilys, M., Choat, J.H., Myers, R., To, A., Ma, K., Nair, R., Suharti, S., Law, C., Amorim, P. & Russell, B. 2018. *Cephalopholis igarashiensis*. The IUCN Red List of Threatened Species 2018: e.T132791A100455099. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T132791A100455099.en>

Copyright: © 2018 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Actinopterygii	Perciformes	Epinephelidae

Taxon Name: *Cephalopholis igarashiensis* Katayama, 1957

Synonym(s):

- *Cephalopholis igarashiensis* Katayama, 1957
- *Cephalopholis igarasiensis* Katayama, 1957
- *Cephalopholis swanius* Tsai, 1960

Common Name(s):

- English: Garish Hind, Goldbar Grouper, Japanese Cod
- French: Vielle Voyant
- Spanish: Cherna Chillona

Assessment Information

Red List Category & Criteria: Least Concern [ver 3.1](#)

Year Published: 2018

Date Assessed: November 17, 2016

Justification:

Cephalopholis igarashiensis is a widely distributed, naturally rare and deep-living species. There are no known major threats; therefore, it is listed as Least Concern. The change in status from the previous assessment reflects an improved application of the Red List categories and criteria, as well as a better understanding of available data.

Previously Published Red List Assessments

2008 – Data Deficient (DD)

<http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T132791A3452612.en>

Geographic Range

Range Description:

This species is distributed from southern Japan to Fiji and French Polynesia. Its depth range is 64-250 metres.

The specific localities in which it occurs are as follows: Japan, Ogasawara Islands (Japan), Taiwan, Philippines, Guam, Tahiti, and French Polynesia (Heemstra and Randall 1993). It is also reported in Fiji and Tuvalu by Chapman and Cussack (1990). Palau, Marianas, Marshall Islands, Federated States of Micronesia, Rotuma (Fiji), Samoa, Cook Islands, Society Islands and Tuamotus Islands (French Polynesia) and Christmas Island. Also recorded from Scott Reef and Rowley Shoals in waters off northwestern Australia (Rome and Newman 2010). Its occurrence in Tahiti, French Polynesia is reported from a

specimen caught at 250 m.

Country Occurrence:

Native: American Samoa; Australia; Christmas Island; Cook Islands; Disputed Territory (Paracel Is., Spratly Is.); Fiji; French Polynesia; Indonesia; Japan; Marshall Islands; Micronesia, Federated States of ; New Caledonia; Niue; Northern Mariana Islands; Palau; Philippines; Samoa; Taiwan, Province of China; Timor-Leste

FAO Marine Fishing Areas:

Native: Indian Ocean - eastern, Pacific - eastern central, Pacific - western central, Pacific - northwest

Distribution Map

Cephalopholis igarashiensis




Sources: Esri, HERE, Garmin, Intermap, increment P Corp., GEBCO, USGS, FAO, NPS, NRCAN, GeoBase, IGN, Kadaster NL, Ordnance Survey, Esri Japan, METI, Esri China (Hong Kong), swisstopo, © OpenStreetMap contributors, and the GIS User Community

Range

Extant (resident)

Compiled by:

IUCN Grouper and Wrasse Specialist Group



The boundaries and names shown and the designations used on this map do not imply any official endorsement, acceptance or opinion by IUCN.



Population

This deep-living species is presumed to be naturally rare to scarce. In Guam, catch in 1985 was 10 kg and 260 kg in 1999 (Zeller 2006). It is exploited commercially in Indonesia, but catch data are not available (P. Mous and J. Pet unpublished data 2017). A length-based assessment of commercial deepwater grouper fisheries in eastern Indonesia reported 131 individuals, 5% of which were immature (≥ 17 cm to ≤ 22 cm total length). The spawning potential ratio was estimated at near 100% and the stock is currently considered not overfished (P. Mous and J. Pet unpublished data 2017).

Current Population Trend: Unknown

Habitat and Ecology (see Appendix for additional information)

This deep-living, demersal species is associated with steep reef drop-offs, seamounts and offshore banks (Mathieu *et al.* 2002). The maximum total length is 44.8 cm. The upper depth limit was originally reported as 80 m, but there is a photo available of this species at a depth of 64 m (reef.com). At lengths between 14 to 19 cm, this species is two to five years of age (Fry *et al.* 2006). Natural mortality is estimated as 0.35, longevity is 40.6 years and length at maturity is about 23-25.4 cm (P. Mous and J. Pet unpublished data 2017).

Systems: Marine

Use and Trade

Juveniles of this species are highly-valued in the aquarium fish trade, and retails at USD\$6,000 per individual (reef.com). It is caught incidentally and retained in deep-water subsistence and commercial fisheries, including in Guam, Indonesia and Papua New Guinea (Zeller 2006, Fry *et al.* 2006, P. Mous and J. Pet unpublished data 2017). Rare occurrences in markets in some parts of its range indicate that it is not likely being targeted.

Threats (see Appendix for additional information)

There are no known major threats.

Conservation Actions

In Indonesia, there is a total length limit (23.9 cm) for take of this species, and this is considered sufficient to maintain a sustainable fishery (P. Mous and J. Pet unpublished data 2017).

Credits

Assessor(s): Rhodes, K., Samoily, M., Choat, J.H., Myers, R., To, A., Ma, K., Nair, R., Suharti, S., Law, C., Amorim, P. & Russell, B.

Reviewer(s): Linardich, C.

Facilitators(s) and Compiler(s): Carpenter, K.E.

Bibliography

Chapman, L.B. and Cusack, P. 1990. South Pacific Commission Deep Sea Fisheries Development Project Report on Second Visit to Tuvalu 30 August - 7 December 1983. In: South Pacific Commission. (ed.). SPC (Noumea), Noumea, New Caledonia.

Fry, G.C., Brewer, D.T. and Venables, W.N. 2006. Vulnerability of demersal fishes to commercial fishing: Evidence from a study around a tropical seamount in Papua New Guinea. *Fisheries Research* 81: 126-141.

Heemstra, P.C. and Randall, J.E. 1993. *FAO species catalogue. Vol. 16. Groupers of the world (Family Serranidae, Subfamily Epinephelinae). An annotated and illustrated catalogue of the grouper, rockcod, hind, coral grouper and lyretail species known to date.* FAO, Rome.

IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-2. Available at: www.iucnredlist.org. (Accessed: 15 November 2018).

Mathieu, L. F., Allison, E.H. and Tinch, R. 2002. Seeking patterns of population variability from fish catch and stock biomass time series. University of East Anglia, Overseas Development Group, Norwich (UK).

Zeller, D. 2006. Unpublished data base; Creel & Offshore surveys American Samoa, Guam and the Northern Mariana Islands. Available at: <http://www.seararoundus.org>.

Citation

Rhodes, K., Samoily, M., Choat, J.H., Myers, R., To, A., Ma, K., Nair, R., Suharti, S., Law, C., Amorim, P. & Russell, B. 2018. *Cephalopholis igarashiensis*. The IUCN Red List of Threatened Species 2018: e.T132791A100455099. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T132791A100455099.en>

Disclaimer

To make use of this information, please check the [Terms of Use](#).

External Resources

For [Images and External Links to Additional Information](#), please see the [Red List website](#).

Appendix

Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
9. Marine Neritic -> 9.2. Marine Neritic - Subtidal Rock and Rocky Reefs	-	Suitable	-
9. Marine Neritic -> 9.8. Marine Neritic - Coral Reef -> 9.8.3. Foreslope (Outer Reef Slope)	-	Suitable	-
11. Marine Deep Benthic -> 11.1. Marine Deep Benthic - Continental Slope/Bathyl Zone (200-4,000m) -> 11.1.1. Hard Substrate	-	Marginal	-
11. Marine Deep Benthic -> 11.5. Marine Deep Benthic - Seamount	-	Suitable	-

Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
5. Biological resource use -> 5.4. Fishing & harvesting aquatic resources -> 5.4.1. Intentional use: (subsistence/small scale) [harvest]	Ongoing	-	-	-
	Stresses:	2. Species Stresses -> 2.1. Species mortality		

Research Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Research Needed
1. Research -> 1.2. Population size, distribution & trends
1. Research -> 1.3. Life history & ecology
3. Monitoring -> 3.1. Population trends

Additional Data Fields

Distribution
Lower depth limit (m): 250
Upper depth limit (m): 64
Population
Population severely fragmented: No

The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).