

Note

Biological aspects of the Indo-Pacific sergeant *Abudefduf vaigiensis* (Quoy and Gaimard, 1825) from Gulf of Mannar, Tamil Nadu, India

N. Rajesh^{1*}, Imelda Joseph¹, N. K. Chadha², Bobby Ignatius¹, P. P. Suresh Babu¹, K. K. Anikuttan¹, A. Anuraj¹, Paramita Banerjee Sawant² and A. M. Babitha Rani²

¹ICAR-Central Marine Fisheries Research Institute, Kochi - 682 018, Kerala, India

²ICAR-Central Institute of Fisheries Education, Mumbai - 400 061, Maharashtra, India



Abstract

Food and feeding habits of the commercially important marine ornamental fish *Abudefduf vaigiensis* from the Gulf of Mannar were evaluated. Analysis of gut contents indicated that *A. vaigiensis* is an omnivorous fish, feeding on any available food item in its environment, such as seaweeds, cladocerans, copepods and insects. Most of the sampled fishes had either quarter-full or half-full or three-fourth full stomachs. The index of preponderance indicated seaweeds as the preferred food item among majority of length classes except for those falling within the 80-89 mm and 100-109 mm length classes, in which a preference towards copepods was observed. On a monthly basis, seaweed *Spongomorpha* sp. emerged as the predominant food item. Gastro-somatic index (GSI) for males recorded the highest value in July (1.20) and lowest (0.36) in November, while that of females recorded the highest value in July (0.95) and lowest value (0.47) in January. Hepato-somatic index (HSI) of females recorded the highest in December (2.01) and lowest in July (0.83), while that of males recorded the highest value in January (1.25) and lowest in September (0.81). Gonadosomatic index (GSI) in females recorded maximum in February (4.23) and lowest in June (1.32) while that of males recorded the highest value in February (0.85) and lowest value in July (0.35). The results of the study indicate that adopting seasonal regulatory measures is essential for sustainable exploitation of the species, since their trade relies on wild collection. The study also revealed a negative correlation between the feeding patterns of these fish and their reproductive development.



*Correspondence e-mail:
rajeshnellora@yahoo.co.in

Keywords:

Abudefduf vaigiensis, Feeding intensity, Gastro-somatic index, Gonadosomatic index, Hepato-somatic index

Received : 20.06.2022

Accepted : 02.04.2024

Fishes belonging to the family Pomacentridae are banded marine fishes widely distributed along the Indo-west and Central Pacific tropical and warm temperate waters. They are primarily found in shallow coral reefs, with few species inhabiting depths of 80 m or more (Khalaf and Disi, 1997). The Indo-Pacific sergeant major, *Abudefduf vaigiensis* (Quoy and Gaimard, 1825) is one of the important groups of marine ornamental fishes of this family, which contribute significantly to the total marine ornamental fish trade in India. It feeds mainly on macroalgae, benthic invertebrates and detritus (Sudhakar and Shameen, 2006). The study of food and feeding habits of fishes is essential to understand the species' life

history, including its growth, breeding and migration (Bal and Rao, 1984). Information on the feeding habits of marine fish and their predatory relationship is helpful in assessing their role in the ecosystem (Bachok *et al.*, 2004). Knowledge of the influence of maturity stages, age and season on food and feeding preferences is important in assessing the trophic inter-relationship. Even though several studies on the diet composition in marine ornamental fishes are available (Mohan and Pillai, 1988; Pillai and Mohan, 1990; Pillai and Vijay Anand, 1995; Gandhi, 2002; Pillai *et al.*, 2002), specific studies on the seargent major group are limited. The present study analysed the food spectrum and dietary

preferences of *A. vaigiensis* from Gulf of Mannar, south-east coast of India. It is also the first attempt to investigate these details in relation to gonadal development in *A. vaigiensis* and the results of this study will be of use in planning commercial aquaculture of the species.

Monthly samples of thirty individuals of *A. vaigiensis* were collected from October 2019 to September 2020 from Gulf of Mannar, south-east coast of India (Fig. 1). A total of 360 intact samples of *A. vaigiensis* caught by encircling nets were used for the study. The total length and standard length were measured to the nearest 0.1 cm and weight measurements (total body weight and gut weight) to the nearest 0.1 g using a measuring board and a sensitive weighing balance respectively. *A. vaigiensis* was identified following the description given by Allen (1991) and Randall (2007).

Each specimen was weighed to the nearest 1.0 g with the help of a single pan balance (Sartorius 'PT' 600) and then dissected to collect gut contents for analysis. Gut contents were preserved in 4% formalin solution and kept for further analysis. Qualitative analysis of gut contents was done in the Mariculture Laboratory of Mandapam Regional Centre of ICAR-Central Marine Fisheries Research Institute (ICAR-CMFRI), Mandapam, Tamil Nadu, India with the help of a trinocular microscope (Olympus CH 20i) following the frequency of occurrence methods as per Hynes (1950) and Pillay (1952).

The data were segregated month-wise, sex-wise and size-wise. The gut contents were grouped into different categories such as zooplankton, phytoplankton, plant material, insects and decaying matter. The relative importance of all food contents was quantified by the index of preponderance and was calculated with the help of percentage composition (volume and occurrence) of food contents to follow the equation of Natrajan and Jhingran (1963):

$$I = (V1O1 / \sum V1 O1) 100$$

where, I = Index of preponderance; V1= Percentage volume of particular food component; O1 = Occurrence of particular food component

The degree of fullness of stomach and the quantity of food contained in it were noted so as to ascertain the extent of feeding (feeding intensity). The degree of distension of stomachs was designated as gorged (50 points), full (40 points), $\frac{3}{4}$ full (30 points), $\frac{1}{2}$ full (20 points), $\frac{1}{4}$ full (10 points) and empty (0 points) following the method by Hynes (1950). A stomach was considered 'gorged' when it was packed with food and stretched fully with thin, transparent walls, 'full' when filled with food and the walls thick and intact and ' $\frac{3}{4}$ full' when the stomach was partly collapsed with thick walls. Depending on the amount of food present and the appearance of the stomach wall, the stomach was further designated as ' $\frac{1}{2}$ full', ' $\frac{1}{4}$ full' and 'empty'.

The samples collected were adequately cleaned in the laboratory, dissected and the stomachs were removed. The total weight of each stomach with its contents was measured to the nearest 0.01 g. The contents of each stomach and foregut were examined under a microscope and further identification of food content within each taxonomic group was made following appropriate taxonomic identification guides. GaSI based on monthly and seasonal calculations was obtained as described by Biswas (1993):

$$\text{GaSI} = (\text{Total weight of stomach} / \text{Bodyweight}) \times 100$$

Liver was removed and weight was recorded to the nearest 0.001 g. HSI was calculated following Sulistyo *et al.* (2000):

$$\text{HSI} = \frac{\text{Liver weight (g)}}{\text{Body weight (g)}} \times 100$$

The fishes were cut open and the morphology of the gonad was observed for sex determination. Gonads were then removed, weighed and macroscopically assessed to determine their maturation stage. Gonad weight was recorded to the nearest 0.001 g. Body weight and gonad weight were measured using a top-loading electronic balance with an accuracy of 0.001 g and the length was measured using a wooden scale with an accuracy of 1 mm.




Fig.1. Location of sample collection for feeding studies of *A. vaigiensis*

To quantify the changes in gonad weight during the annual sexual cycle and to determine the spawning season, monthly GSI (Devlaming *et al.*, 1982), was calculated using the formula:

$$\text{GSI} = \frac{\text{Gonad weight (g)}}{\text{Body weight (g)}} \times 100$$

Monthly analysis of the stomach contents of *A. vaigiensis* revealed that the diet consisted of seaweeds, polychaetes, copepods, shrimps, insect larvae, decapod larvae, mollusc, sagitta and organic debris. Index of preponderance of *A. vaigiensis* for various length classes and during various months is given in Table 1 and 2 respectively. Preferred food items for various length classes were: Copepods (80-89 mm), *Spongomorpha* sp. (90-99 mm), Copepods (100-109 mm), *Gracilaria* sp. (110-119 mm), *Spongomorpha* sp. (120-129 mm), *Chaetomorpha* sp. (130-139 mm), *Spongomorpha* sp. (140-149 mm) and *Chaetomorpha* sp. (150-159 mm).

On an average, for all months of the study period, the seaweed *Spongomorpha* sp. dominated with a percentage of 29% and the other food items in descending order were seaweed *Chaetomorpha* sp. (19.54%), copepods (17.96%), *Halobates* sp. (15.94%), *Gracilaria* sp. (7.92%), *Kappaphycus* sp. (3.33%), *Ulva* sp. (2.88%), *Caulerpa* sp. (1.03%), molluscs (0.63%) and shrimps (0.63%). The preference for food items were in the order: - *Spongomorpha* sp. (I), *Chaetomorpha* sp. (II), Copepods (III), *Halobates* sp. (IV), *Gracilaria* sp. (V), *Kappaphycus* sp. (VI), *Ulva* sp. (VII), *Caulerpa* sp. (VIII), molluscs (IX) and shrimps (X). The results of the present study indicated that *A. vaigiensis* is an omnivorous species feeding on any available food item in its environments such as seaweeds, cladocerans, copepods and insects.

Degree of fullness (%) of stomachs of *A. vaigiensis* observed in different months is given in Table 3. Out of the 360 specimens evaluated, the percentage of empty stomach was minimum (1.1 to 5.5%), the full stomach was comparatively less (2.2 to 56.3%) and

Table 1. Index of preponderance of *A. vaigiensis* for various length classes (mm)

Organism	80-89	90-99	100-109	110-119	120-129	130-139	140-149	150-159
<i>Chaetomorpha</i> sp.	0.2	29.5 (II)	31.5 (II)	22.3 (II)	25.2 (II)	35.2 (I)	18 (II)	26.3 (I)
<i>Spongomorpha</i> sp.	1.9 (II)	45.3 (I)	15.2 (III)	5.4	28.3 (I)	31.2 (II)	52 (I)	24.3 (II)
<i>Ulva</i> sp.	0.9	3.6	9.6	4.05	8.6	1.2	0.2	3.5
<i>Caulerpa racemosa</i>	0.12	0.5	1.1	1.38	1.2	1.5	4.5	5.2
<i>Kappaphycus</i> sp.	0.9	1.6	1.2	15.3(III)	5.2	5.6	9.15	12.5
<i>Gracilaria</i> sp.	0.95(III)	2.82	0.51	30.25 (I)	8.2	15.2 (III)	12.1 (III)	19.3 (III)
Polychaetes	0	0	0.15	1.3	0.25	0	0	0
Copepods	90.93 (I)	15.2 (III)	35 (I)	12.3	12.5 (III)	0.1	2.6	5.3
Shrimps	0	0.53	0.2	0.5	0	8.94	1.2	0
<i>Halobates</i> sp.	0	0	1.5	4.5	4.86	0	0	0
Molluscs	0	0.2	0.04	0	0	0.5	0	0
<i>Sagitta</i> sp.	0.52	0	0.8	0.5	0	0	0	0
Decapod larvae	0.26	0.25	0	0.02	0.02	0	0	0
Miscellaneous	3.32	0.5	3.2	2.2	5.67	0.56	0.25	3.6
Total	100	100	100	100	100	100	100	100

*Most preferred food item for each length class given in parenthesis

Table 2. Month-wise Index of preponderance of *A. vaigiensis*

Organism	Oct	Nov	Dec	Jan	Feb	Mar	Apl	May	Jun	Jul	Aug	Sep
<i>Chaetomorpha</i> sp.	18.4 (II)	22.2 (III)	2.1	2.5	13.3 (II)	2.6	53.4 (I)	7.8 (II)	38.5 (II)	34.8 (II)	6.2 (III)	32.66 (II)
<i>Spongomorpha</i> sp.	1.0	3.4	0.6	1.2	65 (I)	54.2 (I)	28.5 (II)	70.2 (I)	48.5 (I)	51.2 (I)	23 (II)	1.2
<i>Ulva</i> sp.	1.8	0.6	0.3	0.24	0.32	2.6	1.5	4.4	3.2	3.1	1.1	15.4 (III)
<i>Caulerpa racemosa</i>	0.2	0.5	1	3.26 (III)	1.58	0	1.3	0.6	0.1	0.25	2.4	1.2
<i>Kappaphycus</i> sp.	2.1	1.2	2.9 (III)	0.5	5.2	9.4 (III)	7.1	5.6 (III)	0.8	0.1	2.5	2.5
<i>Gracilaria</i> sp.	3.6 (III)	36.2 (I)	1.2	15.2 (II)	8.5 (III)	5.5	7.6 (III)	4.9	8.5 (III)	0.9	2.1	0.8
Polychaetes	0	0	0	0.2	0.1	0	0	0	0	0	0	0
Copepods	70.1 (I)	2.5	40.2 (II)	0	0	0	0	0	0	9.1 (III)	55.5 (I)	38.1 (I)
Shrimps	0	1.2	1.05	1.5	0.6	0	0	2.5	0	0	0	0.44
<i>Halobates</i> sp.	0	32.2(II)	50.4 (I)	75.2(I)	4.7	25.1(II)	0	1.9	0	0	0.3	1.5
Molluscs	0	0	0	0	0.2		0.5	0	0.4	0	5.5	0.9
<i>Sagitta</i> sp.	0	0	0	0.2	0	0	0.1	0	0	0	0.1	0.5
Decapod larvae	0.2	0	0	0	0	0.3	0	2.1	0	0	0.1	2.1
Miscellaneous	2.6	0	0.25	0	0.5	0.3	0	0	0	0.55	1.2	2.7
Total	100	100	100	100	100	100	100	100	100	100	100	100

*Most preferred food item for each length class given in parenthesis

Table 3. Degree of fullness (%) of stomachs of *A. vaigiensis* in different months

2019-20	N	Empty	¼ full	½ full	¾ full	Full	Gorged
Oct	30	1.1	27.5	39.3	31.2	0	0.8
Nov	30	0	20.2	59.4	20.1	0	0.2
Dec	30	1.2	38.9	30.3	20.1	9.4	0
Jan	30	0	25.7	52.7	11.7	9.8	0
Feb	30	2.1	30.6	36.2	31	0	0
Mar	30	0	38.5	30.2	23.1	8.1	0
Apl	30	0	32.5	28.1	39.2	0	0
May	30	5.1	13	30.8	42.5	8.5	0
Jun	30	0	12.3	18.1	12.1	56.3	1.1
Jul	30	3.1	27.5	8.8	40.54	18.1	2.05
Aug	30	8	0	37.7	50.2	2.2	1.8
Sep	30	5.5	16.4	35.4	37.8	3.8	1





most of the fishes had either 1/4 or 1/2 or 3/4-full stomachs. The highest percentage of empty stomachs was in September and the maximum percentage (56.3%) of full stomachs was in June.

GaSI (Fig. 2.) in male *A. vaigiensis* followed regular pattern with an increasing trend after February and reached the highest value (1.20) in July and lowest value (0.36) in November. GaSI in female *A. vaigiensis* followed an irregular pattern with an increasing trend followed by a decrease, with the highest value in July (0.95) and the lowest (0.47) in January.

HSI (Fig. 3.) in male *A. vaigiensis* increased from November (0.85) to January (1.25) and decreased from February to September, with a peak in December. HSI in females increased from August (0.85) to December (1.25) and then decreased up to July with a peak in December.

GSI in male *A. vaigiensis* (Fig. 4) sustained a maximum from September to March with a peak during February and was observed to be lower from April to July. GSI in female *A. vaigiensis* (Fig. 5) sustained maximum values from December to March with a peak during February and was observed to be lower from April to October.

Diet is a key factor that profoundly affects an organism's distribution, growth, reproduction, migration rate and behaviour (Priyadharsini et al., 2012). Feed quality and quantity directly impact fish growth and indirectly impact fish maturity and survival (Sourinejad et al., 2015). Information on the food and feeding habit of a species in its natural habitat provides cues for the selection of suitable cultivable

Fig. 2. Monthly gastro-somatic index in male and female *A. vaigiensis*Fig. 3. Monthly hepato-somatic index in male and female *A. vaigiensis*Fig. 4. Monthly gonado-somatic index in males of *A. vaigiensis*Fig. 5. Monthly gonado-somatic index in females of *A. vaigiensis*

species and for the development of successful farming practices (Manon and Hossain, 2011).

Research on the feeding habits and feeding ecology of fishes is the basis for understanding their roles in ecosystems (Hajisamae *et al.*, 2003). In addition, the knowledge of the feeding behaviour of marine fishes is essential to understand prey selection (Stergiou and Fourtouni, 1991), predator-prey size relationships (Mohanraj and Prabhu, 2012; Priyadharsini *et al.*, 2012), distribution of feeding habits with latitude (Pauly, 2000), and habitat selection (Labropoulou *et al.*, 1999), which contribute to fish stock assessment and ecosystem modelling (Bachok *et al.* 2004; Salavatian *et al.*, 2011; Mohanraj and Prabhu, 2012).

In general, damselfishes living in coral reefs of the Indo-West Pacific region feed mainly on filamentous algae and small planktonic and benthic invertebrates (Allen, 1991; Kuo and Shao, 1991; Frédérich *et al.*, 2009). The results of the present study were on par with other previous reports for the same species (Fishelson, 1970; Emery, 1973; Allen, 1991). *Abudefduf troschelii*, the pacific sergeant-major, was found to be omnivorous, foraging on benthic organisms (Hobson, 1965). Analysis of gut contents in the present study indicated that *A. vaigiensis* is an omnivorous species which agrees with the observations made on *A. vaigiensis* in the Mediterranean Sea (Allen, 1991). Fishelson (1970) noted that *A. vaigiensis* feeds on floating algae, whereas Emery (1973) reported that it feeds on benthic algae. Fishes often display environmentally induced morphological variations (Wimberger, 1992), reflecting different feeding biology (Sreekanth *et al.*, 2012).

The feeding intensity studies of *A. vaigiensis* samples collected from the Gulf of Mannar region revealed that most of the fishes had either 1/4th or 1/2th or 3/4th of the stomach. The highest percentage of empty stomachs was in September, which correlated with the GSI peak observed in this study. Percentage of empty stomachs and feeding intensity in fishes have been observed to be negatively related (Bowman and Bowman, 1980; Pallaoro *et al.*, 2003) and often synchronised with the spawning seasons of the respective fishes (Manon and Hossain, 2011; Salavatian *et al.*, 2011; Sourinejad *et al.*, 2015).

GaSI of male and female fishes followed an almost similar pattern and recorded low value during the period of high GSI in the present study. Perera and Cumaranatunga (2009) reported that GaSI peaked just before the breeding season in *A. vaigiensis* and reduced during reproductive activity as GSI increased. HSI followed a similar trend in both sexes and correlated with the trends of GSI in the present study. Higher liver activities are always associated with yolk development for vitellogenesis and gonadal development. HSI is used as an indicator of energy reserves in the liver (Cerdeira *et al.*, 1996; Hismayasari *et al.*, 2015) and as an indicator of the reproductive period in fishes (Bolger and Connolly 1989; Jayasankar and Alagarwami, 1994; Barbieri *et al.*, 1996; Carvalho *et al.*, 2009; De Giosa *et al.*, 2014).

The results of GSI in the present study revealed that April to August (pre-breeding season) is the preparatory season for the breeding season and the fishes feed voraciously during this phase leading to higher GaSI. During the breeding season (September to March), the gonadal development is rigorous, leading to higher GSI. GSI has been used to improve accuracy in determining the maturity stage (McPherson *et al.*, 2011) and has also been widely used

to determine reproduction timing (Lowerre-Barbieri *et al.*, 2011). During the high GSI, the alimentary canal will be thinner and the peritoneal space will be occupied by the developing gonad. Keivany and Soofiani (2004) reported that the feeding intensity declines as the fish approaches the peak spawning season. The present study gives the first report on the food preferences and feeding indices of *A. vaigiensis* in the Gulf of Mannar and also give a clue to the breeding season of *A. vaigiensis* in this region, which can be used for regulatory measures on their capture in the wild and breeding of these species in controlled conditions.

Acknowledgements

The authors are thankful to Dr A. Gopalakrishnan, Director, ICAR-CMFRI, Kochi for providing necessary facilities to carry out this work. We also acknowledge the support and logistics rendered by Mandapam Regional Centre of ICAR-CMFRI during the study period. Authors are thankful to ICAR, Govt. of India, New Delhi, for financial support.

References

- Allen, G. R. 1991. *Damselfishes of the world*. Hans A. Baensch, Mergus Publishers, Melle, Germany, 271 p.
- Bachok, Z., Mansor, M. I. and Noordin, R. M. 2004. Diet composition and food habits of demersal and pelagic marine fishes from Terengganu waters, east coast of Peninsular Malaysia. *NAGA, WorldFish Center Quarterly*, 27(3&4): 41-47.
- Bal, D. V. and Rao, K. V. 1984. *Marine fisheries of India*. Tata McGraw Hill Publishing Co. Pvt. Ltd., New Delhi, India, 296 p.
- Barbieri, G., Hartz, S. M. and Verani, R. J. 1996. The condition factor and gonadosomatic index as indicator of the spawning period for *Astyanax fasciatus* (Cuvier, 1819) in the Lobo Reservoir, São Paulo (Osteichthyes, Characidae). *Iheringia, Serie Zoologia*, 81: 97-100.
- Biswas, S. P. 1993. *Manual of methods in fish biology*. South Asian Publishers Pvt. Ltd., New Delhi, India, 157 p.
- Bolger, T. and Connolly, P. L. 1989. The selection of suitable indices for the measurement and analysis of fish condition. *J. Fish Biol.*, 34: 171-182. <https://www.researchgate.net/publication/227832970>.
- Bowman, R. E. and Bowman, E. W. 1980. Diurnal variation in the feeding intensity and catchability of silver hake (*Merluccius bilinearis*). *Can. J. Fish. Aquat. Sci.*, 37: 1565-1572.
- Carvalho, P. A., Paschoalini A. L., Santos, G. B., Rizzo, E. and Bazzoli, N. 2009. Reproductive biology of *Astyanax fasciatus* (Pisces: Characiformes) in a reservoir in southeastern Brazil. *J. Appl. Ichthyol.*, 25: 306-313. <https://doi.org/10.1111/j.1439-0426.2009.01238.x>.
- Cerdeira, J., Calman, B. G., LaFleur, Jr, G. J. and Limesand, S. 1996. Pattern of vitellogenesis and follicle maturational competence during the ovarian follicular cycle of *Fundulus heteroclitus*. *Gen. Comp. Endocrinol.*, 103(1): 24-35. <https://doi.org/10.1006/gcen.1996.0090>.
- De Giosa, M., Czerniejewski, P. and Rybczyk, A. 2014. Seasonal changes in condition factor and weight length relationship of invasive *Carassius gibelio* (Bloch, 1782) from Leszczynskie Lakeland, Poland. *Adv. Zool.*, 2014, Article ID 678763 <http://dx.doi.org/10.1155/2014/678763>.
- Devlaming, V., Grossman, G. and Chapman, F. 1982. On the use of the gonosomatic index. *Comp. Biochem. Physiol. Part A: Physiol.*, 73(1): 31-39. [https://doi.org/10.1016/0300-9629\(82\)90088-3](https://doi.org/10.1016/0300-9629(82)90088-3).

- Emery, A. R. 1973. Comparative ecology and functional osteology of fourteen species of the damselfish (Pisces: Pomacentridae) at Alligator Reef, Florida Keys. *Bull. Mar. Sci.*, 23(3): 649-770.
- Fishelson, L. 1970. Behaviour and ecology of a population of *Abudefduf saxatilis* (Pomacentridae, Teleostei) at Eilat (Red Sea). *Anim. Behav.*, 18: 225-237.
- Frédérich, B., Fabri, G., Lepoint, G., Vandewalle, P. and Parmentier, E. 2009. Trophic niches of thirteen damselfishes (Pomacentridae) at the Grand Récif of Toliara, Madagascar. *Ichthyol. Res.*, 56(1): 10-17. <https://doi.org/10.1007/s10228-008-0053-2>.
- Gandhi, V. 2002. Studies on the food and feeding habits of cultivable butterfish *Scatophaps arps* (Cuv. and Val.). *J. Mar. Biol. Ass. India*, 44(1&2): 115-121.
- Hajisamae, S., Chou, L. M. and Ibrahim, S. 2003. Feeding habits and trophic organization of the fish community in shallow waters of an impacted tropical habitat. *Estuar. Coast. Shelf Sci.*, 58(1): 89-98. [https://doi.org/10.1016/S0272-7714\(03\)00062-3](https://doi.org/10.1016/S0272-7714(03)00062-3).
- Hismayasari, I. B., Marhendra, A. P. W., Rahayu, S., Saidin, S. D. and Supriyadi, D. S. 2015. Gonadosomatic index (GSI), hepatosomatic index (HSI) and proportion of oocytes stadia as an indicator of rainbowfish *Melanotaenia boesemani* spawning season. *Int. J. Fish. Aquat. Stud.*, 2(5): 359-362.
- Hobson, E. S. 1965. Diurnal-nocturnal activity of some inshore fishes in the Gulf of California. *Copeia*, 1965(3): 291-302.
- Hynes, H. B. N. 1950. The food of freshwater stacle backs *Gasterosteus aculeatus* and *Pygosteus pungitus*, with a review of methods used in studies of the food of fishes. *J. Anim. Ecol.*, 19: 36-58. <https://doi.org/10.2307/1570>.
- Jayasankar, P. and Alagaswami, K. 1994. Histological and histochemical observations on the oocytes in the sand whiting *Sillago sihama* (Forsskal). *Proceedings of the Indian National Science Academy, B. Biological Sciences*, 60: 173-182.
- Keivany, Y. and Soofiani, N. 2004. Contrbtion to the biology of Zagros tooth-carp, *Aphanius vladkovi* (Cyprinodontidae) in central Iran. *Environ. Biol. Fishes*, 71(2):165-169. <https://doi.org/10.1007/s10641-004-0106-y>.
- Khalaf, M. A. and Disi, A. M., 1997. *Fishes of the Gulf of Aqaba*. Marine Science Station, Aqaba, Jordan, 252 p.
- Kuo, S. R. and Shao K. T. 1991. Feeding habits of damselfish (Pomacentridae) from the southern part of Taiwan. *J. Fish. Soc. Taiwan*, 18: 165-176.
- Labropoulou, M. and Papadopolou Smith, K. N. 1999. Foraging behavior patterns of four sympatric demersal fishes. *Estuar. Coast. Shelf Sci.*, 49: 99-108. [https://doi.org/10.1016/S0272-7714\(99\)80014-6](https://doi.org/10.1016/S0272-7714(99)80014-6).
- Lowerre-Barbieri, S. K., Ganas, K., Saborido-Rey, F., Murua, H. and Hunter, J. R. 2011. Reproductive timing in marine fishes: Variability, temporal scales, and methods. *Mar. Coast. Fish.*, 3(1): 71-91. <https://doi.org/10.1080/19425120.2011.556932>.
- Manon, M. R. and Hossain, M. D. 2011. Food and feeding habit of *Cyprinus carpio* var. *specularis*. *J. Sci. Found.*, 9(1&2): 163-181. <https://doi.org/10.3329/jsf.v9i1-2.14658>.
- McPherson, L. R., Ganas, K. and Marshall, C. T. 2011. Inaccuracies in routinely collected Atlantic herring (*Clupea harengus*) maturity data and correction using a gonadosomatic index model. *J. Mar. Biol. Assoc. UK*, 91(7): 1477-1487.
- Mohan, M. and Pillai, C. S. G. 1988. The biology of the convict surgeonfish *Acanthurus triostegus triostegus* (Linnaeus) from Minicoy Atoll, Lakshadweep. *J. Mar. Biol. Assoc. India.*, 30(1&2): 182-191.
- Mohanraj, T. and Prabhu, K. 2012. Food habits and diet composition of demersal marine fishes from Gulf of Mannar, Southeast coast of India. *Adv. Biol. Res.*, 6(4): 159-164. <https://doi.org/10.5829/idosi.abr.2012.6.4.6486>.
- Natarajan, A. V. and Jhingran, A. G. 1963. On the biology of *Catla catla* (Ham.) from the river Jamuna. *Proceedings of the National Institute of Sciences of India*, 29B: 326-355.
- Pallaoro, A., Santic, M. and Jardas, I. 2003. Feeding habits of the Saddled bream, *Oblada melanura* (Sparidae), in the Adriatic Sea. *Cybiurn*, 27(4): 261-268.
- Pauly, D. 2000. Herbivory as a low-latitude phenomenon. In: Froese, R. and Pauly, D. (Eds.), *FishBase 2000: Concepts, design and data sources*. International Centre for Living Aquatic Resources Management, Laguna, Philippines, 179 p.
- Perera, W. A. F. C. and Cumaranatunga, P. R. T. 2009. Feeding ecology, reproductive biology and aspects related to selected marine ornamental fish, *Abudefduf vaigiensis*. *Proceedings of the Annual Academic Sessions of University of Ruhuna*, 6: 50-57.
- Pillay, T. V. R. 1952. A critique of the methods of study of food of fishes. *J. Zool. Soc. India*, 4: 185-200.
- Pillai, C. S. G. and Mohan, M. 1990. Ecology and biology of *Abudefduf glaucus* (Cuvier) (Pomacentridae, Pisces) from Minicoy Atoll, Lakshadweep. *Indian J. Fish.*, 37(1): 15-23.
- Pillai, N. G. K. and Vijay Anand, P. E. 1995. Studies on some aspects of biology and ecology of coral reef fishes of Lakshadweep with observations on other coral reef ecosystems of India. *CMFRI Special Publication No. 61*. ICAR-Central Marine Fisheries Research Institute, Kochi, India, pp. 99-111.
- Pillai, P. P., Gopakumar, G. and Said Koya, K. P. 2002. Spatial distribution pattern and biology of tuna live-bait fishes in Lakshadweep. *J. Mar. Biol. Ass. India*, 44(1&2): 231-236.
- Priyadharsini, S., Manoharan, J., Varadharajan, D. and Subramaniyan, A. 2012. Interpretation on the food and feeding habits of *Dascyllus trimaculatus* (Ruppell, 1829) from Gulf of Mannar, South East coast of India. *Arch. Appl. Sci. Res.*, 4(4): 1758-1762.
- Randall, J. E. 2007. *Reef and shore fishes of the Hawaiian Islands*. University of Hawaii Press Sea Grant College Program, Honolulu, Hawaii, 560 p.
- Salavatian, M., Gholiev, Z., Aliev, A. and Abassi, K. 2011. Feeding behavior of brown trout, *Salmo trutta fario*, during spawning season in four rivers of Lar National Park, Iran. *Caspian J. Environ. Sci.*, 9: 223-233.
- Sreekanth, B., Chakraborty, S. K., Jaiswar, A. K., Nair, J. R., Pazhayamadom, D. G., Renjith, R. K., Ratheesh K., Sandeep, K. P. and Anuraj, A. 2012. Site specific differences in food and feeding biology of *Nemipterus japonicus* (Bloch, 1791) along Indian coast. *Indian J. Fish.*, 59(3): 25-31.
- Sourinejad, I., Nikkhah Khaje Ataei, S., Kamrani, E. and Ghodrati Shojaei, M. 2015. Feeding habits of yellowfin seabream, *Acanthopagrus latus* in the northern region of the Persian Gulf. *Caspian J. Environ. Sci.*, 13(1): 31-39.
- Stergiou, K. I. and Fourtouni, H. 1991. Food habits, ontogenetic diet shift and selectivity in *Zeus faber* Linnaeus, 1758. *J. Fish Biol.*, 39(4): 589-603. <https://doi.org/10.1111/j.1095-8649.1991.tb04389.x>.
- Sudhakar, G. V. S. and Shameem, A. 2006. Food and feeding habits of Sergeant major, *Abudefduf vaigiensis* (Quoy and Gaimard, 1825) from shallow waters off Visakhapatnam. *J. Mar. Biol. Ass. India*, 48(1): 12-123.
- Sulistyio, I., Fontaine, P., Rincarh, J., Gardeur J. N., Migaud, H. and Capdeville, B. 2000. Reproductive cycle and plasma level of steroid in male eurasian perch (*Perca fluviatilis*). *Aquat. Living Resour.*, 13(2): 99-106. [https://doi.org/10.1016/S0990-7440\(00\)00146-7](https://doi.org/10.1016/S0990-7440(00)00146-7).
- Wimberger, P. H. 1992. Plasticity of fish body shape: The effects of diet, development, family and age in two species of *Geophagus* (Pisces: Cichlidae). *Biol. J. Linn. Soc.*, 45: 197-218. <https://doi.org/10.1111/J.1095-8312.1992.TB00640.X>.