### J. mar. biol. Ass. India, 1998, 40 (1 & 2) : 105 - 124

# FOOD AND FEEDING HABITS OF THE SPOTTED SEER, SCOMBEROMORUS GUTTATUS (BLOCH AND SCHNEIDER), IN THE GULF OF MANNAR AND PALK BAY

#### M. DEVARAJ

## Central Marine Fisheries Research Institute, Cochin

#### ABSTRACT

The spotted seer, essentially a surface feeder, feeds on a limited number of about five species, of which the sardines are the most important and the whitebaits the next. The ration per active feeding (R) in 1968-69 declined by 50% of the 1967-68 level, owing to shortage of food. The shift from whitebaits which constitute the exclusive diet of the young, to sardines which form the major food of the adult, takes place at about 315 mm, about 1/4 the asymptotic length. The fish is far less aggressive than the kingseer or the streaked seer as evident from its inability to compete ably in times of food shortage, the much larger size at which it shifts to feeding on larger forage species and the orientation of 68.4% of the forage fish in its stomach in its reverse axis. The spotted seer is numerically more abundant, but gravimetrically less abundant than the kingseer, and therefore, its poor catch by weight must be seen more as a function of its small size than as a function of competence or competitiveness.

There are two active feedings a day, one between 7 and 10 p.m. and the other between 6 and 10 a.m. Food intake did not slacken in maturing or ripe fish and was generally high from September to March. Young fish less than 300 mm length, distributed in nearshore areas were found much better fed than the adults in grounds beyond the 20 meter depth line. Young spotted seer (say, 1 to 3 year old) consume more food per unit body weight than old fish (say, 4 year old), but the former require much less food to produce a unit weight growth than the latter. The T-line suggests that the ration available and consumed by the spotted seer is sufficient enough to sustain the normal growth of the fish, and that the process of normal food consumption and growth goes on in spite of severe interspecific competition. The K-lines reveal that the general level of gross growth efficiency is within the normal range of 0.25 to 0.75. The Paloheimo-Dickie weight growth model is not valid for the spotted seer, owing to the slope of the T-line being significantly different from the value describing the relation between metabolism and body weight under normal non-stress conditions.

#### INTRODUCTION

STUDIES concerning the food of the spotted seer of the seas around India are limited to those by Venkataraman (1960) for the juveniles from Calicut, Kumaran (1964), of the postlarvae and juveniles from Vizhinjam and Srinivasa Rao (1964) of the juveniles and adults from Waltair. The results of a systematic study on the food and feeding habits of the spotted seer scomberomorus guttatus from the northern Gulf of Mannar and Palk Bay along the southeast coast of India, undertaken during 1967-'69, are dealt with in this account.

I owe my sincere thanks to Dr. S. Jones, Dr. R.V. Nair, Dr. E.G. Silas and Dr. P.S.B.R. James of the Central Marine Fisheries Research Institute, Cochin and Dr. S.N. Dwivedi of the Central Institute of Fisheries Education, Bombay for their guidance and encouragement. I am also thankful to Mr. M. Stephen, fleet owner, for his help in the procurement of most of the samples for this study.

#### MATERIAL AND METHODS

710 spotted seer comprising 65 from Palk Bay (zone I) and 645 from the northern Gulf of Mannar (zone II) sampled from the commerical drift gillnet fishery in the Rameswaram Island twice or thrice a week between July, 1967 and July, 1969, formed the basic material for the study. The method of examination of the stomach contents and the treatment of the data for the determination of food composition, qualitative and quantitative variations of food according to zones and seasons, according to the size and maturity of the fish, variation in average ration and ration per unit body weight, aggressiveness in feeding, and food intake and utilisation are the same as followed in a similar study of the king seer (Devaraj, MS).

#### RESULTS

#### Food composition

The food of the spotted seer was represented by fishes belonging to three families, one cephalopod and one penaeid (Table 1).

### (i) Analysis by volume

710 spotted seer from both zones I (Palk Bay) and zone II (northern Gulf of Mannar) were found to have consumed 1,651.1 ml food comprising 99.6% fish (teleosts), 0.3% Loligo and 0.01% prawn. Sardinella contributed the greatest volume (86.8%) of all food elements. Anchoviella was second in importance (6.3%). Carangids formed a meagre 0.3%. Unidentified fishes ('fishes') accounted for 2.1%. Fish scales formed 0.6% and digested matter, 3.6% (Table 1).

#### (ii) Analysis by frequency of occurrence

99.6% of the total frequencies (487) were contributed by fish to which were also added the frequencies of digested matter on the presumption that they were of fish origin. Squids and prawns contributed 0.2% each. Sardinella alone accounted for nearly half (49.3%) with an index of preponderance 60 and Anchoviella 11.3% with an IP 35.3. 29.4% of the stomachs contained only digested matter. Other items occurred less frequently; carangids with an IP of 0.0002 and fishes with an IP of 0.34. Prawns and Loligo recorded a meagre 0.0001% and 0.0002% respectively by IP (Table 1)

### Zonal variations

The food dispensation is similar in both the zones. The volume of 282 ml food taken from 65 stomachs sampled from zone I was composed exclusively of fish (*Sardinella* 236 ml, *Anchoviella* 37.5 ml and 'fishes' 7.8 ml) including 2.5 ml digested matter assumed to be of fish origin. By volume (V), occurrence (O), and *IP*, *Sardinella*, *Anchoviella* and 'fishes' ranked in descending order.

Besides the organisms noted above, negligible quantities of carangids, squids and

 

 TABLE 1. Food composition by volume (V), frequency occurrence (O) and index of preponderance (IP) for a total of 710 spotted seer sampled from Palk Bay (zone I) and the northern Gulf of Mannar (zone II) during 1967-'69

| b flest owner.  | Vo      | lume | and the second of | Frequency of              | ccurrence | i india | Index of | preponderance         |
|-----------------|---------|------|-------------------|---------------------------|-----------|---------|----------|-----------------------|
| Food items      | ml      | %    | rank              | Actual                    | %         | rank    | %        | rank                  |
| Sardinella      | 1432.1  | 86.8 | spiel for         | 240                       | 49.3      | 1.00    | 60       | from Calicut, N       |
| Anchoviella     | 102.7   | 6.3  | 2                 | 55                        | 11.3      | 3       | 35.3     | 2                     |
| Carangids       | man Ska | 0.3  | 6                 | 1 or whe                  | 0.2       | 6       | 0.002    | 6                     |
| 'Fishes'        | 35      | 2.1  | 4                 | 34                        | 7         | 4       | 0.34     | t bas4bool asi        |
| Fish scales     | 10.5    | 0.6  | 5                 | 12                        | 2.4       | 5       | 0.22     | mom5 mdmona           |
| Prawns          | 0.2     | 0.01 | 8                 | uthessi Bay<br>69. arc of | 0.2       | nol6 y  | 0.001    | of Main <b>e</b> r an |
| Loligo          | 5       | 0.3  | 6                 | 100 i                     | 0.2       | 6       | 0.002    | ni ibiw theat         |
| Digested matter | 60.6    | 3.67 | 3                 | 143                       | 29.4      | 2       | 4.18     | 3                     |
| Total           | 1651.1  | 100  |                   | 487                       | 100       | e asis  | 100      | aield V X ad          |

prawns also formed the diet in zone II. In the first year (August, 1967 to July, 1968), the food consisted of 99.8% fish and 0.2% prawns. Sardinella formed 91.3% by V, 60.7% by O and 98.3 by IP (Table 3). The other six items

food in all months in 1967-'68 with *IP* varying from 75 to 100 except in July and September (*IP* = 20 to 29) during which *Anchoviella* dominated the diet (*IP* = 52 to 61). During the other months, *Anchoviella* was very low

 TABLE 2. Index of preponderance (IP) of organisms in the diet according to months for zone I (Palk Bay) during 1967-68

| Months               | Aug.<br>(1967) | Sep. | Oct. | Dec. | Jan.<br>(1968) | Jun. | Jul. | Aug. | Sep. | Oct. | Total | Rank |
|----------------------|----------------|------|------|------|----------------|------|------|------|------|------|-------|------|
| No. of fish examined | 5              | 5    | 15   | 5    | 6              | 4    | 8    | 10   | 6    | 5 1  | 65    |      |
| Sardinella           | 89.6           | 93.8 | 98   | 100  | 99             | 0    | 0    | 53.3 | 52.2 | 0    | 92.2  | 1    |
| Anchoviella          | 10.4           | 6.2  | 2    | 0    | 0              | 100  | 37.2 | 30.1 | 43.5 | 0    | 7.2   | 2    |
| 'Fishes'             | 0              | 0    | 0    | 0    | 1              | 0    | 0    | 15.4 | 0    | 0    | 0.4   | 3    |
| Digested matter      | 0              | 0    | 0    | 0    | 0              | 0    | 62.8 | 1.2  | 4.3  | 100  | 0.2   | 4    |
| Total                | 100            | 100  | 100  | 100  | 100            | 100  | 100  | 100  | 100  | 100  | 100   | 8    |

had the levels of *IP* ranging from a mere 0.002 (prawns) to 0.912 (digested matter) (Table 3a).

In the second year (August, 1968 to May, 1969), 332.8 ml food from 240 stomachs consisted mainly of fish items and only 5 ml of squid remains. *Sardinella* was the most dominant item of food by IP = 80.2, but O for *Sardinella* declined to 31% as majority of stomachs contained only digested matter (Table 3b).

#### Seasonal variations

In 1967-'68, in Palk Bay (zone I), Sardinella formed a major item of food from August through January with IP (90 to 100), during which Anchoviella was much less significant IP = 0 to 10. In June and July, Sardinella was absent in the diet, but Anchoviella dominated with IP ranging from 37 to 100. In the two succeeding months, August and September, Sardinella attained a marginal increase over that of Anchoviella (Table 2).

In northern Gulf of Mannar (zone II), Sardinella was the most significant item of in the food with an IP = 4 in August and absent during October to January and April. Carangids and prawns were present only in July and September (Table 3a).

In 1968-'69, Sardinella was absent in the diet in January, ranked low in December and May and high in all other months (IP = 81 to 92). In those months in which Sardinella was either absent or the least in the diet, either digested matter alone was present as in December and in January the Anchoviella dominated the diet as in May (Table 3b).

### Food variations according to size

710 Spotted seer in the length range between 31 mm and 720 mm, sampled during the period 1967-'69 from zones I and II were divided into 20 length groups at 30 mm intervals. Fish in the range 61 mm to 300 mm were found to have fed exclusively on *Anchoviella* which dominated in the succeeding group, but progressively reduced in the succeeding length ranges and was absent in fish beyond 601 mm. *Sardinella* first appeared in the diet of fish in the 301-330 mm group

| 100 10 10 10 10 10 10 10 10 10 10 10 10 | 904   | <u>55</u> | . 3  | 2.8.3 | - <u>-</u> | 1. 14. – |       |       |              |            | <u>5</u> . 3 | - 24 E | 2.81 |
|---|-------|-----------|------|-------|------------|----------|-------|-------|--------------|------------|--------------|--------|------|
| Months                                  | Aug.  | Sep.      | Oct. | Nov.  | Jan.       | Feb.     | Mar.  | Apr.  | May          | Jun.       | Jul.         | Total  | Rank |
| No. of fish examined                    | 22    | 32        | 18   | 55    | 26         | 34       | 51    | 59    | 6 <b>4</b> 0 | 38         | 30           | 405    | 12   |
| Sardinella                              | 96.19 | 20.14     | 100  | 99.94 | 99.68      | 99.87    | 99.42 | 95.95 | 87.01        | 75.31      | 28.63        | 98.306 | 10,1 |
| Anchoviella                             | 3.81  | 60.87     | 0    | 0     | 8.0        | 0.07     | 0.02  | 0     | 0.05         | 2.88       | 52.15        | 0.438  | 3    |
| Carangids                               | 0     | 9 8 0     | 0    | 0     | 0          | 0        | 0     | 80    | 0 0          | 0          | 5.11         | 0.004  | 5    |
| 'Fishes'                                | 0     | 18.88     | 0    | 0.06  | 0.28       | 0.02     | 0.4   | 0.18  | 0.23         | 0.55       | 3.07         | 0.337  | 4    |
| Fish scales                             | 0     | 0         | 0    | 0     | 0          | 0        | 0     | 0     | 0.68         | 0          | 0            | 0.004  | 5    |
| Prawns                                  | 0     | 0.11      | 0    | 0     | 0          | 0        | 0     | 80    | 0            | ⊂ <b>0</b> | 0            | 0.002  | 6    |
| Digested matter                         | 0     | 0         | 0    | 0     | 0.04       | 0.04     | 0.16  | 3.87  | 12.03        | 21.26      | 11.04        | 0.912  | 2    |
| Total                                   | 100   | 100       | 100  | 100   | 100        | 100      | 100   | 100   | 100          | 100        | 100          | 100    |      |

TABLE 3a. IP of organisms in the diet according to months for zone II (northern Gulf of Mannar) during 1967-'68.

TABLE 3b. IP of organisms in the diet according to months for zone II during 1968-69.

| <u> </u>             | S. G. M. | <u>11 2 3 0</u> |       | Q     | 0 12 9 2 |       | 214   |       |       |       |          |
|----------------------|----------|-----------------|-------|-------|----------|-------|-------|-------|-------|-------|----------|
| Months               | Aug.     | Sep.            | Oct.  | Nov.  | Dec.     | Jan.  | Mar.  | April | May   | Total | Rank     |
| No. of fish examined | 45       | 18              | 32    | 13    | 24       | 9     | 22    | 51    | 26    | 240   | apy.     |
| Sardinella           | 80.78    | 82.3            | 85.32 | 92.29 | 3.54     | 0     | 92.46 | 82.59 | 7.85  | 80.29 |          |
| Anchoviella          | 7.27     | 17.49           | 0.69  |       | 0        | 0     | 8 0   | 0.34  | 50.26 | 2.69  | 8 X 3    |
| 'Fishes'             | 0.33     | 0               | 1.35  | 0     | 0        | 0     | 0     | 0.71  | 0     | 0.02  | 8 1 B    |
| Fish scales          | 0 8 0    | 0               | 0     | 1.63  | 12.42    | 7.69  | 0.45  | о     | 0     | 0.44  | la la la |
| Loligo "             | 0.84     | 1 2 <b>0</b> 0  | 0     | 0     | 0        | 0     | 0     | o     | 0     | 0.03  | 5 ≥ 5    |
| Digested matter      | 10.78    | 0.11            | 12.64 | 6.08  | 84.04    | 92.31 | 7.09  | 16.36 | 41.89 | 16.53 | 11.2     |
| Total                | 100      | 100             | 100   | 100   | 100      | 100   | 100   | 100   | 100   | 100   |          |

ETTORS M. DEVARAL AUGUST (D)

| Length groups (mm)       | 61-90   | 91-120   | 121-1 50     | 151-180 | 181-210    | 271-300 | 301-330 | 331-360 | 361-390 | 391-420 |
|--------------------------|---------|----------|--------------|---------|------------|---------|---------|---------|---------|---------|
| No. of fish examined     | L       | 6        | 2 H 19       | 1 1 5   | 5          | ÷ 2.5   | 15      | 31      | 34      | 62      |
| Food items               | 2       | E B B    | al de la com | 4 - 2   | da         | 100     |         | in a    |         | in a b  |
| Sardinella               | 6 5 5 0 | • 0      | 0            | 0       | 0          | 1 0     | 11.56   | 83.01   | 78.8    | 86.95   |
| Anchoviella              | 100     | 100      | 100          | 100     | 100        | 100     | 57.81   | 12.09   | 19.32   | 9.67    |
| 'Fishes'                 | 0       |          | 0            | 0       | 0 00       | 0       | 0       | 0       | 0.54    | 0.01    |
| Fish scales              | 0       | o o      | 0            | 0       | . 0        | 8 o     | 2.89    | 0.29    | 0       |         |
| Digested matter          | 0       | 0        | 0            | 0       | <b>0</b>   | 0       | 27.74   | 4.61    | 1.34    | 3.36    |
| Total                    | 100     | 100      | 100          | 100     | 100        | 100     | 100     | 100     | 100     | 100     |
| Length groups (mm)       | 421-450 | 451-480  | 481-510      | 511-540 | 541-570    | 571-600 | 601-630 | 631-660 | 661-690 | 691-720 |
| Length groups (mm)       | 421-450 | 451-480  | 481-510      | 511-540 | 541-570    | 571-600 | 601-630 | 631-660 | 661-690 | 691-720 |
| No. of fish examined     | 93      | 104      | 100          | 95      | 59         | 36      | 33      | 8       | 6       | 1       |
| Food items in a pair (1) |         |          |              |         |            | 714.2   |         |         |         |         |
| Sardinella               | 97.41   | 98.07    | 76.48        | 95.97   | 91.91      | 93.05   | 91.66   | 98.92   | 97      | 100     |
| Anchoviella              | 0.49    | 0.02     | 1.69         | 0.01    | 2.5        | 0.62    | Ó       | 0       | 0       | 0       |
| 'Fishes'                 | 0.18    | 0.14     | 1.77         | 0.41    | 2.16       | 0.87    | 0.59    | 0.17    | 0       | 0       |
| Fish scales              | 0       | 0        | 0            | 1.23    | 10014 0.14 | 0       | 0       | 0       | 0       | 0       |
| Digested matter          | 1.92    | 233 1.72 | 20.08        | 2.38    | 3.43       | 5.46    | 7.75    | 0.91    | 1 113   | 0       |
| Total                    | 100     | 100      | 100          | 100     | 100        | 100     | 100     | 100     | 100     | 100     |

TABLE 4. IP of organisms in the diet according to length groups for zones I and II during 1967-69.

II (northern (hulf of Monuter) during 1967-68 and 1968-69.

(NRL 5. Fabri wishing of from, everage ration (R) and ration per unit body workfit of 1000 g (B.1) to not according to months for somer 1 (Palk Bay) and

FOOD AND FEEDING HABITS OF SPOTTED SEER

| Bav   | 2       |
|-------|---------|
| alk   |         |
| I (P  | 1       |
| San   |         |
| 102   |         |
| s for |         |
| mthe  |         |
| m     |         |
| R to  | ,       |
| ordin |         |
| acot  |         |
| THE   |         |
| in (  |         |
| (RI   |         |
| 0 8   |         |
| 100   |         |
| fo 1  |         |
| eigh  |         |
| ž     |         |
| poq   |         |
| unit  | Ş.      |
| Der   | 1968    |
|       | pu      |
| rati  | 88<br>8 |
| and   | 10      |
| R     | 8 19    |
| ion   | urin    |
| rat   | r) d    |
| rage  | nna     |
| ave   | Ma      |
| Xod,  | if of   |
| of fe | 3       |
| me    | hern    |
| volu  | north   |
| otal  | 11 (9   |
| T.    |         |
| LE S  |         |
| LVB   |         |

|                         | monter for fam |       | 10/1 9 | L'I THUN O |        |       |         |           |        |      |                      |       |          |
|-------------------------|----------------|-------|--------|------------|--------|-------|---------|-----------|--------|------|----------------------|-------|----------|
| 1967-68<br>Moaths       | Aug.           | Sep.  | Oct.   | Nov.       | Dec.   | Jan.  | 00 Feb. | Mar.      | Apr    | May  | 100 <sup>J</sup> un. | Jul.  | Combined |
| Mean weight of fish (g) | 478.2          | 527.7 | 587.5  | 588.2      | 567.7  | 592.5 | 684.2   | 696.4     | 640    | 640  | 1027.7               | 437.7 | 727.1    |
| Total melana            | 17             | 10 22 | 55     | 55         | 5      | 32    | 8       | <b>S1</b> | 20     | 4    | 42                   | 38    | 453      |
|                         | FI -           | 6.61  | 145.8  | 221.9      | 33.5   | 100.4 | 161.7   | 187.3     | 177.3  | 80.8 | 24.5                 | 33    | 1268.1   |
| A more K                | 0.7            | 1.9   | 4.4    | 4          | 6.7    | 3.1   | 4.7     | 3.6       | 3      | 2.2  | 0.5                  | 0.8   | 2.7      |
| RI                      | 1.4            | 3.6   | 7.4    | 6.8        | 11.8   | 5.2   | 6.8     | 5.1       | 4.6    | 2.1  | 1.1                  | 1.1   | 4.4      |
| 1968-69<br>Months       | Aug.           | 98.0  | Sep.   | Oct.       | Nov.   | 100   | Dec.    | Jan.      | Mar.   | 10   | Apr.                 | May.  | Combined |
| Mean weight of fish (g) | 675.6          | l s   | 23.3   | 818.1      | 1151.5 |       | 750     | 714.2     | 1187.5 |      | 112.5                | 800   |          |
| No. of fish examined    | 55             |       | 24     | 33         | 13     |       | 24      | 6         | 22     |      | 51                   | 56    | 257      |
| Total volume            | 131.5          |       | 50.3   | 27.3       | 50.5   |       | 14.4    | 2.5       | 40.8   |      | 61                   | 4.7   | 383      |
| rudig Rools (am)        | 23             |       | 2      | 0.8        | 3.8    |       | 9.0     | 0.2       | 1.8    |      | ņ                    | 0.1   | 1.4      |
| RI                      | 3.4            |       | 3.8    | 0.0        | 33     |       | 0.8     | 0.2       | 15     | 3    | 13                   | 0.05  |          |

M. DEVARAJ

and formed the major diet of all groups above 331 mm Prawns, squids, and carangids were found to have been fed upon by fish in the length groups of 391-420 mm, 451-480 mm and 481-510 mm respectively, in very negligible quantities (Table 4).

Variations in average ration and ration per unit body weight

In 1967-'68, the spotted seer consumed an average ration (R) of 2.7 ml food per one active feeding period (vide infra). R ranged from 0.5 ml in June to 6.7 ml in December . R was less than 2.2 ml from May to September, and higher than 3.0 ml from October to April (Table 5). The ration taken by fish of different length groups steadily increased from 0.2 ml in the 121-150 mm group to 6.0 ml in the 661-690 mm group, with only minor fluctuations between this range. The fish in the largest length group, 691-720 mm, had a ration of 50 ml food (Table 6).

In 1968-'69, the average ration dropped to 1.4 ml. The ration ranged from 0.1 ml in May to 3.8 ml in November. In spotted seer less than 300 mm in length, the ration ranged from 0.05 ml to 1.0 ml. In fish ranging from 301 mm to 630 mm length, ration never exceeded 2.1 ml except in the 331-360 mm group (4.3 ml). In the largest length groups, 631-660 mm and 661-690 mm, it attained the maximum values of 8.8 ml and 4.3 ml respectively (Table 6).

In 1967-'68, ration per unit body weight of 1000 g  $(R_1)$  was found to be 4.4 ml.  $R_1$ ranged from 1.1 ml in June and July to 11.8 in December.  $R_1$  recorded 10.3 to 13.8 ml in fish ranging from 121 mm to 210 mm length. It declined to 7.1 ml in the 271-300 mm group. In the length groups of 301-690 mm,  $R_1$  varied from 2 ml to 5.5 ml except for the 361-390 mm group in which it recorded the lowest of

110

and

|                      |                             | the state of the second st |              |                        |             |             |         |              |        |                |
|----------------------|-----------------------------|--|--------------|------------------------|-------------|-------------|---------|--------------|--------|----------------|
| 1967-'68             |                             |  | 10 - Q.M.    | 0 0                    | 7 S.T       | 1968-'69    | 0.7 5.8 | 0.3          | 9.0    | 0 7            |
| Length group<br>(mm) | Mean weight<br>of fish in g | No. of fish<br>examined  | Total volume | Average<br>ration :(R) | Body weight | No. of fish | volume  | 078 <b>R</b> |        | R <sub>1</sub> |
| 31-60                | 1.2                         | 0  | 0.0          | 0.0                    | 0.0         | 1           | 0.0     | 0.0          | 1      | 0.0            |
| 61-90                | 2.9                         | 0  | 0.0          | 0.0                    | 0.0         | 4           | 0.5     | 0.1          | 0      | 34.5           |
| 91-120               | 7.2                         | 0  | 0.0          | 0.0                    | 0.0         | 6           | 1.2     | 0.2          | × .    | 27.8           |
| 121-150              | 19.4                        | 2  | 0.4          | 0.2                    | 10.3        | 7           | 0.5     | 0.1          | - 11   | 3.6            |
| 151-180              | 29.4                        | 2  | 0.8          | 0.4                    | 13.6        | 3           | 2.0     | 0.6          | -01    | 20.4           |
| 181-210              | 36.0                        | 1  | 0.5          | 0.5                    | 13.8        | 4           | 0.2     | 0.1          | 0      | 1.4            |
| 271-300              | 140.0                       | 4  | 4.0          | 1.0                    | 7.1         | 1           | 1.0     | 1.0          |        | 7.1            |
| 301-330              | 199.4                       | 11   | 4.4          | 0.4                    | 2.0         | 2           | 4.3     | 2.1          |        | 11.0           |
| 331-360              | 247.6                       | 30   | 20.8         | 0.7                    | 2.8         | 0 07        | 4.3     | 4.3          | 10     | 17.3           |
| 361-390              | 321.7                       | 25   | 24.9         | 1.0                    | 3.1         | 9           | 10.2    | 0 1.1        | 0      | 3.4            |
| 391-420              | 389.5                       | 45   | 96.7         | 2.1                    | 5.3         | 17          | 16.8    | 0.9          | 0      | 2.3            |
| 421-450              | 476.3                       | 58   | 136.6        | 2.3                    | 4.8         | 35          | 66.5    | 0 1.8        | 9      | 3.7            |
| 451-480              | 594.0                       | 76   | 240.3        | 3.3                    | 5.5         | 28          | 17.3    | 0.6          | 0      | 1.0 58         |
| 481-510              | 633.2                       | 69   | 239.9        | 3.4                    | 5.3         | 31          | 51.6    | 018          | 0      | 2.5            |
| 511-540              | 864.3                       | 65   | 250.8        | 3.8                    | 4.3         | 30          | 48.8    | 0 1.6        | -0     | 1.7            |
| 541-570              | 992.9                       | 33   | 83.3         | 2.5                    | 2.5         | 26          | 30.3    | 0 1.1        | 0      | 0 1.1 13       |
| 571-600              | 1258.0                      | 11   | 28.2         | 2.7                    | 2.1         | 25          | 48.6    | 0 1.9        | ġ.     | 1.5            |
| 601-630              | 1433.5                      | 13   | 56.4         | 4.3                    | 2.9         | 20          | 35.4    | 1.7          | ď.     | 1.1            |
| 631-660              | 1586.2                      | 5  | 18.1         | 3.6                    | 2.2         | 3           | 26.5    | 8.8          |        | 5.5            |
| 661-690              | 1822.5                      | 2  | 12.0         | 6.0                    | 3.2         | an 201-240  | 17.3    | 4.3          |        | 2.4            |
| 691-720              | 2070.0                      | int of mation)   | 50.0         | 50.0                   | 24.1        | 0           | 0.0     | 0.0          | 00 401 | 0.0            |
| Combined             | 13125.2                     | 453  | 1268.1       | 2.7                    | 4.4         | 257         | 383.3   | 1.4          | 1000   | 2.0            |

TABLE 6. Total volume of food, average ration per unit body weight of 1000 g (R1) in ml according to length groups for Zone I and Zone II during 1967-58 and 1968-69

FOOD AND FEEDING HABITS OF SPOTTED SEER

| 1974 and       |                |         |         | -1      |         | 0       |         | 2011    |         |         |         | <u>[]</u> |         |         |         |                  | 37620         |
|----------------|----------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|-----------|---------|---------|---------|------------------|---------------|
| Length<br>(mm) | groups         | 271-300 | 301-330 | 331-360 | 361-390 | 391-420 | 421-450 | 451-480 | 481-510 | 511-540 | 541-570 | 571-600   | 601-630 | 631-660 | 661-690 | 691-720          | Com-<br>bined |
| A. Male        |                |         |         |         |         | 18.1    |         | 3.6     |         | 12      |         | F         | 26.3    |         | 8.8     | ан <sup>ан</sup> | 22            |
| Immature       | R              | 2       | 1       | 1.6     | 1.6     | 1.8     | 4       | 0       | 0       | 0       | 0       | 0         | 0       | 0       | 0       | 0                | 2             |
| 211-60         | R <sub>1</sub> | 14.3    | 5       | 6.5     | 5       | 4.7     | 8.4     | 0       | 0       | 0       | 0       | 0         | 0       | 0       | 1.9 0   | 0                | 7.3           |
| Intermedia     | te R           | 0       | 8.5     | 0.3     | 1.2     | 2.8     | 3       | 2.6     | 1.1     | 3.4     | 2.5     | 0         | 0.5     | 0       | 0       | 0                | 1.8           |
| 511.54         | R <sub>1</sub> | 0       | 2.6     | 1.3     | 3.7     | 7.2     | 6.3     | 4.4     | 1.7     | 3.9     | 2.5     | 0         | 0.4     | 0       | 0       | 0                | 3.4           |
| Maturing       | R              | 0       | 0       | 0       | 0.5     | 1.1     | 3.2     | 3.3     | 2.7     | 1.8     | 0.5     | 3.3       | 0       | 0.8     | TR 0    | 0                | 1.9           |
| 151-485        | R <sub>1</sub> | 0       | 0       | 0       | 1.6     | 2.8     | 6.6     | 5.5     | 4.3     | 2.1     | 0.5     | 2.6       | 0       | 0.5     | 0       | 0                | 2.9           |
| Ripe R         |                | 0       | 0       | 0       | 0       | 0       | 2.5     | 0.7     | 11.9    | 0.7     | 0       | 0.3       | 0       | 0       | 118 0   | 0                | 3.2           |
| R <sub>1</sub> |                | 0       | 0       | 0       | 0       | 0       | 7.7     | 1.2     | 18.8    | 0.8     | 0       | 0.2       | 0       | 0       | 0.90    | 0                | 5.7           |
| Spent R        |                | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 0.5     | 0         | 0       | 0       | 11 O    | 0                | 0.2           |
| R <sub>1</sub> |                | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 0.5     | 0         | 0       | 0       | 0       | 0                | 0.2           |
| B. Female      | ,              | 1400    |         | 4       |         | 90      |         | TO      | 3       | T       | , i i i | /         | 10      |         | 1.0     |                  | 11            |
| Immature       | R              | 0       | 0.2     | 0.8     | 0.6     | 2.4     | 0.8     | 072 - 4 | 0       | 0       | 0       | 0         | 0       | 0       | 01      | 0                | 0.9           |
|                | R <sub>1</sub> | 0       | 0.2     | 3.2     | 2       | 6.3     | 1.7     | 6.7     | 0       | 0       | 0       | 0         | 0       | 0       | 0,6     | 0                | 3.5           |
| Intermedia     | te R           | 0       | 0       | 1.8     | 0       | 1.8     | 3.8     | 4.6     | 3.3     | 4       | 0.4     | 1.7       | 2       | 0.8     | 01      | 50               | 6.1           |
|                | <b>R</b> 1     | 0       | 0       | 7.3     | 0       | 4.6     | 8       | 7.7     | 5.2     | 4.6     | 0.4     | 1.3       | 1.4     | 0.5     | 0.2     | 24.2             | 5.1           |
| Maturing 1     | R              | 0       | 0       | 0       | 0       | 0       | 8       | 0       | 3.7     | 7       | 4.1     | 8.2       | 1.9     | 16      | 017     | 0                | 7             |
| (mm)           | R <sub>1</sub> | 0       | . 0     | 0       | 0       | 0       | 16.8    | 0       | 5.8     | 8.1     | 4.1     | 6.5       | 1.3     | 10      | 010     | ) 0              | 7.5           |
| Ripe R         | mb ja          | 0       | 0       | 0 0     | 0       | 2.5     | 0       | 0.4     | 3.6     | 6.7     | 2.3     | 0.9       | 8.3     | 0.3     | ж       | 0                | 2.9           |
| R <sub>1</sub> |                | 0       | 0       | 0       | 0       | 6.4     | 0       | 0.7     | 5.7     | 7.7     | 2.3     | 0.7       | 5.8     | 0.2     | 0.6     | 6 0              | 3.3           |
| Spent R        |                | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 0       | 5       | 0.3     | 0.3       | 0       | 0       | (       | ) 0              | 1.9           |
| THE OB         |                |         | 0001 0  | antia a | uon lon | -       |         |         | 1.000   |         | 0.5     | 0.4       |         |         |         |                  |               |

TABLE 7. Average ration (R) and ration per unit body weight of 1000g (R1) in ml according to length groups and maturity stages for zones I (Palk Bay) and II (northern Gulf of Mannar) during 1967-68.

## FOOD AND FEEDING HABITS OF SPOTTED SEER

|                       | Imma                      | ature                 | Intermed                                | late                                | Maturi                                      | B                 | Rip            |                            | Spe                        | at a               |
|-----------------------|---------------------------|-----------------------|---|-------------------------------------|---|-------------------|----------------|----------------------------|----------------------------|--------------------|
| Length<br>groups (mm) | ×                         | R1                    | (93)<br>89)<br>8                        | R1                                  | able<br>sea th<br>c <b>B</b><br>med         | R1                | a<br>a         | R1                         |                            | R1                 |
| A. Male               | in la<br>la<br>Ta         | ai<br>ioi<br>io<br>io | 68-<br>tal<br>a.                        | 1320<br>1321<br>132                 | f)<br>el<br>tit                             | (5)<br>(4)<br>(4) | si q           | ion<br>tul<br>utu          |                            | 113<br>151<br>1549 |
| 301-330               | 4                         | 20.1                  | 0                                       | 0                                   | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| 331-360               | 0                         | 0                     | <b>1</b>                                | 8.1                                 | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| 361-390               | 2.5                       | 7.8                   | 1.8                                     | 5.6                                 | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| 391-420               | ig<br>be                  | 2.6                   | 15                                      | 3.9                                 | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| 421-450               | 0.3                       | 0.6                   | 2.2                                     | 4.6                                 | ę   | 9.0               | 2.5            | 5.2                        | 0                          | 0                  |
| 451-480               | 0                         | 0                     | 1.5                                     | 2.5                                 | 0   | 0                 | 2.1            | 3.5                        | 0.3                        | 0.5                |
| 481-510               | 0                         | 0<br>0<br>34          | 4.7                                     | 7.4                                 | 0   | 0                 | 0.8            | 13                         | 0.3                        | 0.4                |
| 511-540               | 0                         | 0                     | 13                                      | 1.5                                 | 0   | 0                 | 2.8            | 3.2                        | 6.5                        | 7.5                |
| 541-570               | 0                         | 0                     | 2.5                                     | 2.5                                 | 0   | 0                 | 6.0            | 6.0                        | 0.3                        | 0.3                |
| 271-600               | 0                         | 0                     | 0                                       | 0                                   | 0   | 0                 | 0.5            | 0.4                        | 0                          | 0                  |
| 601-630               | 0                         | 0                     | 0<br>019<br>019                         | 0                                   | 0   | 0                 | 0              | 0                          | 10.2                       | 1.1                |
| 631-660               | 0                         | 0                     | 0                                       | 0<br>0<br>0                         | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| 661-690               | 0                         | 0                     | 0                                       | 0                                   | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| Combined              | 1.7                       | 6.2                   | 2.2                                     | 4.5                                 | 1.5   | 3.1               | 1.6            | 2.4                        | 2.9                        | 2.6                |
| B. Female             | n s<br>ml)<br>7.5<br>c (T |                       |   |                                     | : ma<br>ont<br>onta<br>2.0                  |                   |                | 6).<br>4,5<br>10,5<br>20,4 | .lm<br>>)<br>>001          |                    |
| 271-300               | III III                   | 7.1                   | ible<br>oss<br>ces<br>dec               | 0                                   |   | 0                 | 0              | 0                          | 0                          | 0                  |
| 301-330               | 0.3                       | 1.5                   |   | 0                                   | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| 331-360               | 0                         | 0                     |   | 0                                   | 10<br>0<br>0<br>0<br>0<br>0                 | 0                 | 0              | 0                          | 0                          | 0                  |
| 361-390               | 0.2                       | 0.5                   | 0<br>(11)<br>(11)<br>(11)               | 0                                   | 0   | 0                 | 0              | 0                          | 0                          | 0                  |
| 391-420               | 0.1                       | 0.3                   | 000000000000000000000000000000000000000 | 0                                   | 0   | 0<br>sdi          | 0              | 0                          | 0                          | 0                  |
| 421-450               | 3                         | 6.3                   | 0                                       | 0                                   | 0<br>0<br>1011<br>8                         | 0                 | 0              | 0                          | 0                          | 0                  |
| 451-480               | 1.7                       | 2.9                   | 0.3                                     | 0.5                                 | 1 a<br>0<br>1<br>1<br>1<br>1<br>1<br>1<br>1 | 0                 | 90<br>10       | 0                          | 2.5                        | 4.2                |
| 481-510               | 0.5                       | 0.8                   | 0.7                                     | II.                                 | 0.2   | 0.3               | 0              | 0                          | 2.2                        | 3.5                |
| 511-540               | 0                         | 0                     | 0.8                                     | (1)<br>(12)<br>(12)<br>(12)<br>(12) | 0.5   | 9.0               | 1.1            | 1.3                        | 2.4                        | 2.7                |
| 541-570               | 0                         | 0                     | 0.4                                     | 0.4                                 | 0.8   | 0.8               | 0.9            | 0.0                        | 2.1                        | 2.1                |
| 571-600               | 0                         | 0                     | 1.6                                     | 1.3                                 | 1.5   | 1.2               | 0.8            | 0.6                        | 3.7                        | 2.1                |
| 601-630               | 0                         | 0                     | 0.7                                     | 0.5                                 | 0.3   | 0.2               | 0.4            | 0.3                        | 0.6                        | 0.5                |
| 631-660               | 0                         | 0                     | 0<br>T                                  | 0                                   | 0.5   | 0.3               | 13             | 8.2                        | 9                          | 3.5                |
| 661-690               | 0                         | 0                     | 0.2                                     | 0.1                                 | 0<br>(1                                     | 0                 | d)<br>2<br>130 | 3.8                        | 3<br>1<br>1<br>1<br>1<br>1 | 1.7                |
| combined              | a<br>a<br>a<br>a          | 2.8                   | 0.6                                     | 0.6                                 | 0.6   | 0.6               | 3.0            | 2.5                        | 28                         | 20                 |

M. DEVARAJ

0.3 ml. The exceptionally high value of  $R_1$  (24.1 ml) in the largest length group, 691-720 mm, is due to the high of R recorded for the only fish in this group (Table 6).

In 1968-'69,  $R_1$  declined to 2 ml. It remained at a low level in all months (< 1.5 ml) except in August (3.4 ml), September (3.8 ml) and November (3.3 ml) (Table 6).  $R_1$ recorded maximum values (27.8 to 34.5 ml) in the smaller length groups in the range of 61-120 mm. It ranged from 11.0 to 20.4 ml in the 151-180 mm, 301-330 mm and 331-360 mm groups, but less than 3 ml in most of the length groups beyond 361 mm (Table 6).

### Effect of maturation on feeding

In 1967-'68, among the males of different maturity stages, the values for R and  $R_1$  attained the maximum 3.2 ml and 5.7 ml in ripe males. Spent males consumed very poor ration (0.2 ml). In the immature, intermediate and maturing males, R did not vary much (1.8 to 2.0 ml), but  $R_1$  steadily decreased from 7.3 ml in immature to 2.9 ml in the maturing male, apparently because of size effect (Table 7). In the following year (1968-'69), the values of Rfor the males did not vary significantly from one maturity stage to the other (Table 8).

The values for  $R_1$  steadily increased from immature to maturing females, but declined in the ripe and spent females in the first year of the study. The length groups to which the intermediate, maturing and ripe females belonged were nearly the same, and hence, the drop in the ration in the ripe females (R = 2.9)ml;  $R_1 = 3.3$ . ml) from the high levels in the intermediate (R = 6.1 ml;  $R_1 = 5.1$  ml) and maturing females (R = 7.0 ml;  $R_1 = 7.5$  ml) might indicate slackening in food intake (Table 8). On the other hand, the ripe females in the following year (1968-'69) consumed more food  $(R = 3.9 \text{ ml}; R_1 = 2.5 \text{ ml})$  than the intermediate and maturing females (R and  $R_1 = 0.6$  ml each) (Table 8).

With increase in fish length, R exhibited generally an increasing trend, and  $R_1$  a decreasing trend in both sexes (Tables 8 and 9). (Since the results of the observation in the first year are not repeated the following year, no definite conclusions could be arrived at regarding the effect of gonad maturation on feeding). However, the overall results of both years' observations seem to indicate that the maturing and ripe females might be feeding as normally as the immature and intermediate fish.

#### Condition of food and feeding periodicity

In 1967-'68, all the samples were taken from the catches by drift gillnets operated from about 6.30 p.m. to 4 a.m. (first haul at 10 p.m. and second haul at 4 a.m.). 41.7% of the fish examined were well or partly fed and the rest poorly fed or starving (Table 9). The stomachs of fish in the samples from the second haul were mostly empty, but the analysis of the first haul material, which formed the bulk of the samples, showed the duration between 7 p.m. and the time of first haul at about 10 p.m. to be one of the active feeding periods of the spotted seer.

In the second year (1968-'69), all fish below 150 mm in length were taken from shore seines operated from about 5 a.m. to 10 a.m. and fish above 150 mm in length from the drift gillnets operated between about 6.30 p.m. and 4 a.m. The very low ration received by the fish in the second year of observation was also confirmed by the degree of fullness of the stomachs. During this year, 62.7% of the fish examined were poorly fed and 17.1% starving. Well fed and partly fed fish accounted for only 20.2% of the total (Table 9). In the pooled shore seine samples, 50 to 75% of fish (61-120 mm in length) were found well fed indicating that the spotted seer fed actively for some period in the forenoon also.

| (init)             | TABLE 9. | Feedin | g intensity | according      | to mont | hs for zon      | es I (Palk | k Bay) and      | II (northe     | ern Gulf o                      | of Mannar)                   | during 19 | 067-'69. |          |
|--------------------|----------|--------|-------------|----------------|---------|-----------------|------------|-----------------|----------------|---------------------------------|------------------------------|-----------|----------|----------|
| Months<br>1967-'68 | 광 700    | Aug.   | Sep.        | Oct.           | Nov.    | Dec.            | Jan.       | Feb.            | Mar.           | April.                          | May.                         | Jun.      | Jul.     | Combined |
| Well fed           | No.      | 2      | 4           | 8              | 12      | 1               | 6          | 12              | 8              | 11                              | 5                            | 2         | 1        | 72       |
|                    | %        | 7.4    | 10.8        | 24.3           | 21.9    | 20              | 18.7       | 35.3            | 15.6           | 18.7                            | 12.5                         | 4.8       | 2.6      | 15.8     |
| Partly fed         | No.      | 6      | 12          | 12             | 16      | 3               | 15         | 10              | 15             | 9                               | 10                           | 5         | 4        | 117      |
|                    | %        | 22.3   | 32.8        | 36.4           | 29      | 60              | 46.9       | 29.5            | 29.5           | 15.2                            | 25                           | 11.9      | 10.5     | 25.9     |
| Poorly fed         | No.      | 2      | 10          | <sub>0</sub> 7 | 13      | 2d. 0           | 0 3379.4   | 5               | 10             | 29                              | 17                           | 16        | 16       | 128      |
|                    | %        | 7.4    | 27.1        | 21.2           | 21.9    | 0               | 12.5       | 14.7            | 19.6           | 49.2                            | 42.5                         | 38.1      | 42.1     | 28.2     |
| Starving           | No.      | 17     | 11          | 6              | 15      | 1               | 7          | 7               | 18             | 10.8                            | 19                           | 17        | 136      |          |
|                    | %        | 62.9   | 29.7        | 18.1           | 27.2    | 20              | 21.9       | 20.5            | 35.3           | 16.98                           | 20                           | 45.2      | 44.8     | 30.1     |
| Total              | a: 0     | 27     | 37          | ° 33           | 55      | 5               | 32         | 34              | 51             | 59                              | 40                           | 42        | 38       | 453      |
| 1968-'69           |          |        |             |                | 0 3     | 4 1             | 80 - 3     | 15 18           | 20             | 15 IQ                           | - ð - 3                      | e 3       | 1 1      | 1 189    |
| Well fed           | No.      | °7     | 3           | 0              | 2       | 0               | 0          | 0               | 117 <b>1</b> 8 | 1 <sup>2</sup> - <sup>2</sup> 1 | 0                            | 0         | 0        | 14       |
|                    | %        | 12.7   | 12.5        | 0              | 15.1    | 0               | 0          | <sup>13</sup> 0 | 4.5            | 1.9                             | <sup>13</sup> 0 <sup>4</sup> | 0         | 0        | 5.4      |
| Partly fed         | No.      | 8      | 5           | 5              | 3       | 2               | 0          | 0               | 4              | 11                              | 0                            | 0         | 0        | 38       |
|                    | %        | 14.5   | 20.8        | 15.1           | 23.1    | 8.3             | 0          | 0               | 18.2           | 21.6                            | 0                            | 0         | 0        | 14.8     |
| Poorly fed         | No       | 26     | 3           | 23             | 7       | 22              | 9          | 0               | 12             | 34                              | 25                           | 0         | 0        | 161      |
|                    | %        | 47.3   | 12.5        | 69.8           | 53.9    | 91.7            | 100        | 0               | 54.6           | 66.7                            | 96.2                         | 0         | 0        | 62.7     |
| Starving           | No.      | 14     | 13          | 5              | 1       | 0               | 0          | 0               | 5              | 5                               | 1                            | 0         | 0        | 44       |
|                    | %        | 25.5   | 54.2        | 15.1           | 7.7     | 330733 <b>0</b> | 0          | -30431-40       | 22.7           | 9.8                             | 3.8                          | 0.001     | 0        | 17.1     |
| Total              |          | 55     | 24          | 33             | 13      | 24              | 9          | 0               | 22             | 51                              | 26                           | 0         | 0        | 257      |

FOOD AND FEEDING HABITS OF SPOTTED SEER

| Length<br>groups (mm) | 1   | 31-60 6 | 1-90 9 | 1-120 1 | 21-1501 | 51-18018 | 1-2102 | 71-30030 | 1-3303 | 31-3603 | 61-3903 | 91-4204 | 21-4504 | 51-4804 | 81-5105 | 11-5405 | 41-5705 | 71-6006        | 01-6306 | 31-6606 | 61-6906 | 91-720 | Com-<br>bined |
|-----------------------|-----|---------|--------|---------|---------|----------|--------|----------|--------|---------|---------|---------|---------|---------|---------|---------|---------|----------------|---------|---------|---------|--------|---------------|
| 1967-'68              |     |         |        |         | 12      |          | с.     |          |        | 10      |         |         | 20      |         | 1       |         |         | 1.             |         | 0       | 0       |        | 10            |
| Well fed              | No. | 0       | 0      | 0       | 2       | 0        | 0      | 0        | 0      | 1       | 1       | 8       | 12      | 15      | 10      | 18      | 4       | 2              | 1       | 1       | 0       | 0      | 75            |
|                       | %   | 0       | 0      | 0       | 100     | 0        | 0      | 0        | 0      | 3.3     | 4       | 17.8    | 20.6    | 19.8    | 14.4    | 27.7    | 12.1    | 18.2           | 7.6     | 20      | 0       | 0      | 16.5          |
| Partly fed            | No. | 0       | 0      | 0       | 0       | 2        | 1      | 2        | 3      | 5       | 8       | 10      | 14      | 22      | 21      | 16      | 7       | 3              | 4       | 0       | 1       | 0      | 11.9          |
|                       | %   | 0       | 0      | 0       | 0       | 100      | 100    | 50       | 27.2   | 16.7    | 32      | 22.2    | 24.2    | 28.9    | 30.5    | 24.6    | 21.2    | 27.2           | 30.8    | 0       | 50      | 0      | 26.3          |
| Poorly fed            | No. | 0       | 0      | 0       | 0       | 0        | 0      | 0        | 4      | 5       | 8       | 12      | 14      | 13      | 21      | 18      | 13      | <sup>0</sup> 4 | 5       | 3       | 0       | 0      | 120           |
|                       | %   | 0       | 0      | 0       | 0       | 0        | 0      | 0        | 36.4   | 16.7    | 32      | 26.7    | 24      | 17.1    | 30.5    | 27.7    | 39.4    | 36.4           | 38.5    | 60      | 0       | 0      | 26.5          |
| Starving              | No. | 0       | 0      | 0       | 0       | 0        | 0      | 2        | 4      | 19      | 8       | 15      | 18      | 26      | 17      | 13      | 9       | 2              | 3       | 1       | 1       | 1      | 139           |
|                       | %   | 0       | 0      | 0       | 0       | 0        | 0      | 50       | 36.4   | 63.3    | 32      | 33.3    | 31      | 34.2    | 24.6    | 20      | 27.3    | 18.2           | 23.1    | 20      | 50      | 100    | 30.7          |
| Total                 |     | 0       | 0      | 0       | 2       | 2        | 1      | 4        | 11     | 30      | 25      | 45      | 58      | 76      | 69      | 65      | 33      | 11             | 13      | 5       | 2       | 1      | 453           |
| 1968-69               |     |         |        |         |         |          |        |          |        |         |         |         |         |         |         |         |         |                |         |         |         |        |               |
| Well fed              | No. | 0       | 3      | 3       | 0       | 0        | 0      | 0        | 1      | 0       | 2       | 1       | 2       | 5       | 2       | 2       | 0       | 1              | 1       | 1       | 2       | 0      | 26            |
|                       | %   | 0       | 75     | 50      | 0       | 0        | 0      | 0        | 50     | 0       | 22.3    | 5.9     | 5.7     | 17.9    | 6.5     | 6.6     | 0       | 4              | 5       | 33.3    | 50      | 0      | 10.           |
| Partly fed            | No. | 0       | 0      | 0       | 2       | 0        | 2      | 1        | 0      | e0 1    | 1       | 2       | 9       | 1       | 1       | 3       | 8       | 6              | 2       | 1       | 102     | 0      | 4             |
|                       | %   | 0       | 0      | 0       | 28.6    | 0        | 50     | 100      | 0      | 100     | 11.1    | 11.7    | 25.7    | 3.5     | 3.2     | 10      | 30.8    | 24             | 10      | 33.3    | 25      | 0      | 15.9          |
| Poorly fed            | No. | 0       | 0      | 0       | 1       | 1        | 0      | 0        | 1      | 0       | 3       | 7       | 17      | 15      | 26      | 20      | 16      | 16             | 13      | 1       | 0       | 0      | 13            |
|                       | %   | 0       | 0      | 0       | 14.2    | 33.3     | 0      | 0        | 50     | 0       | 33.03   | 41.2    | 48.5    | 53.6    | 83.8    | 66.7    | 61.6    | 64             | 65      | 33.3    | 0       | 0      | 53.           |
| Starving              | No. | 1       | 1      | 3       | 4       | 2        | 2      | 0        | 0      | 0       | 3       | 7       | 7       | 2       | 5       | 2       | 2       | 4              | 0       | 1       | 0       | 53     |               |
|                       | %   | 100     | 25     | 50      | 57      | 66.7     | 50     | 0 0      | 0      | 0       | 33.3    | 41.02   | 20      | 25      | 6.5     | 16.7    | 7.6     | 8              | 20      | 0       | 25      | 0      | 20.           |
| Total                 |     | 1       | 4      | 6       | 7       | 3        | 4      | 1        | 2      | 1       | 9       | 17      | 35      | 28      | 31      | 30      | 26      | 25             | 20      | 3       | 4       | 0      | 25            |

TABLE 10. Feeding intensity according to length groups for zones I (Palk Bay) and II (northern Gulf of Mannar) during 1967-'69.

116

M. DEVARAJ

### FOOD AND FEEDING HABITS OF SPOTTED SEER

8

1967-

Mannar)

5

Gulf

H

and

from zones I (Palk Bay)

fish

5

stomach

the

forage in

5

orientation

Percentage

BLE

In 1967-'68, both well fed and partly fed fish together ranged from 44 to 80% during September to March, but declined to 13 to 38% in April to August during which a majority of fish (62 to 87%) were either poorly fed or starving (Table 9). In 1968-'69, no well fed fish were met with during October, December, January and May. The well fed and partly fed fish together formed more than 25% only in August (27.2%), September (33.3%) and November (38.2%), and less than 25% in other months. The incidence of poorly fed fish was remarkably high in all months (47.3 to 100%) but September (12.5%), and there were large number of starving fish (54.2%) (Table 9). The period of maximum incidence of well fed and partly fed fish was nearly the same in both the years.

Most fish of less than 300 mm length, which normally were distributed along the nearshore waters, were found to be well fed or partly fed both the years. However, in 1967-'68, in all the length groups between 361 mm and 690 mm which were normally taken from areas beyond the 20 meter depth line the incidence of these two categories together ranged from 33.3% to 52.3% (Table 10). In 1968-'69, both well fed and partly fed fish formed less than 33.3% in all the length groups from 361 mm to 630 mm and 66.7% and 75.0% respectively in the two largest groups, 631-660 mm and 691-720 mm (Table 10). Based on these observations, it is inferred that the juveniles find good pastures in the nearshore areas, while the adults meet with competition from other predators in the area beyond the 20 meter depth line. However, among the adult spotted seer, the larger fish do not seem to compete with the smaller fish, especially during years of forage abundance as in 1967-'68, but may do so in lean years as in 1968-'69.

| nthwise        |                 | kt<br>st | 190                                  | 1000      | 1        | -1.75  | 89<br>80 |                  | R)<br>26 | i da    | bah<br>ani<br>> 0 | ili<br>De |                         |          | 108<br>8193<br>8194 |                         |
|----------------|-----------------|----------|--------------------------------------|-----------|----------|--------|----------|------------------|----------|---------|-------------------|-----------|-------------------------|----------|---------------------|-------------------------|
| i y            | , 1             | Aug.     | Sep.                                 | Ō         | ct.      | Nov.   | Jan.     | E                | eb.      | Mar.    | Apr.              | May       |                         | une      | July                | Combined                |
| teriorly<br>d  |                 | 33.3     | 47.4                                 | vitas     | 33       | 34     | 100      | seer<br>as in    | 8        | 29.2    | 17.6              | 40.9      | ed fa<br>ba<br>1. Ba    | 06       | 6                   | 36.1<br>(78 actual)     |
| steriorly<br>d | and to<br>LOT22 | 66.7     | 52.6                                 | ۶<br>۵    | 1.3      | 8      |          | bettor<br>pilevo | 0        | 70      | 82.4              | 59.1      | a mug<br>bqA<br>a ta ta | 10       | 99                  | 63.9<br>(138 actual)    |
| troups (mm)    | 121-150 15      | 1-180 27 | 1-300 33                             | 31-360 36 | 51-390 3 | 91-420 | 421-450  | 151-480          | 481-510  | 511-540 | 541-570 5         | 11-600 60 | 1-630 6                 | 31-660 6 | 61-690 6            | 91-720 Com-             |
| teriorly       | 100             | . 001    | 100                                  | 33.3      | 25       | 37     | 22.7     | 37.5             | 36.7     | 35.1    | 28.6              | 45.5      | 20                      | 66.7     | 50                  | 0.36.1<br>(78 actual)   |
| steriorly      | 0               | 0        | 2, <b>6</b><br>,2, <b>6</b><br>98,1) | 66.7      | 75       | 63     | 77.3     | 62.5             | 63.3     | 64.9    | 71.4              | 54.5      | 80                      | 33.3     | 50                  | 10063.9<br>(138 actual) |

## Aggressiveness in feeding

That the spotted seer is a passive predator is suggested by the orientation of prey in their stomachs: 36.1% of the prey were found characterising anteroposteriorly oriented aggressive seizure and 63.9% of prey in the reverse, denoting passive capture (Table 11). percentage of reverse orientation which dominated in all the months ranged from 52.6% in September to 82.4% in April. However, in fish up to 300 mm in length which were found to have fed exclusively on anchovies, all the prey were anteroposteriorly oriented denoting aggressive predation. Beyond 300 mm and up to 630 mm passive predation dominated. Aggressive predation increased (66.7%) in only one group, i.e., 631-660 mm, but in the next group, 661-690 mm, both types of predation were in equal proportions while all fish in the largest group, 691-720 mm, exhibited passive predation (Table 11). Sardines formed the major food organisms of the spotted seer of length beyond 300 mm. The observations indicate that the adult spotted seer, however large they be, are not able to exert as much aggression on sardines as their juveniles are on anchovies.

### Food intake and utilisation

From the knowledge of a density of 1.25 for the stomach contents, 2 active feedings per day and the average ration per active feeding, the annual rations in grams wet weight for 1,2,3 and 4 year old spotted seer (Devaraj, 1981) were estimated by using the method followed in Devaraj (MS). The relation betwen grams wet weight of fish (W) and grams wet weight of annual ration ( $R\Delta t$ ) at age 1,2,3 and 4 years (Table 12) is found to be,

$$\ln R\Delta t = 0.6495 + 1.0722 \ln W \dots (1)$$

Since the annual ration for the 4 year old fish has been found to be extremely high owing to one fish in the sample, the above relation has been restricted to 1 to 3 year old fish only (Eq. 2),

$$\ln R\Delta t = 2.8902 + 0.6800 \ln W \dots (2)$$

For the same reason as above, data from 4 year old fish have been excluded in fitting T-line and K-line. The lines fitted for 1 to 3 year old fish are,

 $\ln T = 2.5637 + 0.6773 \ln W \dots (3)$  $\ln K = -1.2531 + 0.000007223 \text{ R}\Delta t \dots (4)$ 

Weight growth model developed by Palmoheimo and Dickie (1965) for species whose K-line has a slope close to zero,

W = {(1-q) c (t-t<sub>o</sub>) + W<sub>0</sub> 1-q}  $^{1/1-q}$  ... (5)

Where q is the slope (0.6773) of the T-line;  $c = pe^{-a}/(1-e^{-1})$  where p is the antilog of a-axis intercept of the T-line, and -a is the y-axis intercept of the K-line; and, W is the weight at 1 = 10.3 cm (derived from the von Bertalanffy equation: 127.8 Ļ  $(1-e^{-0.18007(t-0.4654)})$  (t = 0 year) which has been converted into  $W_0$  by using the length-weight relation  $W = 0.01011 \text{ L}^{2.8605}$  where W = weight in g and L = total length in cm (Devaraj, 1981). Substituting the values of q = 0.6773,  $W_0 = 7.98$ g, c = 5.1910 and  $t_0 = -0.4654$  year in Eq. (7), W for t = 1 year is found to be 99 g which is rather different from the observed value of 162 g and the von Bertalanffy estimate of 161 g. has bel lieve doed . 88 . 880 ]

Following Paloheimo and Dickie's (1965) view that their growth model holds only for a particular food type, the length of 31.5 cm at which the spotted seer begin to shift to their adult diet of sardines was considered as the  $1_0$  and the growth estimates made according to Eq. (5). Growth estimates thus obtained are consistently higher than those from von Bertalanffy growth function. This holds good for comparison with estimates obtained by summation of weight increments ( $\Sigma \Delta W$ ) derived from the difference between the annual ration increments (R $\Delta t$ ) and total metabolic rate

(T) even though the difference is of the order of 22% for ages 5 to 8 years but much higher during lower and higher ages. Thus the Paloheimo and Dickie (1965) weight growth model does not seem to be valid for spotted seer. Growth estimates by the summation of weight increments ( $\Sigma \Delta W$ ), derived from the differences between the annual ration increment (R $\Delta t$ ) and total metobolic rate (T), are found to agree with the empirical or von Bertalanffy estimates up to the fourth year of life, but significantly higher at subsequent ages (Table 13).

#### DISCUSSION

The spotted seer feed on a very limited range of about 5 food items, many of which are species low in the food chain, unlike the kingseer which feed on a large spectrum of items. In spite of this striking dissimilarity, the lesser sardines (Sardinella albella and S. gibbosa) were found to be the most favourite food of both the species, with whitebaits (anchovies), the most important among the supplementary diets, being in the much greater favour with the spotted seer than with the king seer. Between Palk Bay (zone I) and the northern Gulf of Mannar (zone II), the spotted seer diet included more of whitebaits in the former, apparently due to greater abundance, although they are not known to be of any commercial fishery importance in zone I. In contrast, the spotted seer, both juveniles and adults, along the Waltair coast (Lawson's Bay) are reported to feed primarily on whitebaits and secondarily on sardines (Srinivasa Rao, 1964), obviously due to the predominance of whitebaits in this area. Whitebaits are known to undertake diurnal vertical migration, but whether the spotted seer follow them to the sea bottom during daytime is not known. Despite substantial stocks of silverbellies and rainbow sardines in the bottom and midwaters of zone I, the total absence of these items in the diet may suggest the spotted seer to be a surface feeder as the king seer, although its demersal existence for a part of the day is known from its occurrence in the daytime bottom trawl catches along the northwest coast of India. Therefore, bottom feeding of spotted seer cannot be totally ruled out.

The contribution of sardines to the spotted seer diet in zone II declined marginally in 1968-69 from the previous year's level, apparently owing to the shortage of young sardines and predominance of older ones which the spotted seer could not more effectively prey upon unlike the king seer. In spite of a significant decline in the abundance of spotted seer in both zones I and II (Table 14), the decline in the spotted seer ration per active feeding in zones I-II in 1968-69 by about 50% of the 1967-68 level denotes acute scarcity of forage leading to an aggravation of the already severe competition from the king seer and other predators. That the decline in the king seer ration in 1968-69 was only 15% of the 1967-68 level is proof enough that the spotted seer is a far less competent species. Unlike the king seer which shifts to feeding on larger items like the sardines at a size about 1/13 of its asymptotic size, such a change over occurs at a much larger size (about 1/4 its  $L_{\infty}$ ) in the spotted seer, because of its far less aggressive nature which is also evident from most of its prey being oriented in its reverse axis (indicating passive predation as the prey comes face the predator). The presence of a relatively large number of gill rakers (8-12) on the outermost gill arch and a relatively less fusiform and much deeper body further support the view that this species is much less predatory than the kingseer (Nilson, 1958). The relatively poor spotted seer catches, forming about 33% (second rank), 18% (third rank) and 19% (third rank) of the average annual seerfish catches from the

TABLE 12. Empirical lengths in mm(L), weights in g(W), ration per one active feeding period in ml(R), R per 1000g body weight in ml(R<sub>1</sub>), annual R, i.e., R  $\Delta$  t in g, annual R<sub>1</sub> i.e., R  $\Delta$ t/1000 g W in g, conversion factor (C), growth efficiency (K) and annual total metabolic rate in g (T = R $\Delta$ t -  $\Delta$ Wt) according to age in years (t). Increments are given between brackets. N = numbers of fish studied for the estimation of food intake and utilisation parameters.

| 1 out | L        | W(\DeltaWt) | R          | <i>R</i> <sub>1</sub> | R∆t   | RΔt/1000 g<br>W | С    | K    | 5 <b>T</b> | N     |
|-------|----------|-------------|------------|-----------------------|-------|-----------------|------|------|------------|-------|
| 1.    | 297(297) | 167(167)    | 0.6(0.6)   | 11.4(11.4)            | 548   | 3281            | 3.3  | 0.3  | 381        | 268   |
| 2.    | 453(156) | 551(384)    | 1.7(1.1)   | 5.3(-6.1)             | 1551  | 2815            |      | 0.25 | 1167       | 358   |
| 3.    | 594(141) | 1188(637)   | 2.2(0.5)   | 2.6(-2.7)             | 2008  | 1690            | 3.2  | 0.32 | 1371       | 83    |
| 4.    | 705(111) | 1998(810)   | 12.4(10.2) | 6.4(3.8)              | 11315 | 5663            | 8 14 | 0.07 | 10505      | 3 8 1 |

TABLE 13. Estimated annual ration increment (R $\Delta t$ , Eq. 2), total metabolic rate (T; Eq. 5), weight increment ( $\Delta W$ ) and weight (W) at age in years, t (vB = weight estimate by the von Bertalanffy growth equation (Devaraj, 1981);  $\Sigma \Delta W$  = weight estimated by the summation of  $\Delta W$ ; PD = weight estimate by the Paloheimo-Dickie growth equation, Eq. 7; bR = reference point where b is the slope of the K - line of Eq. 6).

|      | R<br>(g) | T <sub>(g)</sub> | Δ <i>W</i><br>(g) | vB   | W(g)<br>Σ ΔW | PD<br>(for l <sub>o</sub> =10.3<br>cm) | PD<br>(for l <sub>o</sub> =31.5cm) | bR     |
|------|----------|------------------|-------------------|------|--------------|--|------------------------------------|--------|
| 1. H | 568      | 406              | 162               | 161  | 162          | 130                                    | 337                                | 0.0002 |
| 2.   | 1351     | 957              | 394               | 572  | 556          | 177                                    | 424                                | 0.0004 |
| 3.   | 2244     | 1581             | 663               | 1201 | 1219         | 234                                    | 524                                | 0.0007 |
| 4.   | 3132     | 2203             | 929               | 1957 | 2148         | 302                                    | 640                                | 0.001  |
| 5.   | 3962     | 2777             | 1185              | 2759 | 3333         | 383                                    | 772                                | 0.0012 |
| 6.   | 4707     | 3294             | 1413              | 3550 | 4746         | 477                                    | 920                                | 0.0015 |
| 7. 8 | 5357     | 3744             | 1613              | 4289 | 6359         | 585                                    | 1088                               | 0.0017 |
| 8.   | 5913     | 4130             | 1783              | 4957 | 8142         | 710                                    | 1275                               | 0.0018 |
| 9.   | 6384     | 4455             | 1929              | 5545 | 10071        | 851                                    | 1482                               | 0.002  |
| 10.  | 6779     | 4729             | 2050              | 6053 | 12121        | 1010                                   | 1711                               | 0.0021 |
| 11.  | 7106     | 4953             | 2153              | 6485 | 14274        | 1188                                   | 1963                               | 0.0022 |
| 12.  | 7377     | 5141             | 2236              | 6850 | 16510        | 1386                                   | 2239                               | 0.0023 |
| 13.  | 7600     | 5294             | 2306              | 7155 | 18816        | 1606                                   | 2541                               | 0.0024 |
| 14.  | 7782     | 5421             | 2361              | 7488 | 21177        | 1847                                   | 2868                               | 0.0024 |
| 15.  | 7930     | 5524             | 2406              | 7616 | 23583        | 2112                                   | 3223                               | 0.0025 |

M. DEVARAJ

Indian seas (1960-81), the Indian Ocean (1965-77) and the Pacific Ocean (1965-77) respectively, is attributable, *inter alia*, to the inability of this species to compete effectively for the available food.

However, the fact that in zone II, the spotted seer was numerically more abundant than the kingseer in both 1967-68 (2.4 times) and 1968-69 (1.2 times) would mean the former to be a more successful species than the latter, from the numerical angle. Therefore, its low abundance in terms of weight in this zone (1967-68: 0.75 times the kingseer; 1968-69): 0.7 times the kingseer; should be seen more as a function of its size, i.e., its growth characteristics ( $L_{\infty} = 1278$  mm;  $W_{\infty} = 8,540$ g; Devaraj, 1981) relative to the much larger kingseer ( $L_{\infty} = 2081 \text{ mm}; W_{\infty} = 39,027 \text{ g}$ ), than as a function of competence or competitiveness. One could add credence to this argument from the fact that the seerfish fishery in the neighbouring Palk Bay (zone I) comprised almost exclusively the spotted seer until about 1955 but progressively got replaced

by the kingseer (Table 14) not because of less competence, but because of the change in the salinity regime after the construction of the Vaigai Dam across the Vaigai river draining into Palk Bay.

Although the spotted seer is far less predatory than the kingseer and the streaked seer, on no occasion were planktonic items noticed in its diet. Moreover, Srinivasa Rao's (1964) observations from Waltair show that juvenile spotted seer feed predominantly on teleosts and negligible quantities of crustaceans while the adults feed on teleosts and macrocrustaceans. The insignificant occurrence of diatoms in the gut of 20-190 mm (total length) spotted seer from Vizhinjam together with the major items like fishes crustaceans and molluscs would appear to have resulted from their release from the gut of planktivorous forage fish already ingested and being digested by the spotted seer.

The fact that food intake by maturing and ripe fish did not decrease is not necessarily

TABLE 14. Age composition of spotted seer expressed in number of fish per 1000 units of 76 mm mesh type drift gillnet effort (boat days) for Palk Bay (zone I) and the northern Gulf on Mannar (zone II) during 1967-68 and 1968-69 together with a comparison of the total number and weight of spotted seer per 1000 units against the total number and weight of kingseer and streaked seer each per 1000 units (percentages between brackets).

| their amouth that may be considered | Palk        | Bay               | Northern Gulf   | f of Mannar  |
|-------------------------------------|-------------|-------------------|-----------------|--------------|
| Age group                           | 1967-68     | 1968-69           | 1967-68         | 1968-69      |
| 0                                   | 6 (0.4)     | 1 (0.32)          | 3 (0.02)        | 0            |
| and the process of 1000 consumption | 909 (60.52) | 167 (53.25)       | 7459 (52.44)    | 1695 (20.72) |
| h considered norms for zones [-1]   | 542 (36.09) | 126 (40.91)       | 6150 (43.24)    | 5400 (66.03) |
| in spite of severe intenspecific    | 44 (2.93)   | 17 (5.52)         | 597 (4.2)       | 1073 (13.12) |
| a in times of food scarcity. The    | 1 (0.07)    | s Ophiditions in  | 15 (0.11)       | 11 (0.13)    |
|                                     |             |                   | R. 1956; 1961), |              |
| Total No. of spotted seer           | 1502        | 220112-000 308mm  | 14224           | 8179         |
| Total weight of spotted seer (kg)   | 503         | nic officines the | 5374            | 4555         |
| Total No. of kingseer               | 8402        | 7246              | 5948            | 6916         |
| Total weight of kingseer (kg)       | 3603        | 2852              | 7192            | 6562         |
| Total No. of streaked seer          | 734         | 458               | 1677            | 1660         |
| Total weight of streaked seer (kg)  | 447         | 362               | 2433            | 2579         |

indicative of a lack of spawning stress as conditions of stress could indeed aggravate food intake to compensate the energy lost in stress. However, the marked fall in R increment between 2 and 3 years of age seems to suggest some kind of a depression following first maturation at an age of about 1.7 year, but that there has been a quick recovery is evident from a marked increase in the R increment as well as R<sub>1</sub> between 3 and 4 years of age. The steady decrease in R with age except at the 4th year denotes that young fish consume more food per unit body weight owing to much faster growth and much greater physical (e.g., swimming) activity. Evidently, comparatively younger fish, say, 1 to 3 year old fish, require much less amount of food (in absolute terms) to produce a unit growth (C = 3.2 to 4; K =0.25 to 0.32; T = 381 to 1371 g) while the 4 year old fish require large amounts of food to produce a unit growth (C = 14; K = 0.07; T = 10,505 g). However, in view of the very limited sample of 4 year old fish (N = 1; Table 13), no definite conclusion is possible, although at older ages close to the asymptotic age, growth efficiency is bount to be at its minimum.

The T-line fitted for 1 to 3 year spotted seer seems more realistic than that fitted for 1 to 4 year fish as the only 4 year old fish in the sample gave rather unrealistic values of  $R\Delta t$  and T (Table 13). Since the slope, q (which defines the rate of change of metabolism with body weight), of the T-line for 1 to 3 year old fish (0.6773) and for 1 to 4 year old fish (1.1534) are significantly different from the value of 0.8 describing the relation betwen metabolism and body weight in most fish species under normal non-stress conditions in nature (Winberg, 1956; 1961), the spotted seer do not appear to live in a normal non-stress state. The y-axis intercept (which defines the level of metabolic expenditure per unit time), p = 0.8555 (log p = -0.0678), of the T-line for 1 to 4 year old fish in zones I-II, denotes

low metabolic level, and hence, a ration only marginally higher than the maintenance ration, perhaps owing to competition from much abler predators including the kingseer. The maximum size attained by the spotted seer in differnt areas bears a distinct relation to its abundance in relation to the kingseer. For example, in zones I-II where the kingseer is the most dominant (88.38% by weight in 1964-76) among the seerfish, the spotted seer is much less abundant (8.14% by weight in 1964-76) and generally of smaller size (maximum recorded size is 723 mm) unlike along the northwest coast of India where it attains 800-900 mm length because of the generally poor abundance of the kingseer (1981 spotted seer catch by weight in total seerfish catch; 75% in the Bombay region; 72.6% in the Gujarat coast; 39% in the Maharashtra coast). In contrast, the T-line for 1 to 3 year old fish in zones I-II denotes a rather high metabolic level of 13.00  $(\log p = 1.1134)$  and high food intake, and therefore, the competition theory does not look plausible, in spite of reduced levels of R in 1968-69 when there was an obvious food shortage. What therefore emerges from this analysis is that the average ration available and taken by the spotted seer at least of 1 to 3 year age during 1967-69 was sufficient enough to sustain their growth that may be considered characteristic of the spotted seer stock in zones I-II, and that the process of food consumption and growth considered normal for zones I-II goes on in spite of severe interspecific competition in times of food scarcity. The K-lines also show that the general level of gross growth efficiency ( $K = e^{-a} = e^{-0.4603} =$ 0.63 of Eq.4 for 1 to 4 year old fish; K = $e^{-0.5405} = 0.58$  of Eq. 6 for 1 to 3 year old fish), which does not change from one species or food type to another, falls between the range

of 0.25 and 0.75 observed in feeding experiments, although the value of  $e^{-a}$  in nature is reportedly less than this level (Paloheimo and Dickie, 1965). Estimates of weight growth by the Paloheimo and Dickie (1965) model are far less than either the empirical or von Bertalanffy values in relation to younger age upto 7 years, apparently because of the slope of the *T*-line being significantly different from the value describing the relation between metabolism and body weight under normal nonstress conditions in nature. However, the higher values in the later years indicate recovery to normal growth condition.

REFERENCES

DESHPANDE, S. D. AND T. M. SIVAN. 1969. On the troll line investigations off Cochin during five fishing seasons. *Fish. Tech.*, 6 (1) : 26-35.

DEVARAJ, M., 1981. Age and growth of three species of seerfishes (Scomberomorus commerson, S. guttatus and S. lineolatus) in India. Indian J. Fish., 28: 104-127.

DHAWAN, R. M., P. V. S. NAMBOOTHIRI AND V. G. GOPINATHAN. 1972. Results of trolling line operations in Goa waters during 1965-68. *Ibid.* 16 : 181-187 (for 1969).

HASTINGS, W. H. AND L. M. DICKIE. 1972. Feed formulation and evaluation. In : Fish Nutrition. Academic Press, New York and London : 327-374.

IVLEV, V. S. 1955. Experimental ecology of the feeding of fishes. Translated from the Russian by Donglas Scott. New Haven, Yale University Press, 1961 : 1-302.

KUMARAN, M. 1964. Observations on the food of juveniles of *Scomberomorus commerson* (Lacepede) and *S. guttatus* (Bloch & Schneider) from Vizhinjam, west coast of India. Mar. Biol. Ass. India, *Proc. Symp. Scombroid Fishes.* Part-II : 586-590.

LEWIS, A. D., B. R. SMITH AND R. E. KEARNEY. 1974. Studies on tunas and baitfish in Papua New Guinea waters-II, Res. Bull. DASfP. Moresby (10): 1-112.

MOORE, H. B. 1972. Aspects of stress in the tropical marine environment. Adv. Mar. Biol. 10: 217-269.

NATARAJAN, A. V. AND A. G. JHINGRAN. 1962. Index of preponderance a method of grading the food elements in the stomach analysis of fishes. *Indian J. Fish.*, **8**: 54-59 (for 1961). NILSON, N. R. 1958. On the food competition between two species of *Coreqonus* in a North Swedish lake. *Inst. Freshw. Res. Drottingholm, Rept.*, (39).

PALOHEIMO, J. E. AND L. M. DICKIE. 1965. Food and growth of fishes. 1. A growth curve derived from experimental data. J. Fish. Res. Bd. Canada, 22: 521-542.

SILAS, E. G., 1967. Tuna fishery of the Tinnevelly coast, Gulf of Mannar. Mar. Biol. Ass. India, *Proc. Symp. Scombroid Fishes*, Part II : 1083-1117.

SLOBODKIN, L. B. 1961. Preliminary ideas for a productive theory of ecology. *Amer. Nat.*, **95** (882) : 145-153.

SRINIVASA RAO, K. 1964. Observations on the food and feeding habits of *Scomberomorus guttatus* (Bloch and Schneider) and juveniles of *S. lineolatus* (Cuvier and Valenciennes) and *S. commerson* (Lacepede) from the Waltair coast. Mar. Biol. Ass. India, *Proc. Symp. Scombroid Fishes*, Part II : 591-598.

SUBRAMONIA PILLAI, N., R. S. MANOHARDOSS AND SULOCHANAN, P., 1972. Standardisation of specifications for different trolling lures. *Fish. Tech.*, 9 (1): 68-75.

UNDP/FAO. 1974. Survey results 1972/73. Pelagic Fishery Project (IND 69/593), Progress Report, (6) : 1-141.

VENKATARAMAN, G. 1960. Studies on the food and feeding relationship of the inshore fishes off Calicut on the Malabar coast. *Indian J. Fish.*, 7 : 275-306.

WHITELY, G. P. 1964. Scombroid fishes of Australia and New Zealand. Mar. Biol. Ass. India, Proc. Symp. Scombroid Fishes, Part-I: 221-253.

WILLIAMS, F. 1964. The scombroid fishes of East Africa. Mar. Biol. Ass. India, *Ibid.*, Part I : 107-164. WINBERG, G. G. 1956. Rate of metabolism and food requirements of fishes. Nauchney, Trudy Belorusskogo Gosudarstvennogo Universiteta, Minsk. : 1-253 (Translated from Russian by Fish. Res. Bd. Canada, Transl. Ser., (194), 1960).

WINBERG, G. G. 1961. New information on metabolic rate in fishes. Vopt. ikhtiol., 1 : 157-165 (Translated from Russian by Fish. Res. Bd. Canada, Transl. Ser., 362).

WOODHEAD, P. M. J. 1966. The behaviour of fish in relation to light in the sea. Oceanogr. Mar. Biol. Ann. Rev., 4 : 337-404.

DESERVATE, S. D. AND T. M. SWARE 1969 On the IroH line lovestigations off Cochis during five fishing assessme. Pick. Tech., 6 (1),: 26-35.

DEVALUE M., 1983. Agt and growth of three sporter of seerflahes (Scomberroment conservates 5 guildar and 5 Joneolaturi (5) Indea / Adim J. Fath., 28 : 104-127.

DREEVEL S. M., F. V. S. NAMEOOTHIN AND V. G. GOWIGERAN, 1972, Results of wolling line operations in Gos waves during 1965-68. *Ibid.* 16 - 181-187 (for 1969).

HASTING, W. H. AND J., M. DSIGH 1972. Fred forensistion and excitation. In J. Fish Nutrition. Academic Press, New York and London. 327-374.

Iviaw, V. S. 1955. Experimental coology of the feeding of flates. Translated from the Russing by Denglas Scott. New Heven, Yale University Press, 1961 1-302.

KLEARAR, M. 1964. Observations on the food of juvenites of Scowberomorus commercion (Lacepade) and S. guanais (Bloch & Scheelder) from Vizhiajam; went coast of India. Mar. Biol. Ass. India, Proc. Symp. Scombroid Plates. Part-II: 586-590.

LEWES, A. D., B. R. SMITH AND R. E. KRADNEY. 1974. Studies on tunes and builfahi in Papus New, Guinea watera-II, Rea. Buill. DASP. Moresby (10) - 1-112.

MOORE, H. B. 1972. Aspects of stress in the trapical matrice environment, Adv. Mer. Biol. 19: 217-269

NATADUAR, A. V. AND A. G. BROUNL 1962. Index of preposiderators a method of grading the food observate in the stomach analysis of fishes. *Index J. Fish.*, \$ : 54-59 (for 1991)

Naustra, N. R. 1958, Jan the Inter competition browers, we appear of Corregonars in a thirth Swedish Julie. Jaur. Nauben: Res. Dreitingholm, Nept., 1991.

PALOHEDHO, J. E. AND L. M. DURIE 1965 FOCA and growth of fishes. 1: A growth carve derived from experimental data. J. Fish. Res. Bd. Canada, 22 1 521-542.

SRAN, E. G., 1967. Tuan fishery of the Timewelly coast, Oulf of Mannar, Mar. Blot. Ass. tudia, Proc. Symp. Scorebroid Fisher, Purt II: 1083-1117

Suppopersk L. B. 1961, Prelimenary Joza (or a production theory of ecology Amer. Nec. 95 (582) 145-153.

SREIVASA RAO, K. 1964. Observations on the flood and feeding tables of Scienderwowns gutanar (Bloch and Scinnester) and juveniles of S. fineolana. (Cavier and Valenciennes) and S. commersen (Lecepede) from the Walthi coast. Mar. Biol. Ans. India, Proc. Symp. Scienderaul Fishes. Part B: 591-598.

SUBRANOMA PRIAL N., R. S. MANDIARDOSS AND SULCCHARAN, P., 1972. Standardification of specifications for different tralling intes. Fish. Tech., 9 (1): 58-75.

UNDIFAC 1974, Survey MUBBA 1972/73, Pelagic Hiltery Project (IND 697393), Progress Haport, (6) : I-141.

VERGALAMAN, G. 1960. Studies on the food and, feeding relationship of the inshore fishes off Calicul on the Maisher coset. Indian J. Fish., 7 : 225-306