

Mussel Farming

Rajesh N, Suresh Babu P P and Bobby Ignatius

Mariculture Division, ICAR-CMFRI, Kochi

Mariculture, the farming of marine organisms in coastal and offshore waters, is an important component of sustainable aquaculture. Among mariculture practices, bivalve farming is gaining importance because it provides nutritious, affordable and eco-friendly animal protein. Bivalves such as mussels, oysters and clams are filter feeders and do not require artificial feed, making their culture economical and environmentally sustainable. They also support food security, coastal livelihoods, employment generation and water quality improvement through natural filtration.

Mussels are among the most commercially important bivalves cultured worldwide. Mussel farming has a history of more than 700 years, with early organized culture practices reported from France during the thirteenth century. Since then, farming methods have improved and expanded across temperate and tropical coastal regions. Mussels are preferred for culture because of their fast growth, high reproductive capacity, efficient use of natural food, good meat yield and strong consumer demand.

Distribution and Biology of Mussels

Globally, the European blue mussel (*Mytilus edulis*) is widely cultured in temperate regions, while the green mussel (*Perna viridis*) is important in tropical countries. Major mussel-producing countries include Spain, France, the Netherlands, Denmark, China, the Republic of Korea and New Zealand. In India, mussel farming is still developing, though the country has vast coastal resources suitable for expansion. The annual cultured mussel production is about 20,000 tonnes. The two major species cultured in India are the green mussel (*Perna viridis*) and the brown mussel (*Perna indica*). These species are valued for fast growth, high meat yield, adaptability and good market demand.

Mussel farming in India is mainly practiced along the southwest coast, especially in Kerala and Karnataka, where environmental conditions are favourable. The Central Marine Fisheries Research Institute (CMFRI) has played a major role in developing and promoting low-cost, eco-friendly and farmer-friendly mussel farming technologies. CMFRI has standardized rack, raft, longline and rope culture methods and promoted them through training, demonstrations and extension programmes. These efforts have improved farmer participation, household income and women's involvement through self-help groups.

Mussels are found along the Indian coastline and are known by local names such as Kallumekai, Kadukka and Chippi. They naturally inhabit rocky shores, estuaries,

backwaters and coastal waters where suitable substrates are available. They form dense natural beds in intertidal and subtidal zones, which support capture fisheries and provide seed for farming. Among Indian mussels, the green mussel is the most widely distributed and commercially important. It attaches to rocks, wooden structures, mangrove roots and other hard surfaces using strong byssal threads.

Kerala is one of the most productive regions for green mussels in India. Important mussel beds occur at Neendakara, Chaliyam, Narakkal, Koduvally, Thikkodi, Port Kollam, Chombala, Chellanam, Elathur, Moodadi, Mahe, Anchangadi, Ethai and Azhikkal. These areas favour spat settlement, fast growth and high survival. Along the Karnataka coast, important mussel beds occur at Gangoli, Malpe, Karwar, Someswara, Bhatkal, Byndur, Uchila, Gokarna, Coondapur, Suratkal and Dhareshwar. Green mussels are also reported from Goa, Maharashtra, Tamil Nadu, Andhra Pradesh and the Andaman and Nicobar Islands, though in lower densities.

The green mussel is a suspension-feeding marine bivalve. It filters seawater through the mantle cavity and traps food particles using cilia and mucus on the gills. Its diet includes phytoplankton, microalgae, small zooplankton, bacteria, detritus and suspended organic matter. Since it depends entirely on natural food, no supplementary feeding is required.

Perna viridis grows rapidly under favourable conditions, with shell length increasing by about 8–14 mm per month. In about five months, it can reach 80–90 mm shell length and 35–40 g body weight. Growth depends on temperature, salinity, dissolved oxygen, pH, plankton availability, water movement, stocking density and fouling. The species is dioecious, with separate males and females. Females have orange to reddish gonads, while males have creamy yellow or pale white gonads. Fertilization is external, and females release millions of eggs. Larvae pass through trochophore, veliger and pediveliger stages before settling as spat on hard substrates. Along the southwest coast, spat settlement usually occurs from July to September, and the juveniles reach seed size by September for use in farming.

Farm Site Selection for Mussel Culture

Farm site selection is a key factor in the success of mussel culture because growth, survival, meat quality, production and profitability depend on the environmental conditions of the culture area. A suitable site should have clean, well-oxygenated water free from sewage, industrial effluents, agricultural runoff, oil pollution and other contaminants, as mussels are filter feeders and can accumulate pollutants. The site should have abundant natural food such as plankton and suspended organic matter, with good primary productivity, suitable salinity of 27–35 ppt and temperature of 26–32°C. Moderate water movement is essential to supply food and oxygen and remove wastes, while the bottom should be firm with low siltation to prevent gill clogging. The area should

be protected from strong waves, cyclones and heavy swells, and should have access to roads, landing centres, labour, seed areas, markets and storage facilities. Mussel farming can be done in open sea and estuarine areas. Open sea sites require stable salinity, good water quality, moderate currents, limited wave action and a minimum depth of about 5 m; longline and raft systems are commonly used, though fouling, poaching, storms and rough seas are major risks. Estuarine sites are shallow, sheltered and easier to manage, with depths below 4 m preferred; rack and horizontal rope culture are commonly practiced. However, estuarine farming is affected by monsoon-related salinity reduction, pollution and sedimentation. Therefore, open sea sites are better for large-scale production, while estuarine sites are cheaper and suitable for small-scale farmers, making proper site selection essential for sustainable and profitable mussel farming.

Mussel Farming Methods

Mussel farming is a sustainable form of mariculture because mussels are filter feeders that depend on naturally available plankton and suspended organic matter. Therefore, no artificial feed is required, making the practice economical and environmentally friendly. The selection of a farming method depends on local conditions such as water depth, tidal range, current speed, bottom type, wave exposure, seed availability, investment capacity, and labour availability. Mussel farming methods are broadly classified into bottom culture and off-bottom or suspended culture. Suspended culture methods generally give better growth, higher production, and improved meat quality because mussels receive better water circulation and food supply.

1. Bottom Culture

Bottom culture involves spreading mussel seed directly on prepared seabed plots. It is commonly practiced in temperate countries where extensive intertidal flats and firm bottoms are available. Seed collected from natural mussel beds is transferred to culture grounds at suitable stocking densities. This method requires low investment and simple management. However, it is more affected by predators, sedimentation, poor water movement, and environmental disturbances.

2. Pole or Bouchot Culture

Pole culture is one of the oldest mussel farming methods and originated in France. In this system, wooden or bamboo poles are fixed vertically in intertidal areas. Mussel spat either settles naturally on the poles or is attached after collection. The mussels grow by attaching firmly to the poles with byssus threads. Protective netting may be used to prevent loss from predators and waves. This method is suitable for areas with a wide tidal range and moderate water movement.

3. Stake Culture

Stake culture is widely practiced in tropical countries, especially in Southeast Asia. Bamboo or wooden stakes are driven into soft-bottom coastal areas where natural spatfall occurs. Juvenile mussels settle on the submerged portions of the stakes and grow to marketable size. This method is inexpensive, simple, and suitable for sheltered bays and estuaries. However, continuous use may cause sediment accumulation and reduce water depth in the farming area.

4. Rack Culture

Rack culture is the most common method used for green mussel farming in India, particularly in shallow estuaries and backwaters. The system consists of bamboo or wooden poles supporting horizontal or vertical ropes on which mussels are grown. It is easy to construct, uses locally available materials, requires low capital investment, and allows simple maintenance and harvesting. CMFRI has standardized and promoted rack culture technology among coastal communities, especially in Kerala. Due to its low cost and good returns, it is suitable for small-scale farmers and women's self-help groups.

5. Raft Culture

Raft culture is practiced in sheltered coastal waters, bays, and lagoons with sufficient depth and good plankton availability. Floating rafts made of timber, steel, or HDPE pipes support vertical ropes called droppers, on which mussels are cultured. Good water circulation around the ropes provides food and oxygen, leading to fast growth and high meat yield. This method is suitable for commercial farming in protected waters.

6. Longline Culture

Longline culture is an efficient method for offshore mussel farming. It consists of buoy-supported horizontal ropes anchored to the seabed, with several vertical culture ropes suspended from them. Mussel seed is attached to the ropes using cotton or synthetic mesh, after which the mussels attach naturally. Longline systems allow good water exchange, high stocking density, and large-scale production. However, regular cleaning is needed to remove fouling organisms such as barnacles, algae, ascidians, and unwanted juvenile mussels.

Overall, rack and stake culture are suitable for shallow estuarine areas and small-scale farmers, while raft and longline culture are better for deeper coastal waters and commercial production. Bottom culture is simple and low-cost but more vulnerable to predators and sedimentation. Therefore, the farming method should be selected according to site conditions, available resources, and market demand.

Mussel farming in India has grown significantly during the last two decades, especially along the southwest coast. The development and demonstration of low-cost farming technologies by CMFRI have encouraged adoption by coastal farmers in Kerala, Karnataka, Goa, Tamil Nadu, and Maharashtra. Mussel farming has become an important

livelihood activity, providing additional income to fishing communities, women's self-help groups, and coastal entrepreneurs. With increasing domestic demand and export potential, the sector has good scope for expansion through improved seed production, value addition, cold-chain facilities, and post-harvest processing.

Successful mussel farming depends mainly on the availability of healthy seed and proper seeding techniques. Mussel seed or spat is collected from clean natural beds, preferably from subtidal rocky areas free from sewage, industrial waste, agricultural runoff, and other pollutants. Good-quality seed is usually 15–25 mm in shell length, as it withstands handling and attaches well after stocking. After collection, the seed is sorted to remove damaged, undersized mussels and fouling organisms such as barnacles, algae, tube worms, and ascidians. It is then washed with clean seawater before use.

Culture ropes made of coir or nylon are used for seeding because their rough surface helps mussels attach firmly with byssus threads. About 500–750 g of seed is required per metre of rope. The cleaned seed is spread evenly on a biodegradable cotton mesh, the rope is placed over it, and the mesh is wrapped and tied securely. Within 2–3 days, the mesh decomposes and the mussels attach naturally to the rope. Knots at 25 cm intervals prevent seed from slipping.

Seeded ropes are suspended from racks, rafts, or longlines. The upper end should remain submerged during low tide, while the lower end should stay above the seabed to avoid mud, sedimentation, and predators. Regular inspection, thinning, removal of dead mussels, and cleaning of fouling organisms are essential. Ropes should be spaced about 25 cm apart to ensure good water flow, food availability, uniform growth, high survival, and better meat yield.

Seeding, Rearing, Harvesting and Health Management in Mussel Farming

Successful mussel farming depends on healthy seed, proper seeding techniques, good farm management, safe harvesting, and disease prevention. Mussel seed or spat is usually collected from clean natural beds, preferably from subtidal rocky areas with minimum pollution. Collection sites should be free from sewage, industrial effluents, agricultural runoff, oil pollution, and other contaminants to ensure healthy seed and safe seafood production.

Good-quality mussel seed is generally 15–25 mm in shell length. Seed of this size can tolerate handling and transport and attaches well after stocking. After collection, the seed is sorted to remove damaged, undersized, or weak individuals, along with fouling organisms such as barnacles, algae, tube worms, and ascidians. The selected seed is then washed thoroughly with clean seawater to remove mud, silt, and debris.

Culture ropes made of coir or nylon are commonly used because their rough surface helps mussels attach firmly using byssus threads. About 500–750 g of seed is required

for one metre of rope. For seeding, cleaned mussel seed is spread evenly on a biodegradable cotton mesh, and the culture rope is placed along the centre. The mesh is wrapped around the rope and seed and tied securely. Within 2–3 days, the cotton mesh decomposes in seawater, while the mussels attach firmly to the rope. Knots are tied at intervals of about 25 cm to prevent seed from slipping and to maintain uniform distribution.

Seeded ropes are suspended vertically from racks, rafts, or longlines. The upper end should remain submerged even during low tide, while the lower end should stay above the seabed to avoid mud, sedimentation, and predators. After stocking, regular inspection is necessary to check attachment, mortality, fouling, and overcrowding. Thinning of dense clusters improves water circulation, reduces competition, and promotes uniform growth. Ropes should be spaced about 25 cm apart to ensure proper water flow and food availability.

Mussels generally grow faster in suspended culture than on the seabed because they receive better oxygen, plankton, and water circulation. The upper and middle portions of the rope often show faster growth than the lower portions due to better food availability and less sedimentation.

Harvesting is usually done after 5–7 months, when mussels reach marketable size. In India, harvesting is commonly completed before the southwest monsoon, mainly from April to June. Ropes are lifted manually from racks, rafts, or longlines, and mussels are detached, washed, sorted, and graded. Dead, damaged, and undersized mussels are removed. Fresh mussels should be protected from heat and sunlight and transported under cool, moist conditions.

Since mussels are filter feeders, depuration is important for food safety. In this process, live mussels are kept in clean, filtered, disinfected seawater for about 24 hours to remove contaminants, bacteria, sediments, and gut contents. Proper depuration improves hygiene, shelf life, and consumer acceptance.

Mussels are marketed mainly as live shell-on products, but value-added products such as frozen half-shell mussels, IQF meat, cooked mussels, pickles, smoked products, canned mussels, and ready-to-eat items offer better income opportunities. Good hygiene, cold-chain management, proper packaging, and food safety standards are essential for domestic and export markets.

Although mussel farming is low-risk, disease and mortality may occur due to poor seed, pollution, overcrowding, high temperature, salinity changes, low oxygen, and poor water exchange. Regular monitoring of growth, survival, water quality, fouling, and disease signs is essential. Healthy seed, proper stocking density, good farm layout, adequate spacing, removal of dead mussels, and basic biosecurity practices help maintain healthy stocks and sustainable production.

Integrated Multi-Trophic Aquaculture (IMTA)

Integrated Multi-Trophic Aquaculture (IMTA) is an environmentally sustainable aquaculture system in which different aquatic organisms belonging to different trophic levels are cultured together. It is designed to improve resource use, reduce waste, and increase farm productivity. Unlike intensive monoculture systems, which may cause nutrient enrichment, organic waste accumulation, poor water quality, disease problems, and habitat degradation, IMTA uses natural biological processes to recycle nutrients within the farming system.

The basic principle of IMTA is that the waste produced by one cultured species becomes a useful resource for another. In conventional aquaculture, uneaten feed, faeces, and metabolic wastes from fed species such as marine finfish may enter the environment and cause pollution. In IMTA, these wastes are utilized by extractive species such as bivalves and seaweeds. This reduces environmental impact while producing additional crops of economic value.

A typical marine IMTA system includes three major components: fed species, organic extractive species, and inorganic extractive species. Fed species such as seabass, cobia, pompano, groupers, or other marine finfish are usually cultured in cages or pens and receive formulated feed. They act as the main source of nutrients in the system.

Organic extractive species include filter-feeding bivalves such as mussels, oysters, and clams. These organisms remove suspended organic particles, phytoplankton, and fine particulate wastes from the water. By filtering the water column, they reduce organic pollution and improve water clarity while producing a valuable shellfish crop.

Inorganic extractive species mainly include seaweeds. Seaweeds absorb dissolved nutrients such as ammonia, nitrate, phosphate, and carbon dioxide released from fish culture. Through photosynthesis, they also add oxygen to the water. In addition to improving water quality, seaweeds are useful raw materials for food, fertilizers, pharmaceuticals, cosmetics, and biofuel industries.

The success of IMTA depends on proper selection of species. Cultured organisms should be native or well adapted to local environmental conditions such as temperature, salinity, water depth, and current pattern. They should occupy complementary trophic levels so that nutrients are efficiently transferred from one group to another. Selected species should also have fast growth, good survival, disease resistance, compatible production cycles, and strong market demand.

IMTA provides several environmental benefits. It reduces nutrient loading, eutrophication, organic enrichment, and water quality deterioration. Bivalves remove suspended particles, while seaweeds absorb dissolved nutrients, thereby improving ecosystem health. The system also increases nutrient-use efficiency and reduces the

environmental footprint of aquaculture. Seaweed cultivation can also contribute to carbon sequestration, making IMTA relevant in the context of climate change.

Economically, IMTA allows farmers to harvest more than one product from the same farming area. Finfish, shellfish, and seaweeds can be produced together, reducing dependence on a single crop and lowering financial risk. Since bivalves and seaweeds require little or no external feed, operational costs are reduced. IMTA also creates employment opportunities in seed production, farming, harvesting, processing, value addition, and marketing.

In India, IMTA has good potential for sustainable expansion of marine aquaculture. With proper site selection, suitable species combinations, nutrient modelling, scientific management, and policy support, IMTA can improve farm income while protecting coastal ecosystems.

Thus, IMTA is a holistic and eco-friendly aquaculture approach that converts waste into useful biomass. By integrating fed fish with bivalves and seaweeds, it improves water quality, increases productivity, supports livelihood generation, and promotes sustainable development of mariculture.
